

The Patient Protection and Affordable Care Act (PPACA) and the Health Care and Education Reconciliation Act of 2010 (HCERA) has designated the services listed below as preventive benefits and available with no cost-sharing when provided by an in-network provider for members of non-grandfathered health plans. In addition to the services listed below, your patient may have additional preventive care benefits covered under their health plan that may or may not be covered at 100%. Your patient should check their benefit booklet for details on these additional preventive care benefits. The following tables provide a quick reference guide for submitting claims for preventive services with a "well-person" diagnosis code as the primary (first) diagnosis on the claim. This information is intended as a reference tool for your convenience and is not a guarantee of payment. This guide is subject to change based on new or revised laws and/or regulations, additional guidance and/or BCBSNC medical policy.

IMPORTANT INFORMATION: Services must be billed with the appropriate diagnosis, at the line level of the claim (Block 24E), pursuant to industry standard coding guidelines. Preventive or screening services are intended for those who currently exhibit no signs or symptoms of disease. Services otherwise deemed preventive that are received in an inpatient setting, an emergency room, or that include additional procedures or diagnostic services may be subject to copayment, deductible and coinsurance. Submitting screening service codes (CPT, HCPCs, ICD-9 or ICD-10) when signs or symptoms are present constitutes inappropriate coding which could result in recoupment of monies paid to the provider for those services. Additionally, these services are subject to certain limitations depending on medical necessity and other reasonable medical management techniques. If you have questions, please contact the **Provider Blue Line at 1-800-214-4844**.

Grade A and B			
Recommendations of U.S.			
Preventive Services Task			
Force (USPSTF) currently			
effective unless otherwise			
noted	CPT or HCPCS	ICD-10 Diagnosis	Comments
Screening for Abdominal Aortic	76706	Z13.6	
Aneurysm			
(one time screening for abdominal aortic aneurysm by ultrasonography in men ages			
65-75 who have ever smoked)			
Screening and counseling to reduce	99408, 99409,	Z13.89, F10.10, F10.120, F10.129	These codes are to be used in the absence of a wellness
unhealthy alcohol use The USPSTF	G0442 or G0443,		visit.
recommends that clinicians screen adults aged 18 years or older, including pregnant	G0396, G0397 G2011		
women for unhealthy alcohol use and	02011		
provide persons engaged in risky or			
hazardous drinking with brief behavioral counseling interventions to reduce alcohol			
misuse.			
Aspirin to prevent cardiovascular disease	Not applicable,	Not applicable, administered through Pharmacy	OTC Aspirin (81 mg) is dispensed to member with a
and colorectal cancer in men and women (Low-dose aspirin use for the primary	administered through		physician order with no cost sharing.
prevention of cardiovascular disease (CVD)	Pharmacy		
and colorectal cancer (CRC) in adults aged			
50 to 59 years who have a 10% or greater			
10 year CVD risk, are not at increased risk for bleeding, have a life expectancy of at			
least 10 years, and are willing to take low-			
dose aspirin daily for at least 10 years.)			
Anxiety screening (screening for anxiety in adult women and female adolescents,	Part of preventive visit or 99401,	Z00.00, Z00.01, Z01.411, Z01.419	Annual 99401-99404 are to be used in the absence of a wellness
including those who are pregnant or	99402, 99403,		visit.
postpartum)	99404		
Screening for bacteriuria (screening for	87081, 87084,	Z34.00, Z34.01, Z34.02, Z34.03, Z34.80, Z34.81, Z34.82,	Dipstick or other urinalysis included in global maternity
asymptomatic bacteriuria with <u>urine culture</u> for pregnant women at 12 - 16 weeks'	87086 or 87088	Z34.83, Z34.90, Z34.91, Z34.92, Z34.93, Z33.1, O09.00, O09.01, O09.02, O09.03, O09.10, O09.11, O09.12,	visits. Only urine culture part of preventive services mandate.
gestation or at the first prenatal visit, if later)		009.13, 009.291, 009.292,009.293, 009.299, 009.40,	NOTE: Z33.1 is not an acceptable principle
, ,		009.41, 009.42, 009.43, 009.211, 009.212,009.213,	diagnosis code.
		009.219, 009.30, 009.31, 009.32, 009.33, 009.511,	
		O09.512,O09.513, O09.519, O09.521 O09.522,O09.523, O09.529,O09.611, O09.612,O09.613, O09.619,O09.621,	
		009.622,009.623, 009.629,009.811, 009.812,009.813,	
		O09.819,09.821,O09.822,O09.823,O09.829,O36.80X0,O3	
		6.80X1,O36.80X2, O36.80X3,O36.80X4, O36.80X5,O36.80X9, O09.891, O09.892, O09.893,	
		O09.899, O09.70, O09.71, O09.72, O09.73, O09.90,	
		O09.91, O09.92, O09.93	
Corponing for high blood processes	00205 00207	009.A0, 009.A1, 009.A2, 009.A3	Dort of wallness office winit
Screening for high blood pressure (Recommends screening for high blood	99385-99387 or 99395-99397	Z13.6 or other wellness exam diagnosis code For coverage of ABPM or HBPM diagnosis code R03.0 is	Part of wellness office visit. Effective 1/1/2017, ambulatory blood pressure
pressure in adults aged 18 years or older.	For ambulatory	required	monitoring(ABPM) or home blood pressure monitors
The USPSTF recommends obtaining	blood pressure		(HBPM) for confirmation of high blood pressure is
measurements outside of the clinical setting for diagnostic confirmation before starting	monitoring use 93784, 93786,		covered at 100%. Home blood pressure monitors must be purchased from an in-network durable medical
treatment)	93788, 93790		equipment (DME) vendor.
	For home blood		- d- h///
	pressure monitor		
	use A4670		



Grade A and B			
Recommendations of U.S.			
Preventive Services Task			
Force (USPSTF) currently			
effective unless otherwise			
noted	CPT or HCPCS	ICD-10 Diagnosis	Comments
Counseling related to BRCA screening	Referrals for	Z80.3, Z80.41, Z85.07, Z85.3, Z85.43, Z85.44, Z85.46, or	Services for BRCA 1/BRCA 2 testing will be provided
(Primary care clinicians should assess women with a personal or family history of	counseling included in wellness visit	Z71.83	with no cost sharing to appropriate groups when the medical policy criteria are met.
breast, ovarian, tubal, or peritoneal cancer	codes 99385-99387,		See BCBSNC Medical Policy: Genetic Testing for Breast
or who have an ancestry associated with	99395-99397		and Ovarian Cancer
breast cancer susceptibility 1 and 2	Counseling services		99401-99404 are to be used in the absence of a wellness
(BRCA1/2) gene mutations with an appropriate brief familial risk assessment	use 96040, 99401, 99402, 99403,		visit.
tool. Women with positive screening results	99404, or S0265		
should receive generic counseling and, if	Genetic testing use		
indicated after counseling, BRCA testing.)	81212, 81215,		
	81216, 81217 or 81162		
	81163, 81164,		
	81165,81166, 81167		
Screening for breast cancer	77067, 77063	Z12.31, Z12.39	If the patient has had an abnormal mammogram in the
[mammography]			past, subsequent routine mammograms may be coded
(for women aged 40 or over every 1-2 yrs, with or without clinical breast examination)			as diagnostic: 77061, 77062, 77065, 77066, G0279 with a diagnosis reflective of the abnormality.
Chemoprevention of breast cancer	99401-99402 or is	Z15.01, Z80.3, D24.1, D24.2, D24.9, N60.81, N60.82,	This recommendation applies to asymptomatic women
(Recommends that clinicians engage in	included in wellness	N60.89	aged 35 years or older without a prior diagnosis of breast
shared, informed decision making with	visit 99385-99387,		cancer, ductal carcinoma in situ, or lobular carcinoma in
women who are at increased risk for breast	99395-99397		situ. Generic risk-reducing medications are dispensed to
cancer about medications to reduce their risk. For women who are at increased risk			member with a physician order with no cost-sharing. For members who have swallowing problems or may
for breast cancer and at low risk for adverse			have an intolerance to generic products, brand products
medication effects, clinicians should offer to			may also be made available by completing the Copay
prescribe risk-reducing medications, such as			Waiver Form and faxing it to the number on the bottom
tamoxifen or raloxifene)			of the document. 99401-99402 are to be used in the absence of a wellness
			visit.
			NOTE: Z15.01 is not an acceptable principle diagnosis
Interventions to support breast feeding	99401-99403 or part	O30.93, Z34.03, Z34.83, Z34.93, Z39.1	99401-99403 are to be used in the absence of a wellness
(interventions during pregnancy and after	of other office visit		visit.
birth to promote and support breastfeeding)			
Screening for cervical cancer	88141, 88142,	Z00.00, Z00.01, Z01.411, Z01.419, Z01.42, Z12.4, Z11.51	This recommendation does not apply to women who
(In women age 21–2965 with cervical	88143, 88147,		have received a diagnosis of a high-grade precancerous
cytology (pap testsmear) every 3 years For women age 30–65,	88148, 88150, 88152, 88153,		cervical lesion or cervical cancer, women with in utero exposure to diethylstilbestrol, or women who are
A Pap test every 3 years or A Pap test and	88164, 88165,		immunocompromised (such as those who are HIV
high-risk human papillomavirus (hrHPV) test	88166, 88167,		positive).
every 5 years or HPV test only every 5	88174, 88175,		
years)	87623, 87624, 87625, G0101,		
	G0123, G0101, G0123, G0124,		
	G0141, G0143,		
	G0144, G0145,		
	G0147, G0148, P3000, P3001		
	1 3000, F3001		
Screening for chlamydial infection in	87270, 87320,	Z34.00, Z34.01, Z34.02, Z34.03, Z34.80, Z34.81, Z34.82,	NOTE: Z33.1 is not an acceptable principle diagnosis
non-pregnant and pregnant women (for	87490, 87491 and	Z34.83, Z34.90, Z34.91, Z34.92, Z34.93, Z33.1, O09.00,	code.
all sexually active non-pregnant and	87810	O09.01, O09.02, O09.03, O09.10, O09.11, O09.12, O09.13, O09.291, O09.292, O09.293, O09.299, O09.40,	
pregnant young women ages 24 and younger and for older, non-pregnant and		O09.13, O09.291, O09.292, O09.293, O09.299, O09.40, O09.41, O09.42, O09.43, O09.211, O09.212, O09.213,	
pregnant women who are at increased risk)		O09.219, O09.30, O09.31, O09.32, O09.33, O09.511,	
,		O09.512, O09.513, O09.519, O09.521 O09.522, O09.523,	
		O09.529, O09.611, O09.612, O09.613, O09.619,	
		O09.621, O09.622, O09.623, O09.629, O09.811, O09.812, O09.813, O09.819, O09.821, O09.822,	
		O09.812, O09.813, O09.819, O09.821, O09.822, O09.823, O09.829, O36.80X0, O36.80X1, O36.80X2,	
		O36.80X3, O36.80X4, O36.80X5, O36.80X9, O09.891,	
		009.892, 009.893, 009.899, 009.70, 009.71, 009.72,	
		O09.73, O09.90, O09.91, O09.92, O09.93, Z11.8, O09.A0, O09.A1, O09.A2, O09.A3, Z11.3	
		000.nu, 000.n1, 000.nz, 000.nd, 211.0	



Grade A and B			
Recommendations of U.S.			
Preventive Services Task			
Force (USPSTF) currently			
effective unless otherwise			
noted	CPT or HCPCS	ICD-10 Diagnosis	Comments
Screening for cholesterol abnormalities:	80061, 82465 or	Z13.220	Do not bill the panel lab code in addition to separate tests
*men 35 and older (for lipid disorders) *men younger than 35 (for ages 20-35 for lipid disorders if they are at increased risk for coronary heart disease) *women 20 and older (for lipid disorders if they are at increased risk for coronary heart disease) See also: Statin Use for the Primary Prevention of Cardiovascular Disease in Adults	83718	210.220	included in the panel.
Prevention of dental caries	99188, 99401 or	Z29.3	Fluoride varnish will not be covered beyond the 5th
(recommends that primary care clinicians prescribe fluoride supplementation starting at age 6 months for children whose water supply is deficient in fluoride. Also recommends that primary care clinicians apply fluoride varnish to the primary teeth of all infants and children starting at the age of primary tooth eruption.)	99402 May prescribe oral fluoride during child wellness visit: 99381-99382 or 99391-99392 with appropriate wellness diagnosis		birthday. Oral fluoride supplementation will be covered from ages 6 months through 16 years. BCBSNC recommends counseling/education in addition to varnish application. • For well child visit: Any counseling services received at the time of a well child visit are considered part of the preventive visit and are not reimbursed separately. • For sick visit: If the varnish is applied during a visit other than for wellness, providers should reference diagnosis code Z41.8 or Z29.3 as the primary diagnosis for the line item of the varnish application service, 99188, on the claim in order to process correctly. If education/counseling on prevention and risk factor reduction of dental caries is conducted during a sick visit, providers should again reference diagnosis code Z41.8 or Z29.3 as the primary diagnosis for the line item of the counseling service, 99401/99402, on the claim in order to process correctly. BCBSNC will disallow D codes if submitted by non licensed dental providers 99401-99402 are to be used in the absence of a wellness visit.
Screening for colorectal cancer (using	Fecal occult blood	Screening Procedures:	VISIT. Colorectal cancer screening mandate does not include
fecal occult blood testing, sigmoidoscopy, or colonoscopy, in adults beginning at ages 45-75) Age change effective for eligible members by April 1, 2022	testing: 82270, 82274 or G0328 FIT DNA: 81528 Sigmoidoscopy: 45330, 45333, 45334, 45338, G0104, G6022, 88305 and Effective 1/1/2017 G0500 Colonoscopy: 45378, 45380, 45381, 45382, 45384, 45385, 74263, G0105, G0121, G6024, 00812, 88305 and G0500 Effective 1/1/2023 45390 Consultation or office visit prior to screening colonoscopy: S0285	Z12.11, Z12.12, K51.40, K62.1, K63.5, D12.0, D12.2, D12.3, D12.4, D12.5, D12.6, D12.7, D12.8, or D12.9 Pathology services (88305): Z12.11, Z12.12, K51.40, K62.1, K63.5, D12.0, D12.2, D12.3, D12.4, D12.5, D12.6, D12.7, D12.8, or D12.9 Visit or Consultations: Z01.818	barium enema. ***If a test is performed for screening purposes, screening service codes may be submitted even in the event of positive findings e.g., polyp found during a screening colonoscopy. It is critical to use a screening diagnosis as the primary diagnosis for the claim line of the test performed [not just in the header diagnoses]. Any abnormal findings from a screening test should be listed as a secondary diagnosis code. Failure to list screening as the primary diagnosis code will affect correct claim adjudication. Effective 6/1/2022: If a colonoscopy is performed following a positive stool based (FIT, Cologuard) or direct visualization test (sigmoidoscopy or CT colonography), it is critical to use a screening diagnosis as the primary diagnosis for the claim line of the test performed [not just in the header diagnoses]. Any abnormal findings from a screening test should be listed as a secondary diagnosis code. Failure to list screening as the primary diagnosis code will affect correct claim adjudication. As of 9/1/2016, Certain bowel preparation medications, when medically appropriate and prescribed by a health care provider are allowed without cost sharing when dispensed at a participating pharmacy. If there is a medical reason a member cannot take a generic bowel preparation, the physician should review the clinical criteria and if appropriate, submit fax form. As of 1/1/2017, Moderate sedation (G0500) performed by the physician performing the procedure is allowed but cannot be billed on the same DOS as 00812.



Grade A and B Recommendations of U.S. Preventive Services Task Force (USPSTF) currently			
effective unless otherwise			
noted	CPT or HCPCS	ICD-10 Diagnosis	Comments
Screening for depression: *adults (screening for depression, including pregnant and postpartum women, when staff-assisted depression care supports are in place to assure accurate diagnosis, effective treatment, and follow-up) *adolescents (screening 12-18 yr olds for major depressive disorder when systems are in place to ensure accurate diagnosis, psychotherapy [cognitive-behavioral or interpersonal], and follow-up)	G0444 or preventive visit	Z13.89 Remove effective 9/30/18 Effective 10/1/18: Z13.31	Part of any problem or preventive office visit.
Screening for diabetes Screening for abnormal blood glucose as part of cardiovascular risk assessment. Clinicians should offer or refer patients with abnormal blood glucose to intensive behavioral counseling interventions to promote a healthful diet and physical activity. Adults aged 40 to 70 who are overweight or obese	82947 or 83036	Z13.1,	
Behavioral Counseling to promote a healthful diet and physical activity for cardiovascular disease prevention in adults with cardiovascular risk factors (Recommends offering or referring adults with cardiovascular disease (CVD) risk factors behavioral counseling interventions to promote a healthy diet and physical activity)	99402, 99403, 99404, 97802, 97803, 97804, S9452, S9470, G0270, G0271, G0108, G0109, S9140, S9141, S9455, S9460, S9465, G0446 or part of preventive visit	Z71.3	99402-99404 are to be used in the absence of a wellness visit.
Fall Prevention (Exercise interventions to prevent falls in community-dwelling adults 65 years or older who are at increased risk for falls.)	97001-97002, 97110, 97112, 97116, 97530, G0151, G0157, G0159, S9131 and S9476.	Z91.81	Vitamin D is dispensed to member with a physician order with no cost-sharing.
Supplementation with folic acid (for all women planning or capable of pregnancy to take a daily supplement containing 0.4 - 0.8 mg [400-600 mcg] of folic acid)	Not applicable, administered through Pharmacy	Not applicable, administered through Pharmacy	OTC folic acid supplements are dispensed to member with a physician order with no cost-sharing.
Corponing for Contational District	82950	Z34.00, Z34.01, Z34.02, Z34.03, Z34.80, Z34.81, Z34.82,	In pregnant women between 24 and 28 weeks of
Screening for Gestational Diabetes Mellitus (Recommends screening for gestational diabetes mellitus (GDM) in asymptomatic pregnant women after 24 weeks of gestation.)		Z34.83, Z34.90, Z34.91, Z34.92, Z34.93, Z33.1, O09.00, O09.01, O09.02, O09.03, O09.10, O09.11, O09.12, O09.13, O09.291, O09.292, O09.293, O09.299, O09.40, O09.41, O09.42, O09.43, O09.211, O09.212, O09.213, O09.219, O09.30, O09.31, O09.32, O09.33, O09.511, O09.512, O09.513, O09.519, O09.521 O09.522, O09.523, O09.529, O09.611, O09.612, O09.613, O09.619, O09.621, O09.622, O09.623, O09.629, O09.811, O09.812, O09.813, O09.819, O09.821, O09.822, O09.823, O09.829, O36.80X0, O36.80X1, O36.80X2, O36.80X3, O36.80X4, O36.80X5, O36.80X9, O09.891, O09.892, O09.893, O09.899, O09.70, O09.71, O09.72, O09.73, O09.90, O09.91, O09.92, O09.93 O09.A0, O09.A1, O09.A2, O09.A3	gestation and at the first prenatal visit for pregnant women identified to be at high risk for diabetes NOTE: Z33.1 is not an acceptable principle diagnosis code.
Screening for gonorrhea for women (all sexually active women, including those who are pregnant if they are at increased risk for infection [if they are young or have other individual or population risk factors])	87590-87591 and 87850	Z11.59, Z11.3, Z34.00, Z34.01, Z34.02, Z34.03, Z34.80, Z34.81, Z34.82, Z34.83, Z34.90, Z34.91, Z34.92, Z34.93, Z33.1, O09.00, O09.01, O09.02, O09.03, O09.10, O09.11, O09.12, O09.13, O09.291, O09.292, O09.293, O09.299, O09.40, O09.41, O09.42, O09.43, O09.211, O09.212, O09.213, O09.219, O09.30, O09.31, O09.32, O09.33, O09.511, O09.512, O09.513, O09.519, O09.521 O09.522, O09.523, O09.529, O09.611, O09.612, O09.613, O09.619, O09.621, O09.622, O09.623, O09.629, O09.811, O09.812, O09.813, O09.819, O09.821, O09.822, O09.823, O09.829, O36.80X0, O36.80X1, O36.80X2, O36.80X3, O36.80X4, O36.80X5, O36.80X9, O09.891, O09.892, O09.893, O09.899, O09.70, O09.71, O09.72, O09.73, O09.90, O09.91, O09.92, O09.93, Z11.8 O09.A0, O09.A1, O09.A2, O09.A3	NOTE: Z33.1 is not an acceptable principle diagnosis code.



Grade A and B Recommendations of U.S. Preventive Services Task Force (USPSTF) currently effective unless otherwise noted Prophylactic medication for gonorrhea: newborns (ocular topical medication for all newborns against gonococcal ophthalmia neonatorum)	CPT or HCPCS Not applicable, administered through Facility	ICD-10 Diagnosis Not applicable, administered through Facility	Comments This medication is generally administered to newborn at birth facility.
Healthy Weight and Weight Gain In Pregnancy: Behavioral Counseling Interventions (For pregnant adolescents and adults: Offer effective behavioral counseling interventions aimed at promoting healthy weight gain and preventing excess gestational weight gain in pregnancy.)	99402, 99403, 99404, 97802, 97803, 97804, S9452, S9470, G0270, G0271or part of preventive visit	Z34.00, Z34.01, Z34.02, Z34.03, Z34.80, Z34.81, Z34.82, Z34.83, Z34.90, Z34.91, Z34.92, Z34.93, Z33.1, O09.00, O09.01, O09.02, O09.03, O09.10, O09.11, O09.12, O09.13, O09.291, O09.292, O09.293, O09.299, O09.40, O09.41, O09.42, O09.43, O09.211, O09.212, O09.213, O09.219, O09.30, O09.31, O09.32, O09.33, O09.511, O09.512, O09.513, O09.519, O09.521 O09.522, O09.523, O09.529, O09.611, O09.612, O09.613, O09.619, O09.621, O09.622, O09.623, O09.629, O09.811, O09.812, O09.813, O09.819, O09.821, O09.822, O09.823, O09.829, O36.80X0, O36.80X1, O36.80X2, O36.80X3, O36.80X4, O36.80X5, O36.80X9, O09.891, O09.892, O09.893, O09.899, O09.70, O09.71, O09.72, O09.73, O09.90, O09.91, O09.92, O09.93, O09.A0, O09.A1, O09.A2, O09.A3, Z71.3	NOTE: Z33.1 is not an acceptable principle diagnosis code.
Screening for hearing loss (in all newborn infants)	92558, 99381, 99391 or 92586	Z00.121, Z00.129, Z00.110, Z00.111, Z01.10, Z01.118	Service is typically performed in the birth facility or as part of a wellness office visit in the event of a home birth. CPT code 92586 cannot be billed by professional provider in a facility setting
Screening for hemoglobinopathies (for sickle cell disease in newborns)	83020, 83021, 85660 and S3620	Z13.0	Service is typically performed in the birth facility or as part of a wellness office visit in the event of a home birth.
Screening for Hepatitis B (in pregnant women at first prenatal visit and in persons at high risk for infection.)	80055 , 87340 or 80081	Z34.00, Z34.01, Z34.02, Z34.03, Z34.80, Z34.81, Z34.82, Z34.83, Z34.90, Z34.91, Z34.92, Z34.93, Z33.1, O09.00, O09.01, O09.02, O09.03, O09.10, O09.11, O09.12, O09.13, O09.291, O09.292, O09.293, O09.299, O09.213, O09.219, O09.30, O09.31, O09.32, O09.33, O09.511, O09.512, O09.513, O09.519, O09.521 O09.522, O09.523, O09.529, O09.611, O09.612, O09.613, O09.619, O09.621, O09.622, O09.623, O09.629, O09.811, O09.812, O09.813, O09.819, O09.821, O09.822, O09.823, O09.829, O36.80X0, O36.80X1, O36.80X2, O36.80X3, O36.80X4, O36.80X5, O36.80X9, O09.891, O09.892, O09.893, O09.899, O09.70, O09.71, O09.72, O09.73, O09.90, O09.91, O09.92, O09.93, Z11.59 O09.A0, O09.A1, O09.A2, O09.A3	NOTE: Z33.1 is not an acceptable principle diagnosis code.
Screening for hepatitis C virus (HCV) infection (for persons at high risk for infection.) The USPSTF also recommends offering one-time screening for HCV infection to adults aged 18 to 79	G0472	Z72.51, Z72.52, Z72.53, Z11.59, Z11.4, F10.10, F10.20, F10.21, F10.229, F11.10, F11.20, F11.21, F12.10, F12.20, F12.21, F13.10, F13.20, F13.21, F14.10, F14.20, F14.21, F15.10, F15.129, F15.20, F15.21, F16.10, F16.129, F16.20, F16.21, F17.200, F18.10, F19.10, F19.20, F19.21, F55.2	
Screening for HIV (on all adolescents ages 15 to 65 years, younger adolescents and older adults who are at increased risk, and all pregnant women including those who present in labor who are untested and whose HIV status is unknown)	G0432, G0433, G0435, OR G0475	Z34.00, Z34.01, Z34.02, Z34.03, Z34.80, Z34.81, Z34.82, Z34.83, Z34.90, Z34.91, Z34.92, Z34.93, Z33.1, O09.00, O09.01, O09.02, O09.03, O09.10, O09.11, O09.12, O09.13, O09.291, O09.292, O09.293, O09.299, O09.40, O09.41, O09.42, O09.43, O09.211, O09.212, O09.213, O09.219, O09.30, O09.31, O09.32, O09.33, O09.511, O09.512, O09.513, O09.519, O09.521 O09.522, O09.523, O09.529, O09.611, O09.612, O09.613, O09.619, O09.621, O09.622, O09.623, O09.629, O09.811, O09.812, O09.813, O09.819, O09.821, O09.822, O09.823, O09.829, O36.80X0, O36.80X1, O36.80X2, O36.80X3, O36.80X4, O36.80X5, O36.80X9, O09.891, O09.892, O09.893, O09.899, O09.70, O09.71, O09.72, O09.73, O09.90, O09.91, O09.92, O09.93, Z11.4 O09.A0, O09.A1, O09.A2, O09.A3	NOTE: Z33.1 is not an acceptable principle diagnosis code.
Screening for congenital hypothyroidism (in newborns)	99381 or 99391	Z13.29	Service is typically performed in the birth facility or as part of a wellness office visit in the event of a home birth.



Grade A and B			
Recommendations of U.S.			
Preventive Services Task			
Force (USPSTF) currently			
effective unless otherwise			
noted	CPT or HCPCS	ICD-10 Diagnosis	Comments
Screening for iron deficiency anemia (in asymptomatic pregnant women)	80055, 85013, 85014, 85018, 85025, 85027 or 80081	Z34.00, Z34.01, Z34.02, Z34.03, Z34.80, Z34.81, Z34.82, Z34.83, Z34.90, Z34.91, Z34.92, Z34.93, Z33.1, O09.00, O09.01, O09.02, O09.03, O09.10, O09.11, O09.12, O09.13, O09.291, O09.292, O09.293, O09.299, O09.40, O09.41, O09.42, O09.43, O09.211, O09.212, O09.213, O09.219, O09.30, O09.31, O09.32, O09.33, O09.511, O09.512, O09.513, O09.519, O09.521 O09.522, O09.523, O09.529, O09.611, O09.612, O09.613, O09.619, O09.621, O09.622, O09.623, O09.629, O09.811, O09.812, O09.813, O09.819, O09.821, O09.822, O09.823, O09.829, O36.80X0, O36.80X1, O36.80X2, O36.80X3, O36.80X4, O36.80X5, O36.80X9, O09.891, O09.892, O09.893, O09.899, O09.70, O09.71, O09.72, O09.73, O09.90, O09.91, O09.92, O09.93 O09.A0, O09.A1, O09.A2, O09.A3	NOTE: Z33.1 is not an acceptable principle diagnosis code.
Iron supplementation in children (routine iron supplementation for asymptomatic children 6-12 mo of age who are at increased risk for iron deficiency anemia)	Not applicable, administered through Pharmacy	Not applicable, administered through Pharmacy	OTC iron supplements are dispensed to member with a physician order with no cost-sharing.
Screening for latent tuberculosis (asymptomatic adults 18 years and older at	86480, 86481, 86580	Z11.1	Mandate effective 10/1/2017
increased risk for tuberculosis)			
Low-Dose Aspirin Use for the Prevention of Morbidity and Mortality From Preeclampsia (Recommends the use of low-dose aspirin (81 mg/d) as preventive medication after 12 weeks of gestation in women who are at high risk for preeclampsia.)	Not applicable, administered through Pharmacy	Not applicable, administered through Pharmacy	OTC Aspirin (81 mg) is dispensed to member with a physician order at a participating pharmacy with no cost-sharing.
Screening for Lung Cancer (Recommends annual screening for lung cancer with low-dose computed tomography (LDCT) in adults aged 50 to 80 yrs who have a 20pack-year smoking history and currently smoke or have quite within the past 15 years. Screening should be discontinued once a person has not smoked for 15 years or develops a health problem that substantially limits life expectancy or the ability or willingness to have curative lung surgery.)	G0296, G0297 (termed 12/31/2020) 71271 (effective 1/1/2021)	F17.210, F17.211, F17.213, F17.218, F17.219, Z72.0, and Z87.891	
Screening for and Management of Obesity in Adults (screen all adults; clinicians should offer or refer patients with a body mass index (BMI) of 30 kg/m2 or higher to intensive, multi-component behavioral interventions.)	99385, 99386, 99387, 99395, 99396, 99397, 96150, 96151, 96152, 96153, 99401, 99402, 99403, 99404, G0447 or G0473	Z00.00, Z00.01, Z13.89, Z68.30, Z68.31, Z68.32, Z68.33, Z68.34, Z68.35, Z68.36, Z68.37, Z68.38, Z68.39, Z68.41, Z68.42, Z68.43, Z68.44, Z68.45, E66.9, E66.01, E66.09, E66.1, E66.8, Z13.220, Z13.228, Z13.29	99401-99404 are to be used in the absence of a wellness visit. NOTE: Z68.30, Z68.31, Z68.32, Z68.33, Z68.34, Z68.35, Z68.36, Z68.37, Z68.38, Z68.39, Z68.41, Z68.42, Z68.43, Z68.44, Z68.45 are not acceptable principle diagnosis codes.
Screening and counseling for obesity: children (screen children aged 6 yrs and older for obesity and offer/refer to comprehensive, intensive behavioral interventions to promote improvement in weight status)	99383-99384, 99393, 99394, 99401, 99402, 99403, 99404, G0447 or G0473	Z00.121, Z00.129, Z00.110, Z00.111, Z13.89, Z13.220, Z13.228, Z13.29	99401-99404 are to be used in the absence of a wellness visit.
Screening for osteoporosis (screen women aged 65 and older for osteoporosis and in younger postmenopausal women who are at increased risk of osteoporosis, as determined by a formal clinical risk assessment tool.)	77080 or 77081	Z13.820	This recommendation uses only DEXA scans for testing.
Screening for PKU (in newborns)	84030	Z13.228	Service is typically performed in the birth facility or as part of a wellness office visit in the event of a home birth. 84030 is to be used in the absence of a panel which includes this test.



Grade A and B Recommendations of U.S. Preventive Services Task Force (USPSTF) currently effective unless otherwise noted Screening for preeclampsia	CPT or HCPCS Performed as part of routine prenatal visit	ICD-10 Diagnosis Z34.00, Z34.01, Z34.02, Z34.03, Z34.80, Z34.81, Z34.82, Z34.83, Z34.90, Z34.91, Z34.92, Z34.93, Z33.1, O09.00,	Comments Effective 5/1/18 or upon renewal Service is performed as part of routine prenatal visit not
		O09.01, O09.02, O09.03, O09.10, O09.11, O09.12, O09.13, O09.291, O09.292, O09.293, O09.299, O09.40, O09.41, O09.42, O09.43, O09.211, O09.212, O09.213, O09.219, O09.30, O09.31, O09.32, O09.33, O09.511, O09.512, O09.513, O09.519, O09.521 O09.522, O09.523, O09.529, O09.611, O09.612, O09.613, O09.619, O09.621, O09.622, O09.623, O09.629, O09.811, O09.812, O09.813, O09.819, O09.821, O09.822, O09.823, O09.829, O36.80X0, O36.80X1, O36.80X2, O36.80X3, O36.80X4, O36.80X5, O36.80X9, O09.891, O09.892, O09.893, O09.899, O09.70, O09.71, O09.72, O09.73, O09.90, O09.91, O09.92, O09.93 O09.A0, O09.A1, O09.A2, O09.A3	separately reimbursable NOTE: Z33.1 is not an acceptable principle diagnosis code.
Screening for Rh incompatibility: *first pregnancy visit (recommends Rh (D) blood typing and antibody testing for all pregnant women during their first visit for pregnancy-related care) *24-28 weeks gestation (recommends repeated Rh (D) antibody testing for all unsensitized Rh (D)-negative women at 24-28 weeks gestation unless the biological father is known to be Rh (D)-negative)	86850, 80055 or 80081	Z34.00, Z34.01, Z34.02, Z34.03, Z34.80, Z34.81, Z34.82, Z34.83, Z34.90, Z34.91, Z34.92, Z34.93, Z33.1, O09.00, O09.01, O09.02, O09.03, O09.10, O09.11, O09.12, O09.13, O09.291, O09.292, O09.293, O09.299, O09.40, O09.41, O09.42, O09.43, O09.211, O09.212, O09.213, O09.219, O09.30, O09.31, O09.32, O09.33, O09.511, O09.512, O09.513, O09.519, O09.521 O09.522, O09.523, O09.529, O09.611, O09.612, O09.613, O09.619, O09.621, O09.622, O09.623, O09.629, O09.811, O09.812, O09.813, O09.819, O09.821, O09.822, O09.823, O09.829, O36.80X0, O36.80X1, O36.80X2, O36.80X3, O36.80X4, O36.80X5, O36.80X9, O09.891, O09.892, O09.893, O09.899, O09.70, O09.71, O09.72, O09.73, O09.90, O09.91, O09.92, O09.93 O09.A0, O09.A1, O09.A2, O09.A3	Initial testing is part of the obstetric panel. Use 86850 only if performed separately from the panel for repeat testing. NOTE: Z33.1 is not an acceptable principle diagnosis code.
Counseling for Sexually Transmitted Infections (recommends high-intensity behavioral counseling to prevent STIs for all active adolescents and for adults at increased risk for STIs)	G0445, 99383, 99384, 99385, 99386, 99387, 99393, 99394, 99395, 99396, 99397 or 99401, 99402, 99403, 99404	Z71.7, Z71.89, Z72.51, Z72.52, Z72.53	99401-99404 are to be used in the absence of a wellness visit.
Behavioral Counseling to Prevent Skin Cancer (Counseling young adults, adolescents, children, and parents of young children about minimizing exposure to UV radiation for persons aged 6 months to 24 years with fair skin types to reduce their risk of skin cancer)	99383, 99384, 99385, 99393, 99394, 99395	Z00.121, Z00.129, Z00.110, Z00.111, Z00.00, Z00.01	Considered part of wellness office visit.
Screening for syphilis: *non-pregnant persons (screen persons at increased risk for syphilis infection) *pregnant women (screen all for syphilis infection)	80055, 86592, 86780 or 80081	Z34.00, Z34.01, Z34.02, Z34.03, Z34.80, Z34.81, Z34.82, Z34.83, Z34.90, Z34.91, Z34.92, Z34.93, Z33.1, O09.00, O09.01, O09.02, O09.03, O09.10, O09.11, O09.12, O09.13, O09.291, O09.292, O09.293, O09.299, O09.40, O09.41, O09.42, O09.43, O09.211, O09.212, O09.213, O09.219, O09.30, O09.31, O09.32, O09.33, O09.511, O09.512, O09.513, O09.519, O09.521 O09.522, O09.523, O09.529, O09.611, O09.612, O09.613, O09.619, O09.621, O09.622, O09.623, O09.629, O09.811, O09.812, O09.813, O09.819, O09.821, O09.822, O09.823, O09.829, O36.80X0, O36.80X1, O36.80X2, O36.80X3, O36.80X4, O36.80X5, O36.80X9, O09.891, O09.892, O09.893, O09.899, O09.70, O09.71, O09.72, O09.73, O09.90, O09.91, O09.92, O09.93, Z72.51, Z72.52, Z72.53, Z11.3 O09.A0, O09.A1, O09.A2, O09.A3	NOTE: Z33.1 is not an acceptable principle diagnosis code.
Statin Use for the Primary Prevention of Cardiovascular Disease in Adults (Initiate use of low- to moderate-dose statins in adults aged 40 to 75 years without a history of CVD who have 1 or more CVD risk factors (dyslipidemia, diabetes, hypertension, or smoking) and a calculated 10-year CVD event risk of 10% or greater)	Not applicable, administered through Pharmacy	Not applicable, administered through Pharmacy	Medications covered at 100%: Lovastatin (20 or 40 mg) or Pravastatin (10, 20 40 or 80 mg)



Grade A and B Recommendations of U.S. Preventive Services Task Force (USPSTF) currently effective unless otherwise noted Counseling for tobacco use (Ask all adults about tobacco use, advise them to stop using tobacco, and provide behavioral interventions and U.S. Food and Drug Administration—approved pharmacotherapy for cessation to adults who use tobacco AND ask all pregnant women about tobacco use and provide augmented, pregnancy- tailored counseling for those who smoke)	CPT or HCPCS 99385-99387, 99395-99397 or 99406-99407,	ICD-10 Diagnosis Z87.891, Z72.0, Z71.6, Z34.00, Z34.01, Z34.02, Z34.03, Z34.80, Z34.81, Z34.82, Z34.83, Z34.90, Z34.91, Z34.92, Z34.93, Z33.1, O09.00, O09.01, O09.02, O09.03, O09.10, O09.11, O09.12, O09.13, O09.291, O09.292, O09.293, O09.299, O09.40, O09.41, O09.42, O09.43, O09.211, O09.212, O09.213, O09.219, O09.30, O09.31, O09.32, O09.33, O09.511, O09.512, O09.513, O09.519, O09.521 O09.522, O09.523, O09.529, O09.611, O09.612, O09.613, O09.619, O09.621, O09.622, O09.623, O09.629, O09.811, O09.812, O09.813, O09.819, O09.821, O09.822, O09.823, O09.829, O36.80X0, O36.80X1, O36.80X2, O36.80X3, O36.80X4, O36.80X5, O36.80X9, O09.891, O09.892, O09.893, O09.899, O09.70, O09.71, O09.72, O09.73, O09.90, O09.91, O09.92, O09.93, F17.200, F17.201, F17.210, F17.211, F17.220, F17.221, F17.290, F17.291, O99.330, O99.331, O99.332, O99.333, O99.334, O99.335 O09.A0, O09.A1, O09.A2, O09.A3,	Comments FDA-approved tobacco cessation prescription medications and OTC nicotine replacement therapy (NRT) are covered at 100%. This is limited to a 90 day supply each for two cessation efforts. NRT is available only with a prescription. Telephonic counseling is available by calling 844-8NCQUIT. NOTE: Z33.1 is not an acceptable principle diagnosis code.
Primary care interventions to prevent tobacco use in children and adolescents (including education or brief counseling to prevent initiation of tobacco use among school-aged children and adolescents)	99383, 99384, 99385, 99393, 99394, 99395, 99401, 99402, 99403, 99404	Z00.121, Z00.129, Z00.110, Z00.111, Z00.00, Z00.01, U07.0	99401-99404 are to be used in the absence of a wellness visit.
Screening for visual acuity in children (vision screening for all children at least once between the ages of 3 and 5 years to detect the presence of amblyopia or its risk factors)	99382, 99392, 99173, 99174,or 99177	Z01.00, Z01.01, Z13.5	
HIV Infection Prevention with Pre- exposure prophylaxis (Persons at high risk of acquiring HIV. Offer preexposure prophylaxis (PrEP) with effective antiretroviral therapy to persons who are at high risk of HIV acquisition, including monitoring required prior to initiation of therapy and during drug therapy).	Drug therapy:	Z51.81 Effective 10/1/23 Z29.81	Medications covered at 100%: Truvada. For members who are clinically unable to utilize Truvada, Viread may be made available by completing this form, and faxing it to the number on the bottom of this document Effective 1/15/23: Apretude will be covered at 100% under the member's medical plan when billed with the diagnosis code listed.
Screening for Unhealthy Drug Use (Ask questions about unhealthy drug use in adults 18 years or older. Screening should be implemented when services for accurate diagnosis, effective treatment, and appropriate care can be offered or referred. (Screening refers to asking questions about unhealthy drug use, not testing biological specimens.)	99383, 99384, 99385, 99393, 99394, 99395	Z00.121, Z00.129, Z00.110, Z00.111, Z00.00, Z00.01	Considered part of wellness office visit.
Screening for anxiety in children and adolescents (aged 8-18 years)	99381, 99382, 99383, 99384, 99385, 99391, 99392, 99393, 99394, 99395	Z00.121, Z00.129, Z00.110, Z00.111	Done as part of wellness visit



ight Futures Recommendations for Children	CPT or HCPCS	ICD-10 Diagnosis	Comments	
Sensory Screening - Vision	99381, 99382, 99383, 99384, 99385, 99391, 99392, 99393, 99394, 99395, 99173, 99174 or 99177	Z00.121,Z00.129, Z00.110, Z00.111, Z13.5, Z01.00, Z01.01	May be part of well-child visits.	
ensory Screening - Hearing (beyond newborn screening)	99382, 99383, 99384, 99385, 99392, 99393, 99394, 99395, 92551, 92558 or 92586	Z00.121, Z00.129, Z00.110, Z00.111, Z01.10, Z01.118	May be part of wellness visits.	
Developmental Screening	96110	Z00.121, Z00.129 , Z13.42, Z13.49	96111 is not a screening service and should not be reported for this recommendation.	
Autism Screening	96110	Z13.41	96111 is not a screening service and should not b reported for this recommendation.	
Developmental Surveillance	99381, 99382, 99383, 99384, 99385, 99391, 99392, 99393, 99394, 99395	Z00.121, Z00.129, Z00.110, Z00.111	May be part of well-child visits.	
Psychosocial/Behavioral Assessment	99381, 99382, 99383, 99384, 99385, 99391, 99392, 99393, 99394, 99395	Z00.121, Z00.129, Z00.110, Z00.111	Part of well-child visits.	
Depression Screening (screening at ages 11-21)	99383 99384, 99385 99393, 99394, 99395, G0444	Z00.121, Z00.129, Z00.110, Z00.111 Z13.31		
Alcohol and Drug Use Assessment	99408 , 99409, G0442, G0443, G0396 or G0397 G2011	Z13.89, F10.10, F10.120, F10.129	These codes are to be used when the service is not of a wellness visit.	
Hematocrit or Hemoglobin	85013, 85014, 85018, 85025, 85027	Z00.121, Z00.129, Z00.110, Z00.111, Z13.0		
Congenital Heart Disease Screening	Not applicable, administered through Facility	Not applicable, administered through Facility	Screening for critical congenital heart disease using pul oximetry should be performed in newborns after 24 hou of age before discharge from the hospital. This screeni would be administered to newborn at the birth facility	
Lead Screening (up to 7 yrs)	83655	Z13.88		
Tuberculin Test	86580	Z11.1		
Dyslipidemia Screening (Cholesterol)	80061, 82465, or 83718	Z13.220		
STI/HIV Screening	G0432, G0433 or G0435 or G0475	Z34.00, Z34.01, Z34.02, Z34.03, Z34.80, Z34.81, Z34.82, Z34.83, Z34.90, Z34.91, Z34.92, Z34.93, Z33.1, O09.00, O09.01, O09.02, O09.03, O09.10, O09.11, O09.12, O09.13, O09.291, O09.292, O09.293, O09.299, O09.40, O09.41, O09.42, O09.43, O09.211, O09.212, O09.213, O09.219, O09.30, O09.31, O09.32, O09.33, O09.511, O09.512, O09.513, O09.519, O09.521 O09.522, O09.523, O09.529, O09.611, O09.612, O09.613, O09.619, O09.621, O09.622, O09.623, O09.629, O09.811, O09.812, O09.813, O09.819, O09.821, O09.822, O09.823, O09.829, O36.80X0, O36.80X1, O36.80X2, O36.80X3, O36.80X4, O36.80X5, O36.80X9, O09.891, O09.892, O09.893, O09.899, O09.70, O09.71, O09.72, O09.73, O09.90, O09.91, O09.92, O09.93, Z11.4 O09.A0, O09.A1, O09.A2, O09.A3	Screen adolescents for HIV at least once between the ages of 15 and 21 making every effort to preserve confidentiality of the adolescent, as per "Human Immunodeficiency Virus (HIV) Infection: Screening", and after initial screening, youth at increased risk of HIV infection should be retested annually or more frequently STI screening now references recommendations made it the AAP Red Book. This category was previously titled, "STI Screening". NOTE: Z33.1 is not an acceptable principle diagnosis code.	
Oral Health	99381, 99382, 99383, 99384, 99391, 99392, 99393, 99394			
Anticipatory Guidance	99381, 99382, 99383, 99384, 99391, 99392, 99393, 99394	Z00.121, Z00.129, Z00.110, Z00.111 May be part of well-child visits		
Bilirubin Screening niversal assessment of bilirubin level in infants with gestational age of 35 weeks or greater	82247, 88720	Z00.110 Mandate effective 1/1/18 or upon re Service is typically performed in the birth far of a wellness office visit in the event of a		
Postpartum depression screening Performed as part of 1, 2, 4, and 6 month well baby	96161	Z00.110, Z00.111, Z00.121, Z00.129, Z13.32	Mandate effective 1/1/18 or upon renewal To be billed as part of newborn/infant visit. Not mother	



Women's Preventive Services effective at renewal on or after			
August 1, 2012, unless otherwise			
noted			
Well Woman visits (Well-woman preventive care	CPT or HCPCS 99384, 99385,	ICD-10 Diagnosis Z00.00, Z00.01, Z01.411, Z01.419	Frequency Annual, although HHS recognizes that several visits may be
visit annually for adult women to obtain the recommended preventive services that are age and developmentally appropriate, including preconception and prenatal care. This well-woman visit should, where appropriate, include other preventive services listed in this set of guidelines	99386, 99387 or 99394, 99395, 99396, 99397	200.00, 200.01, 201.411, 201.419	needed to obtain all necessary recommended preventive services, depending on a woman's health status, health needs, and other risk factors
Screening for gestational diabetes	82950	Z34.00, Z34.01, Z34.02, Z34.03, Z34.80, Z34.81, Z34.82, Z34.83, Z34.90, Z34.91, Z34.92, Z34.93, Z33.1, O09.00, O09.01, O09.02, O09.03, O09.10, O09.11, O09.12, O09.13, O09.291, O09.292, O09.293, O09.299, O09.40, O09.41, O09.42, O09.43, O09.211, O09.212, O09.213, O09.219, O09.30, O09.31, O09.32, O09.33, O09.511, O09.512, O09.513, O09.519, O09.521 O09.522, O09.523, O09.529, O09.611, O09.612, O09.613, O09.619, O09.621, O09.622, O09.623, O09.629, O09.811, O09.812, O09.813, O09.819, O09.821, O09.822, O09.823, O09.829, O36.80X0, O36.80X1, O36.80X2, O36.80X3, O36.80X4, O36.80X5, O36.80X9, O09.891, O09.892, O09.893, O09.899, O09.70, O09.71, O09.72, O09.73, O09.90, O09.91, O09.92, O09.A0, O09.A1, O09.A2, O09.A3	In pregnant women between 24 and 28 weeks of gestation and at the first prenatal visit for pregnant women identified to be at high risk for diabetes. NOTE: Z33.1 is not an acceptable principle diagnosis code.
Human papillomavirus (HPV) testing (High-risk human papillomavirus DNA testing in women with normal cytology results.)	87623, 87624 , 87625 or G0476	Z11.51	Screening should begin at 30 years of age and should occur no more frequently than every 3 years.
Counseling for sexually transmitted infections for all sexually active women	Part of preventive visit, 99401-99404 or G0445	Z71.7, Z71.89, Z72.51, Z72.52, Z72.53	Annual. 99401-99404 are to be used in the absence of a wellness visit.
Counseling and Screening for Human Immune-deficiency Virus	Counseling: part of preventive visit or 99401, 99402, 99403, 99404 Screening: G0432, G0433 or G0435 or G0475	Z34.00, Z34.01, Z34.02, Z34.03, Z34.80, Z34.81, Z34.82, Z34.83, Z34.90, Z34.91, Z34.92, Z34.93, Z33.1, O09.00, O09.01, O09.02, O09.03, O09.10, O09.11, O09.12, O09.13, O09.291, O09.292, O09.293, O09.299, O09.40, O09.41, O09.42, O09.43, O09.211, O09.212, O09.213, O09.219, O09.30, O09.31, O09.32, O09.33, O09.511, O09.512, O09.513, O09.519, O09.521 O09.522, O09.523, O09.529, O09.611, O09.612, O09.613, O09.619, O09.621, O09.622, O09.623, O09.629, O09.811, O09.812, O09.813, O09.819, O09.821, O09.822, O09.823, O09.829, O36.80X0, O36.80X1, O36.80X2, O36.80X3, O36.80X4, O36.80X5, O36.80X9, O09.891, O09.892, O09.893, O09.899, O09.70, O09.71, O09.72, O09.73, O09.90, O09.91, O09.92, O09.93, Z71.7, Z11.4	Counseling and screening for human immune-deficiency virus for all sexually active women



Women's Preventive Services effective at renewal on or after August 1, 2012, unless otherwise noted	OPT - HODGO		
Contraceptive method (including-but not limited to, sterilization and IUD), and counseling (Group health plans sponsored by certain religious employers, and group health insurance coverage in connection with such plans, are exempt from the requirement to cover contraceptive services)	CPT or HCPCS Part of preventive visit (99384, 99385, 99386, 99394, 99395, 99396), 99401, 99402, 99403, 99404, S0610, S0612, S0613, Contraceptives A4261, A4264, J7300, J7302, J7306, J7307, J1050, J7301, , J7297, J7298 or J7296, 96372, 11976, S4981, S4989, 11981, 11982, 11983, Sterlization: 57170, 58300, 58600, 58605, 58611, 58670, 00952, 00851, 58301, 74740, 00840, 58661, 58700 Fertility App: 99199 Effective 4/1/2024 A9293 Pregnancy test 81025 Ultrasound 76830, 76857	Z34.00, Z34.01, Z34.02, Z34.03, Z34.80, Z34.81, Z34.82, Z34.83, Z34.90, Z34.91, Z34.92, Z34.93, Z33.1, O09.00, O09.01, O09.02, O09.03, O09.10, O09.11, O09.12, O09.13, O09.291, O09.292, O09.293, O09.299, O09.40, O09.41, O09.42, O09.43, O09.211, O09.212, O09.213, O09.219, O09.30, O09.31, O09.32, O09.33, O09.511, O09.512, O09.513, O09.519, O09.521 O09.522, O09.523, O09.529, O09.611, O09.612, O09.613, O09.619, O09.621, O09.622, O09.623, O09.629, O09.811, O09.812, O09.813, O09.819, O09.821, O09.822, O09.823, O09.829, O36.80X0, O36.80X1, O36.80X2, O36.80X3, O36.80X4, O36.80X5, O36.80X9, O09.891, O09.892, O09.893, O09.899, O09.70, O09.71, O09.72, O09.73, O09.90, O09.91, O09.92, O09.93, Z30.011, Z30.013, Z30.014,Z30.017, Z30.018, Z30.019, Z30.13, Z30.430, Z30.432, Z30.40, Z30.41, Z30.431, Z30.49, Z30.42, Z30.46, Z30.49, Z30.8, Z30.9, Z30.432, O09.A1, O09.A2, O09.A3 Diagnosis codes for urine pregnancy test: Z30.011, Z30.013, Z30.014,Z30.017, Z30.018, Z30.013, Z30.014,Z30.017, Z30.018, Z30.019, Z30.012, Z30.0430	Diaphragms, vaginal rings, contraceptive patches, female condoms, sponges, spermicides, and emergency contraception are available at a participating pharmacy with an MD prescription. Anesthesia services for 00952 and 00851 will be provided with no cost sharing to appropriate groups If there is a medical reason a member cannot take a generic contraceptive, the physician should review the clinical criteria and if appropriate, submit fax form. 99401-99404 are to be used in the absence of a wellness visit. Injectable contraceptives and administration (J1050, 96372) are covered when filed on the same claim and billed with a contraceptive encounter diagnosis. Effective 4/1/24, New instructions for submission of subscription to Natural Cycles. Natural Cycles is currently the only FDA-cleared fertility app. Members must download the Member Claim Form or the State Health Plan Claim Form to submit and include the following: the Natural Cycles receipt. Write the HCPCS code A9293 and diagnosis code Z30.8 on the receipt. Reimbursement will be per month-not annual subscription. Effective 1/1/2023 A pregnancy test prior to the initiation of a contraceptive will be covered with no cost sharing Ultrasound to confirm IUD or implantable contraceptive placement will be covered with no cost sharing NOTE: Z33.1 is not an acceptable principle diagnosis code.
Breastfeeding support, supplies, and counseling (Comprehensive lactation support and counseling, by a trained provider during pregnancy and/or in the postpartum period, and costs for breastfeeding equipment)	99401, 99402, 99403, S9443, E0602, E0603, A4281, A4282, A4283, A4284, A4285, A4286	O30.93, Z34.03, Z34.83, Z34.93, Z39.1	Counseling covered at 100% through in-network providers (i.e., OB/GYNs, midwives, facilities); one breast pump provided per pregnancy through in-network DME providers. Replacement supplies for breast pumps or initial supplies for member owned breast pump will be provided with no cost sharing to appropriate groups Replacement supplies cannot be provided with the initial breast pump as they are included in the initial purchase of the pump 99401-99403 are to be used in the absence of a wellness visit.
Screening and counseling for interpersonal and domestic violence The USPSTF recommends that clinicians screen for intimate partner violence (IPV) in women of reproductive age and provide or refer women who screen positive to ongoing support services.	Part of preventive visit or 99401, 99402, 99403, 99404	Z00.00, Z00.01, Z01.411, Z01.419	Annual 99401-99404 are to be used in the absence of a wellness visit.
Effective 1/1/19 or upon renewal Screening for Urinary Incontinence (Screening should assess whether women experience urinary incontinence and whether it impacts their activities and quality of life. Clinicians should refer women for further evaluation and treatment if indicated)	Part of preventive visit or 99401, 99402, 99403, 99404	Z00.00, Z00.01, Z01.411, Z01.419	Annual 99401-99404 are to be used in the absence of a wellness visit.
Counseling midlife women aged 40 to 60 years with normal or overweight body mass index (BMI) (18.5–29.9 kg/m2) to maintain weight or limit weight gain to prevent obesity. Counseling may include individualized discussion of healthy eating and physical activity.	Part of preventive visit or 99401, 99402, 99403, 99404, G0447 or G0473	Z71.3	Annual 99401-99404 are to be used in the absence of a wellness visit.



Vaccines Recommended by the Centers for Disease Control (CDC)	ICD-10 Diagnosis	Comments
Diphtheria, Tetanus, Pertussis Measles, Mumps, Rubella Haemophilus Infuenzae Type B Hepatitis A Hepatitis B Influenza Pneumococcal Meningococcal Human Papillomavirus (HPV)*. Inactivated Poliovirus Rotavirus Varicella Tetanus-Diphtheria /Tetanus-Diphtheria Acellular Pertussis Herpes Zoster (Shingles) RSV COVID-19 Adult and Child & Adolescent Immunization Schedules (for persons aged 0-6 years, 7-18 years, and "catch-up schedule") Refer to the CDC's posted schedule of immunizations http://www.cdc.gov/vaccines/schedules/index.html *The HPV vaccine (Gardasil) is covered to age 45 per the Food and Drug Administration (FDA) guidelines	Z23, Z29.11	Doses, recommended ages and recommended populations vary. All recommended routine immunizations will be allowed with no cost share. Z29.11 should only be used with CPT codes 90380, 90381

Revision History January 1, 2017 to January 1, 2024

Date of change	Mandate	Change
January 1, 2017	Screening for Abdominal Aortic Aneurysm	Addition of 76706 effective 1/1/17
	Hypertension in Adults-screening and home monitoring	Added coverage of home monitoring (ABPM). Add DX Ro3.o, CPT 93784,93786, 93788, 93790 and HCPCS A4670 for ABPM with criteria
	Screening for Breast Cancer	Added 77067 effective 1/1/18. Removed 77052, 77057 termed 12/31/16
	Colorectal Cancer Screening	Added new HCPCS code Go500 with following instructions: Moderate sedation (Go500) performed by the physician performing the procedure is allowed but cannot be billed on the same DOS as 00810
June 1, 2017	Screening for Breast Cancer	Addition of coverage for 3-D tomosynthesis and CPT code 77063
July 1, 2017	Breast Feeding Support, Supplies and Counseling	Added statement clarifying that replacement supplies cannot be provided at same time as initial breast pump as they are included in purchase of pump
October 1, 2017		Remove ICD-9 codes from all mandates
	Screening for latent tuberculosis	New mandate effective 10/1/17
	Counseling related to BRCA screening	Addition of diagnosis code Z ₇ 1.8 ₃ effective 10/1/17. Removed Z ₃ 1.5 as no longer applicable
	Screening for chlamydial infection in non-pregnant and pregnant women	Removed 87800 and 87801 as more specific codes should be used to report service
	Screening for gonorrhea for women	Removed 87800 and 87801 as more specific codes should be used to report service
	Vaccines	Added Shingrix to covered vaccines
December 1,2017	Statin Use for the Primary Prevention of Cardiovascular Disease in Adults	New mandate effective 12/1/17
January 1, 2018	Bilirubin Screening	New mandate effective 1/1/18 or upon group renewal
	Postpartum depression screening	New mandate effective 1/1/18 or upon group renewal
	Colorectal cancer screening	Added 00812 effective 1/1/18. Removed 00810 termed 12/31/17
	Contraceptives	Added J7296 effective 1/1/18. Removed Q9984 termed 12/31/17
	Screening for Breast Cancer	Removed Go202 termed 1231/17
	Screening for Cervical Cancer	Removed 88154 termed 12/31/17
April 1, 2018	Developmental screening	Added dx codes Zoo.129 and Zoo.120 to covered diagnosis codes effective 4/1/18
May 1, 2018	Screening for preeclampsia	New mandate effective 5/1/18 or upon group renewal
October 1, 2018	Postpartum depression screening	Added dx code Z13.32 effective 10/1/18
	Depression screening (adult and Bright Futures)	Added dx code Z13.31 effective 10/1/18. Removed Z13.89 effective 9/30/18 (should no longer be used as more specific code available)
	Autism screening	Added dx code Z13.41 effective 10/1/18. Removed Z13.89 effective 9/30/18 (should no longer be used as more specific code available)
January 1, 2019	Screening for Diabetes after Pregnancy	Added new mandated coverage for screening in women with previous diagnosis of gestational diabetes effective 1/1/19 or upon group renewal
	Screening and counseling to reduce alcohol misuse	Added HCPCS code G2011 effective 1/1/19
	Counseling related to BRCA screening	Added CPT codes 81163, 81164, 81165, 81166, 81167 effective 1/1/19. Removed CPT codes 81211, 81213, 81214 effective 12/31/18.
	Screening for Urinary Incontinence	New mandate effective 1/1/19 or upon group renewal
An Independent lice	nsee of the Blue Cross and Blue Shield Association. April 1, 2024	



Date of change	Mandate	Change
March 1, 2019	Contraceptives	Added Z30.017 and Z30.46
April 1, 2019	Fall Prevention	Revised description based on updated USPSTF recommendation
	Behavioral Counseling to Prevent Skin Cancer	Revised description based on updated USPSTF recommendation
July 1 2019	Screening for Osteoporosis	Revised description based on updated USPSTF recommendation
August 1, 2019	Colorectal Cancer Screening	Added 81528 (Cologuard) to covered tests.
September 1, 2019	Cervical Cancer Screening	Revised description based on updated USPSTF recommendation
October 1, 2019	Counseling for Tobacco Use	Removed 90-day limit on prescription and OTC medications
	Screening for and Management of Obesity in Adults	Revised description based on updated USPSTF recommendation
November 1, 2019	Screening and counseling to reduce unhealthy alcohol use	Revised description based on updated USPSTF recommendation
	Screening and counseling for interpersonal and domestic violence	Revised description based on updated USPSTF recommendation
July 1, 2020	Primary care interventions to prevent tobacco use in children and adolescents	Added Uo7.o to list of diagnoses
	HIV Infection Prevention with Pre-exposure prophylaxis	New mandate effective 7/1/2020 or upon group renewal
September 1, 2020	Counseling related to BRCA screening	Revised description based on updated USPSTF recommendation
	Screening for hepatitis C virus (HCV) infection	Revised description based on updated USPSTF recommendation
January 1, 2021	Anxiety Screening	New mandate effective 1/1/2021 or upon group renewal
	Screening for Lung Cancer	Removed G0297 termed 12/31/2020 Added 71271 effective 1/1/2021
May 11,2021	Screening for Lung Cancer	Updated the criteria to reflect new recommendation of ages 50-80 and 20 pack year
July 1, 2021	Screening for Unhealthy Drug Use	New mandate effective 7/1/2021 or upon group renewal
January 1, 2022	Contraceptives	Removed codes for Essure (58565,5834) oas it is no longer available commercially
March 7, 2022	Colorectal Cancer Screening	Updated criteria to reflect new recommendation of ages 45-75. Change effective for eligible members no later than 4/1/2022
Jun1 1, 2022	Colorectal Cancer Screening	Addition of coding instructions for colonoscopy after a positive stool based (FIT, Cologuard) or direct visualization test (sigmoidoscopy or CT colonography) to be covered under preventive benefit
	Healthy Weight and Weight Gain In Pregnancy: Behavioral Counseling Interventions	New mandate effective 6/1/2022 or upon group renewal
July 1, 2022	Contraceptives	Addition of salpingectomy and associated anesthesia as surgical method of sterilization covered under preventive benefit
August 1, 2022	Contraceptives	Addition of app-based fertility tools as method of contraception covered under preventive benefit and coding instructions
September 1, 2022	Screening for diabetes	Change in age range from 40-70 to 35-70 and addition of recommendation to refer patients with abnormal blood glucose levels to intensive behavioral counseling interventions to promote a healthful diet and physical activity
January 1, 2023	Contraceptives	Addition of coverage of pregnancy test prior to initiation of contraceptive and ultrasound for use in implant/IUD removal. Added coverage of ancillary services when sterilization procedure performed as outpatient
	Colorectal Cancer Screening	Addition of 45390 as a covered colonoscopy procedure
January 15, 2023	HIV Infection Prevention with Pre-exposure prophylaxis	Addition of Jo739 as covered under medical benefit
May 1, 20223	Aspirin to prevent CVD	Removed mandate. USPTF recommendation revised to C/D
October 1, 2023	HIV Infection Prevention with Pre-exposure prophylaxis	Addition of Z29.81 as a covered diagnosis
	Vaccines	Added Covid-19 vaccines and RSV vaccines(all ages) to list of covered vaccines
	STI/HIV Screening (Bright Futures)	Clarified frequency of screening
	Screening for anxiety in children and adolescents	New Mandate effective 11/1/23
January 1, 2024	HIV Infection Prevention with Pre-exposure prophylaxis	Addition of Goo11, Goo12 as covered codes
	Vaccines	Addition dx code Z29.11 to be used with RSV monoclonal antibody vaccines
April 1, 2024	Contraceptives	Addition of A9293 and revision of submission instructions for Natural Cycles fertility app including switching from annual to monthly reimbursement