

Corporate Medical Policy

Genetic Testing for Diagnosis of Inherited Peripheral Neuropathies AHS – M2072

File Name: genetic_testing_for_diagnosis_of_inherited_peripheral_neuropathies
Origination: 01/01/2019
Last Review: 07/2023

Description of Procedure or Service

The inherited peripheral neuropathies are a heterogeneous group of diseases that may be inherited in an autosomal dominant, autosomal recessive, or X-linked dominant manner. The inherited peripheral neuropathies can be divided into hereditary motor and sensory neuropathies (such as Charcot-Marie-Tooth disease), hereditary neuropathy with liability to pressure palsies, hereditary sensory and autonomic neuropathies, and other miscellaneous types (e.g., hereditary brachial plexopathy, giant axonal neuropathy). In addition to clinical presentation, nerve conduction studies, and family history, genetic testing can be used to diagnose specific inherited peripheral neuropathies (Kang, 2023a).

When pursuing genetic testing for inherited peripheral neuropathies, genetic counseling is strongly recommended.

Related Policies

Nerve Fiber Density Testing AHS – M2112
General Genetic Testing, Germline Disorders AHS – M2145
General Genetic Testing, Somatic Disorders AHS – M2146
Prenatal Screening (Genetic) AHS – M2179
Celiac Disease Testing AHS – G2043

*****Note: This Medical Policy is complex and technical. For questions concerning the technical language and/or specific clinical indications for its use, please consult your physician.**

Policy

BCBSNC will provide coverage for genetic testing for diagnosis of inherited peripheral neuropathies when it is determined the medical criteria or reimbursement guidelines below are met.

Benefits Application

This medical policy relates only to the services or supplies described herein. Please refer to the Member's Benefit Booklet for availability of benefits. Member's benefits may vary according to benefit design; therefore member benefit language should be reviewed before applying the terms of this medical policy.

When Genetic Testing for Diagnosis of Inherited Peripheral Neuropathies is covered

1. For individuals with clinical features of Charcot-Marie-Tooth (CMT) disease, but for whom a definitive diagnosis cannot be made without genetic testing, genetic testing for *PMP22* deletions/duplication, *GJB1* mutations, **and/or** *MFN2* mutations is considered medically necessary.

Genetic Testing for Diagnosis of Inherited Peripheral Neuropathies

AHS – M2072

2. For individuals with clinical features of CMT disease who have tested negative for common deleterious variants (*PMP22* deletions/duplication; *GJB1* or *MFN2* mutations), single gene or multi-gene panel testing for CMT disease risk genes is considered medically necessary.
3. For asymptomatic individuals who have a close blood relative (see Note 1) with a known deleterious mutation in a CMT gene, genetic testing for the known familial mutation is considered medically necessary.
4. For individuals who are clinically suspected of having hereditary neuropathy with liability to pressure palsies (HNPP), but for whom a definitive diagnosis cannot be made without genetic testing, genetic testing for *PMP22* deletions and duplications is considered medically necessary.
5. For individuals who are clinically suspected of having hereditary motor neuropathy (HMN), but for whom a definitive diagnosis cannot be made without genetic testing, genetic testing for *BSCL2* mutations is considered medically necessary.

Note 1: Close blood relatives include 1st-degree relatives (e.g., parents, siblings, and children), 2nd-degree relatives (e.g., grandparents, aunts, uncles, nieces, nephews, grandchildren, and half-siblings), and 3rd-degree relatives (great-grandparents, great-aunts, great-uncles, great-grandchildren, and first cousins).

Note 2: For 5 or more gene tests being run on the same platform please refer to AHS-R2162 Laboratory Procedures Medical Policy.

When Genetic Testing for Diagnosis of Inherited Peripheral Neuropathies is not covered

Genetic Testing for Diagnosis of Inherited Peripheral Neuropathies is not covered when the above criteria are not met.

Policy Guidelines

Background

Peripheral neuropathies encompass the set of disorders that primarily lead to peripheral nerve dysfunction. Symptoms typically include weakness of muscles at extremities, spine curvature, and loss of sensation at extremities (Kang, 2023c; UpToDate, 2023). Neuropathies may be caused by a variety of different factors, such as metabolic issues (including Fabry disease, Niemann-Pick disease, et al.) or present as a secondary symptom to another condition (such as Tangier disease) (Kang, 2023c).

Charcot-Marie-Tooth (CMT) disease, also known as hereditary motor sensory neuropathy, is a group of progressive disorders that affect the peripheral nerves. CMT is caused by a mutation in one of several myelin genes that result in defects in myelin structure, maintenance, or function within peripheral nerves. Charcot-Marie-Tooth disease is one of the most common inherited neurological disorders, affecting approximately 1 in 2,500 people in the United States (Kang, 2023a).

Symptoms

The neuropathy of CMT affects both motor and sensory nerves. Symptoms usually start in childhood and have a gradual progression. The severity of symptoms varies greatly among individuals and even among family members with the disease and gene mutation (Bird, 2023; NINDS, 2007, 2023). Typical symptoms include the following:

Genetic Testing for Diagnosis of Inherited Peripheral Neuropathies

AHS – M2072

- “Weakness or paralysis of the foot and lower leg muscles, which can cause difficulty lifting the foot (foot drop)
- A high-stepped gait with frequent tripping or falling
- Balance problems
- Foot deformities, such as high arches and curled toes (hammertoes)
- Lower legs may take on an "inverted champagne bottle" shape due to the loss of muscle bulk
- Reduced ability to feel heat, cold, and touch
- Weakness and atrophy may occur in the hands, causing difficulty with fine motor skills
- Decreased sense of vibration and position (proprioception)
- Curvature of the spine (scoliosis)
- Hip displacement
- Contractures (chronic shortening of muscles or tendons around joints)
- Muscle cramping
- Nerve pain” (NINDS, 2023)

Pain can range from mild to severe, and some people may need to rely on foot or leg braces or other orthopedic devices to maintain mobility. Some people living with CMT experience tremor, and vision and hearing can also be affected. In rare cases, breathing difficulties may occur if the nerves that control the muscles of the diaphragm are affected (NINDS, 2023).

Causes

CMT is caused by mutations in genes that produce proteins involved in the structure and function of either the peripheral nerve axon or the myelin sheath. Although different proteins are abnormal in different forms of CMT disease, all mutations affect the normal function of the peripheral nerves. There is little correlation between the genotype and phenotype of CMT; it is common to see differing mutations result in various clinical phenotypes all within the same gene (Kang, 2023a).

Pattern of Inheritance

The pattern of inheritance varies with the type of CMT disease. CMT1, most cases of CMT2, and most intermediate forms are inherited in an autosomal dominant pattern. CMT4, a few CMT2 subtypes, and some intermediate forms are inherited in an autosomal recessive pattern. CMTX is inherited in an X-linked pattern. Some cases of CMT disease result from a new mutation and occur in people with no history of the disorder in their family. In rare cases the gene mutation causing CMT disease is a new mutation which occurs spontaneously in the individual's genetic material and has not been passed down through the family (Kang, 2023a).

CMT1

Charcot-Marie-Tooth disease type 1 (CMT1) is a demyelinating peripheral neuropathy characterized by distal muscle weakness and atrophy, sensory loss, and are usually slowly progressive and often associated with *pes cavus* foot deformity and bilateral foot drop (Bird, 2023). The six subtypes of CMT1 shown in

Genetic Testing for Diagnosis of Inherited Peripheral Neuropathies

AHS – M2072

Table 1 are clinically indistinguishable and are designated solely on molecular findings (Saporta et al., 2011)

Table 1: Molecular Genetics of CMT1 (Saporta et al., 2011)

Locus Name	Proportion of CMT1 (excluding CMTX)	Gene	Protein Product
CMT1A	70%-80%	<i>PMP22</i>	Peripheral myelin protein 22
CMT1B	10%-12%	<i>MPZ</i>	Myelin protein P ₀
CMT1C	~1%	<i>LITAF</i>	Lipopolysaccharide-induced tumor necrosis factor-alpha factor
CMT1D	Unknown	<i>EGR2</i>	Early growth response protein 2
CMT1E	~1%	<i>PMP22</i>	Peripheral myelin protein 22 (sequence changes)
CMT1F/2E	Unknown	<i>NEFL</i>	Neurofilament light polypeptide

Charcot-Marie-Tooth disease type 1A (CMT1A) is an autosomal dominant disease that results from a duplication of the gene on chromosome 17 that carries the instructions for producing the peripheral myelin protein-22 (PMP-22). Overexpression of this gene causes the structure and function of the myelin sheath to be abnormal. A different neuropathy distinct from CMT1A called hereditary neuropathy with predisposition to pressure palsy (HNPP) is caused by a deletion of one of the PMP-22 genes. In this case, abnormally low levels of the PMP-22 gene result in episodic, recurrent demyelinating neuropathy (NINDS, 2007, 2023).

Charcot-Marie-Tooth disease type 1B (CMT1B) is an autosomal dominant disease caused by mutations in the gene that carries the instructions for manufacturing the myelin protein zero (P₀), which is another critical component of the myelin sheath. Most of these mutations are point mutations. As a result of abnormalities in P₀, CMT1B produces symptoms similar to those found in CMT1A (NINDS, 2007, 2023).

Other less common genetic causes of CMT1 result from mutations within *LITAF*, *EGR2*, *PMP22*, and *NEFL* genes, respectively (NINDS, 2023).

CMT2

Charcot-Marie-Tooth disease type 2 (CMT2) is an axonal (non-demyelinating) peripheral neuropathy characterized by distal muscle weakness and atrophy. Axonal peripheral neuropathy shows extensive clinical overlap with CMT1 (Bird, 2023). In general, individuals with CMT2 tend to be less disabled and have less sensory loss than individuals with CMT1 (Bird, 2023). It is less common than CMT1. CMT2A, the most common axonal form of CMT, is caused by mutations in Mitofusin 2, a protein associated with mitochondrial fusion. Symptoms are similar to those seen in CMT1, but people with CMT2 often have less disability and sensory loss than individuals with CMT1. Additionally, symptoms for CMT2 may have vocal cord or phrenic nerve involvement, causing speech or respiratory problems (NINDS, 2023).

Table 2: Molecular Genetics of CMT2 (Anthony Antonellis, 2018; Bird, 2023; Peter De Jonghe, 2011; Schindler, 2014; Züchner, 2013)

Locus	Proportion of CMT	Gene / Chromosome Locus	Protein Product
CMT2A1	Unknown	<i>KIF1B</i>	Kinesin-like protein KIF1B
CMT2A2 ¹	20%	<i>MFN2</i>	Mitofusin-2
CMT2B	Unknown	<i>RAB7A</i>	Ras-related protein Rab-7
CMT2B1	Unknown	<i>LMNA</i>	Lamin A/C
CMT2B2	Unknown	<i>MED25</i>	Mediator of RNA polymerase II transcription subunit 25
CMT2C ²	Unknown	<i>TRPV4</i>	Transient receptor potential cation channel subfamily V member 4
CMT2D ³	3%	<i>GARS</i>	Glycyl-tRNA synthetase
CMT2E/1F ⁴	4%	<i>NEFL</i>	Neurofilament light polypeptide
CMT2F	Unknown	<i>HSPB1</i>	Heat-shock protein beta-1
CMT2G	Unknown	12q12-q13	Unknown
CMT2H/2K	5%	<i>GDAP1</i>	Ganglioside-induced differentiation-associated protein-1
CMT2I/2J	Unknown	<i>MPZ</i>	Myelin protein P ₀
CMT2L	Unknown	<i>HSPB8</i>	Heat-shock protein beta-8
CMT2N	Unknown	<i>AARS</i>	Alanine--tRNA ligase, cytoplasmic
CMT2O	Unknown	<i>DYNC1H1</i>	Cytoplasmic dynein 1 heavy chain 1
CMT2P	Unknown	<i>LRSAM1</i>	E3 ubiquitin-protein ligase LRSAM1
CMT2S	Unknown	<i>IGHMBP2</i>	DNA-binding protein SMUBP-2
CMT2T	Unknown	<i>DNAJB2</i>	DnaJ homolog subfamily B member 2
CMT2U	Unknown	<i>MARS</i>	Methionine--tRNA ligase, cytoplasmic

CMT3

Dejerine-Sottas disease (CMT3), is a severe demyelinating neuropathy that begins in infancy. Infants have severe muscle atrophy, weakness, and sensory problems. This rare disorder can be caused by mutations in

Genetic Testing for Diagnosis of Inherited Peripheral Neuropathies

AHS – M2072

multiple genes, including *PMP22*, *MPZ*, and *EGR2*, and can be inherited either dominantly or recessively (NINDS, 2023).

CMT4

Charcot-Marie-Tooth disease type 4 (CMT4) comprises several different subtypes of autosomal recessive demyelinating motor and sensory axonal neuropathies. Each neuropathy subtype is caused by a different genetic mutation, may affect a particular ethnic population, and produces distinct physiologic or clinical characteristics. Affected individuals have the typical CMT phenotype of distal muscle weakness and atrophy associated with sensory loss and, frequently, pes cavus foot deformity. Several genes have been identified as causing CMT4, including *GDAP1* (CMT4A), *MTMR13* (CMT4B1), *MTMR2* (CMT4B2), *SH3TC2* (CMT4C), *NDG1* (CMT4D), *EGR2* (CMT4E), *PRX* (CMT4F), *FDG4* (CMT4H), and *FIG4* (CMT4J) (Kang, 2022, 2023a; NINDS, 2007, 2023).

Table 3: Molecular Genetics of CMT4 (Bird, 2017; Delague, 2013; Hamid Azzedine, 2015; Li, 2013)

Locus Name	Proportion of CMT4	Gene	Protein Product
CMT4A ¹	Unknown	<i>GDAP1</i>	Ganglioside-induced differentiation-associated protein 1
CMT4B1		<i>MTMR2</i>	Myotubularin-related protein 2
CMT4B2		<i>SBF2</i>	Myotubularin-related protein 13
CMT4C ²		<i>SH3TC2</i>	SH3 domain and tetratricopeptide repeats-containing protein 2
CMT4D		<i>NDRG1</i>	Protein NDRG1
CMT4E		<i>EGR2</i>	Early growth response protein 2
CMT4F		<i>PRX</i>	Periaxin
CMT4H ³		<i>FDG4</i>	FYVE, RhoGEF and PH domain-containing protein 4
CMT4J ⁴		<i>FIG4</i>	Phosphatidylinositol 3, 5 biphosphate

CMTX

CMTX is caused by a point mutation in the connexin-32 gene on the X chromosome. The connexin-32 protein is expressed in Schwann cells, which wrap around nerve axons and make up a single segment of the myelin sheath (NINDS, 2007, 2023). CMTX type 1 is characterized by a moderate to severe motor and sensory neuropathy. Hearing loss and central nervous system symptoms may also occur in certain affected families (Abrams, 2020).

Table 4: Molecular Genetics of CMTX (Bird, 2023; Kim, 2013)

Genetic Testing for Diagnosis of Inherited Peripheral Neuropathies

AHS – M2072

Disease Name	Proportion of X-Linked CMT	Gene / Chromosome Locus	Protein Product
CMTX1 ¹	90%	<i>GJB1</i>	Gap junction beta-1 protein (connexin 32)
CMTX2 ²	Unknown	Xp22.2	
CMTX3 ¹			Not applicable
CMTX4 ¹		<i>AIFM1</i>	Apoptosis-inducing factor 1
CMTX5 ²		<i>PRPS1</i>	Ribose-phosphate pyrophosphokinase 1
CMTX6 ¹		<i>PDK3</i>	Pyruvate dehydrogenase kinase isoform 3

Hereditary Brachial Plexopathy (Hereditary Neuralgic Amyotrophy)

This condition is primarily characterized by painful injuries to the brachial plexus nerves as well as episodic weakness of the shoulder and arm. Other symptoms such as winging of the scapula, short stature, neck folds, small face, and hypotelorism may be present. Nerve conduction velocity is typically normal, and the histopathology of this condition is non-specific. The *septin 9 gene (SEPT9)* on chromosome 17 has been associated with this condition (Bromberg, 2023).

Giant Axonal Neuropathy

This condition is characterized by disorganization of cytoskeletal intermediate filaments stemming from a mutated form of gigaxonin. Patients with this disorder often have a signature physical appearance; red and kinked hair, high foreheads, long eyelashes, and pale complexions are all hallmarks of this condition. The central nervous system may be affected as well with cerebellar dysfunction, spasticity, and potentially intellectual disability as possible symptoms. Nerve biopsy may show axonal loss or other axonal dysfunction. This diagnosis is confirmed by testing of the *GAN* gene (Kang, 2023c).

Hereditary Sensory and Autonomic Neuropathies (HSANs)

This subsection of disorders primarily encompasses non-motor neuropathies and are characterized by major loss of myelinated and unmyelinated fibers. These conditions are not as common as hereditary motor neuropathies and primarily present with sensory dysfunction, although motor functions may be affected. There are five main types of HSAN, each caused by different genes. Genes are associated as shown below (Eichler, 2023):

Disease Name (subtype)	Gene(s) or Locus	Examples of symptoms
HSAN1 (A)	<i>SPTLC1</i>	Distal sensory loss, distal muscle wasting
HSAN1 (B)	3p24-p22	Axonal neuropathy with distal sensory impairment
HSAN1 (C)	<i>SPTLC2</i>	Distal sensory loss, distal muscle wasting

Genetic Testing for Diagnosis of Inherited Peripheral Neuropathies

AHS – M2072

HSAN1 (D)	<i>ATL1</i>	Distal sensory loss, distal muscle wasting
HSAN1 (E)	<i>DNMT1</i>	Hearing loss, progressive dementia
HSAN1 (F)	<i>ATL3</i>	Distal sensory impairment
HSAN2 (A)	<i>HSN2</i>	Loss of pain, pressure, touch, and temperature sensation
HSAN2 (B)	<i>FAM134B</i>	Loss of pain, pressure, touch, and temperature sensation
HSAN2 (C)	<i>KIF1A</i>	Loss of pain, pressure, touch, and temperature sensation
HSAN2 (D)	<i>SCN9A</i>	Loss of pain and temperature sensation, hearing loss
HSAN3/Familial Dysautonomia	9q31	Dysautonomic crises, orthostatic hypotension
HSAN4/Congenital Insensitivity to Pain with Anhidrosis	<i>NTRK1</i>	Loss of pain sensation, thermoregulatory dysfunction
HSAN5	<i>NGFB</i>	Loss of pain and temperature sensation
HSAN6	<i>DST</i>	Lack of psychomotor development, respiratory difficulties
HSAN7	<i>SCN11A</i>	Inability to experience pain

Other unclassified HSANs exist, such as spastic paraplegia with ulcerations of the hands and feet (associated with *CCT5*) and sensory neuropathy with ichthyosis and anterior chamber syndrome (Eichler, 2023).

Genetic Testing

Charcot-Marie-Tooth disease is usually diagnosed by an extensive history and physical examination. The clinical diagnosis is then confirmed by electrodiagnostic tests like electromyography and nerve conduction velocity tests, and sometimes by nerve biopsy. Genetic testing is available for most types of CMT, and results are usually enough to confirm a diagnosis. Genetic testing can simplify the diagnosis of CMT by avoiding invasive procedures, such as nerve biopsy. In addition, early diagnosis can facilitate early interventions, including physical therapy. However, most therapies are only supportive (occupational, physical) and generally do not rely on the results of specific genetic testing (Kang, 2023a, 2023b). A positive genetic test can confirm diagnosis in most people with CMT. But a negative result does not exclude the disease, as an unidentified gene may be missed by DNA sampling (Charcot-Marie-Tooth News, 2023).

Genetic testing for CMT is complicated by the extensive underlying genetic heterogeneity. The CMT spectrum of disorders can be inherited in an autosomal dominant, autosomal recessive, or X-linked manner. The most commonly identified CMT subtypes are CMT1A (*PMP22* duplication), CMTX1 (*GJB1* mutation), hereditary neuropathy with liability to pressure palsies (*PMP22* deletion), CMT1B (*MPZ* mutation), and CMT2A (*MFN2* mutation). Together, these five subtypes account for 92 percent of genetically defined CMT cases. All other CMT subtypes and associated mutations each account for <1 percent of genetically defined CMT (CMTA, 2023; Kang, 2023a). Genetic screening for relatives of a patient diagnosed with CMT is an option, but risk assessment depends on several factors, including accuracy of the diagnosis, determination of the mode of inheritance for the individual family, and results of molecular genetic testing (Kang, 2023a). Numerous genetic panels are available for the assessment of peripheral neuropathies, such as GeneDx's panel (64 genes) and Invitae's panel (83 genes) (GeneDx, 2023; Invitae, 2023). Other panels include ones by Athena Diagnostics (23 genes) (Athena Diagnostics, 2023), Claritas Genomics (*PMP22* gene) (Claritas Genomics, 2023), MNG Laboratories (139 genes) (MNG

Genetic Testing for Diagnosis of Inherited Peripheral Neuropathies

AHS – M2072

Laboratories, 2022), Prevention Genetics (44 genes) (Prevention Genetics, 2023), and Variantyx's Genomic Unity panel (25 genes).

Clinical Utility and Validity

DiVincenzo et al. (2014) performed an analysis of the genetic landscape of CMT. 14 genes associated with CMT (*PMP22*, *GJB1*, *MPZ*, *MFN2*, *SH3TC2*, *GDAP1*, *NEFL*, *LITAF*, *GARS*, *HSPB1*, *FIG4*, *EGR2*, *PRX*, and *RAB7A*) were evaluated out of 3312 individuals. Deletions and duplications in the *PMP22* gene consisted of about 78% of positive findings, followed by mutations in the *GJB1* (6.7%), *MPZ* (5.3%), and *MFN2* (4.3%) genes. 71% of the pathogenic mutations found were missense mutations. Overall, 95% of the positive results involved one of four genes (*PMP22*, *GJB1*, *MPZ*, *MFN2*). The authors conclude that these four genes should be screened first before proceeding with further genetic testing (DiVincenzo et al., 2014).

Pareyson (2017) reviewed the current literature on CMT diagnosis stating that data justifies a step-wise algorithm considering a variety of factors, such as phenotype, nerve conduction velocities, and ethnicity. The authors note that NGS is steadily replacing older methods of sequencing in this algorithm. The authors propose evaluating the first few common genes (*PMP22*, *MPZ*, et al) and then considering larger sequencing methods such as NGS. However, due to the growing number of genes associated with CMT, these larger sequencing methods may be considered first-line. Finally, the authors state that due to the growing number of associated genes, newer classifications need to be discussed and validated further (Pareyson et al., 2017).


Rudnik-Schöneborn and colleagues (2016) evaluated the clinical features and genetic results of 1206 CMT patients and 124 affected relatives. Genetic detection rates were 56% in demyelinating CMT and 17% in axonal CMT. "Three genetic defects (*PMP22* duplication/deletion, *GJB1/Cx32* or *MPZ/P0* mutation) were responsible for 89.3% of demyelinating CMT index patients in whom a genetic diagnosis was achieved, and the diagnostic yield of the three main genetic defects in axonal CMT (*GJB1/Cx32*, *MFN2*, *MPZ/P0* mutations) was 84.2%". The authors concluded that "diagnostic algorithms are still useful for cost-efficient mutation detection and for the interpretation of large-scale genetic data made available by next generation sequencing strategies" (Rudnik-Schoneborn et al., 2016).

Vaeth et al. (2019) evaluated the effect of implementing a targeted next-generation sequencing (NGS) approach for identifying CMT. The authors stated that from 1992-2012, a total of 1442 CMT analyses were performed (through Sanger sequencing and other quantitative analyses) and a pathogenic variant was discovered in 21.6% of these cases. From this cohort, 195 samples that did not reach a definitive diagnosis were sequenced by a custom 63-gene panel. The authors identified a 5.6% increase in diagnostic yield using this targeted NGS approach (Vaeth et al., 2019).

Cortese et al. (2020) investigated the effectiveness of NGS panels in CMT. 220 patients were enrolled in the study and a targeted CMT NGS panel was performed. After NGS sequencing, a molecular diagnosis based on a pathogenic variant was found in 30% of the cases and variants of unknown significance were found in 33% of the cases. 39% of the cases held mutations in *GJB1*, *MFN2*, and *MPZ* while the others held mutations in *SH3TC2*, *GDAP1*, *IGHMBP2*, *LRSAM1*, *FDG4*, and *GARS*. Copy number changes were detected in *PMP22*, *MPZ*, *MFN2*, *SH3TC2*, and *FDG4*. The authors conclude that "NGS panels are effective tools in the diagnosis of CMT, leading to genetic confirmation in one-third of cases negative for *PMP22* duplication/deletion, thus highlighting how rarer and previously undiagnosed subtypes represent a relevant part of the genetic landscape of CMT" (Cortese et al., 2020).

Genetic Testing for Diagnosis of Inherited Peripheral Neuropathies

AHS – M2072



Rudnik-Schöneborn et al. (2020) suggested a diagnostic algorithm for genetic testing of suspected hereditary neuropathy. Advanced genetic sequencing allows for comprehensive evaluation of the pathogenic relevance of identified variants. As shown in the chart above, “If *PMP22* copy number analysis is negative, then clinical distinction of HNPP and CMT/dHMN will sort out patients for *PMP22* mutation analysis only and those for broader multigene testing. If a pedigree is compatible with X-linked inheritance, it is recommended to analyze coding and non-coding regions of *GJB1*. Patients who are tested negative for known neuropathy genes may be included in further whole exome or genome sequencing (WES/WGS) to detect mutations in rare and new genes” (Rudnik-Schöneborn et al., 2020).

Yalcintepe et al. (2021) studied the importance of multiple gene analysis for diagnosis of Charcot Marie Tooth Disease. Fifty-five patients with suspected CMT phenotype were examined using a customized multigene panel which was compared to the Multiplex Ligand Probe Amplification (MLPA) method. The custom panel identified 13 cases (7.15%) with a pathogenic/likely pathogenic variant. “The affected genes were *MARS1*, *NDRG1*, *GJB1*, *GDAP1*, *MFN2*, *PRX*, *SH3TC2*, and *FGD4*. Six cases (10.9%) had pathogenic variants in *GJB1* and *FGD4* genes, variants of unknown significance (VUS) in *HSPB3*, *CHRNA1*, *ARHGEF10*, and *KIF5A* genes. 21 cases (11.55%) had VUS with the genes *HSPB3*, *KIF1B*, *SCN11A*, *CHRNA1*, *HSPB1*, *FIG4*, *ARHGEF10*, *DHTKD1*, *SBF1*, *EGR2*, *SBF2*, *IGHMBP2*, *KIF5A*, and *DNAJB2*.” The authors concluded that the NGS customized panel was beneficial, time-saving, and cost-effective in the diagnosis of CMT (Yalcintepe et al., 2021).

Guidelines and Recommendations

American Academy of Neurology (AAN), the American Academy of Neuromuscular and Electrodiagnostic Medicine (AANEM), and the American Academy of Physical Medicine and Rehabilitation (AAPM&R)

The Polyneuropathy Task Force that included 19 physicians with representatives from the American Academy of Neurology (AAN), the American Academy of Neuromuscular and Electrodiagnostic Medicine (AANEM), and the American Academy of Physical Medicine and Rehabilitation (AAPM&R) concluded that “genetic testing is established as useful for the accurate diagnosis and classification of hereditary polyneuropathies (Class I)” (England et al., 2009).

Genetic Testing for Diagnosis of Inherited Peripheral Neuropathies

AHS – M2072

The Task Force stated that “for patients with a cryptogenic polyneuropathy who exhibit a classic hereditary neuropathy phenotype, routine genetic screening may be useful for CMT1A duplication/deletion and Cx32 mutations in the appropriate phenotype (Class III). Further genetic testing may be considered guided by the clinical question.” The Task Force recommended that “genetic testing should be conducted for the accurate diagnosis and classification of hereditary neuropathies (Level A)”. The Task force further recommended that “Genetic testing may be considered in patients with a cryptogenic polyneuropathy and classic hereditary neuropathy phenotype (Level C). Initial genetic testing should be guided by the clinical phenotype, inheritance pattern, and electrodiagnostic features and should focus on the most common abnormalities which are CMT1A [*PMP22*] duplication/HNPP deletion, Cx32 (*GJB1*), and *MFN2* mutation screening. There is insufficient evidence to support or refute the usefulness of routine genetic testing in cryptogenic polyneuropathy patients without a classic hereditary phenotype (Level U)” (England et al., 2009).

These guidelines were reaffirmed on January 22, 2022.

European Federation of Neurological Societies (EFNS)

The EFNS released recommendations on genetic testing for various types of peripheral neuropathies. Regarding CMT, they noted that “Given the rarity of AR CMT in the European population routine diagnostic screening of the many known genes is currently not feasible” but acknowledged that “Currently, molecular genetic testing can be offered for several of the more prevalent CMT genes.” EFNS stated that *PMP22* duplication should be tested first in patients presenting with CMT1, followed by sequencing of *GJB1*, *MPZ*, and *PMP22*. If a patient presents with CMT2, *MFN2* should be screened first, followed by *MPZ*. If a patient presents with intermediate CMT, *GJB1* and *MPZ* should be screened. EFNS notes that in patients with hereditary neuropathy with liability to pressure palsies will be investigated for a *PMP22* deletion at the same time as a screening for a *PMP22* duplication (Burgunder et al., 2011).

However, routine diagnostic screenings for hereditary motor neuropathy (HMN) and hereditary sensory-autonomic neuropathy (HSAN) are not feasible due to low mutation frequencies. If screening is performed for these conditions, EFNS recommends *BSCL2* as the first candidate for screening in HMN. *NTRK1* may also be screened for in congenital insensitivity to pain with anhidrosis patients (CIPA, a sub-phenotype of HSAN) and *RAB7* may be screened in CMT2B patients. Finally, *SEPT9* may be screened in the context of hereditary neuralgic amyotrophy (HNA) (Burgunder et al., 2011).

State and Federal Regulations, as applicable

Many labs have developed specific tests that they must validate and perform in house. These laboratory-developed tests (LDTs) are regulated by the Centers for Medicare and Medicaid (CMS) as high-complexity tests under the Clinical Laboratory Improvement Amendments of 1988 (CLIA '88). LDTs are not approved or cleared by the U. S. Food and Drug Administration; however, FDA clearance or approval is not currently required for clinical use.

Billing/Coding/Physician Documentation Information

This policy may apply to the following codes. Inclusion of a code in this section does not guarantee that it will be reimbursed. For further information on reimbursement guidelines, please see Administrative Policies on the Blue Cross Blue Shield of North Carolina web site at www.bcbsnc.com. They are listed in the Category Search on the Medical Policy search page.

Applicable service codes: 81324, 81325, 81326, 81403, 81404, 81405, 81406, 81448, 96040, S0265

Genetic Testing for Diagnosis of Inherited Peripheral Neuropathies

AHS – M2072

BCBSNC may request medical records for determination of medical necessity. When medical records are requested, letters of support and/or explanation are often useful, but are not sufficient documentation unless all specific information needed to make a medical necessity determination is included.

Scientific Background and Reference Sources

- Abrams, C. (2020). GJB1 Disorders: Charcot Marie Tooth Neuropathy (CMT1X) and Central Nervous System Phenotypes. <https://www.ncbi.nlm.nih.gov/books/NBK1374/>
- Anthony Antonellis, P., Lev G Goldfarb, MD, and Kumaraswamy Sivakumar, MD. (2018). GARS-Associated Axonal Neuropathy. <https://www.ncbi.nlm.nih.gov/books/NBK1242/>
- Athena Diagnostics. (2023). CMT Advanced Evaluation - Comprehensive. <https://www.athenadiagnostics.com/view-full-catalog/c/cmt-advanced-evaluation-comprehensive>
- Bird, T. (2017). GDAP1-Related Hereditary Motor and Sensory Neuropathy. <https://www.ncbi.nlm.nih.gov/books/NBK1539/>
- Bird, T. (2023). Charcot-Marie-Tooth (CMT) Hereditary Neuropathy Overview. <https://www.ncbi.nlm.nih.gov/books/NBK1358/>
- Bromberg, M. (2023, 05/08/2023). *Brachial plexus syndromes*. <https://www.uptodate.com/contents/brachial-plexus-syndromes>
- Burgunder, J. M., Schols, L., Baets, J., Andersen, P., Gasser, T., Szolnoki, Z., Fontaine, B., Van Broeckhoven, C., Di Donato, S., De Jonghe, P., Lynch, T., Mariotti, C., Spinazzola, A., Tabrizi, S. J., Tallaksen, C., Zeviani, M., Harbo, H. F., & Finsterer, J. (2011). EFNS guidelines for the molecular diagnosis of neurogenetic disorders: motoneuron, peripheral nerve and muscle disorders. *Eur J Neurol*, 18(2), 207-217. <https://doi.org/10.1111/j.1468-1331.2010.03069.x>
- Charcot-Marie-Tooth News. (2023). *Genetic Tests and Genetic Counseling*. <https://charcot-marie-toothnews.com/genetic-tests/>
- Claritas Genomics. (2023). PMP22 Deletion/Duplication. <http://www.claritasgenomics.com/test/pmp22-deletionduplication/index.html>
- CMTA. (2023). Genetic Testing. <https://www.cmtausa.org/living-with-cmt/find-help/genetic-testing/>
- Cortese, A., Wilcox, J. E., Polke, J. M., Poh, R., Skorupinska, M., Rossor, A. M., Laura, M., Tomaselli, P. J., Houlden, H., Shy, M. E., & Reilly, M. M. (2020). Targeted next-generation sequencing panels in the diagnosis of Charcot-Marie-Tooth disease. *Neurology*, 94(1), e51-e61. <https://doi.org/10.1212/WNL.00000000000008672>
- Delague, V. (2013). Charcot-Marie-Tooth Neuropathy Type 4H. <https://www.ncbi.nlm.nih.gov/books/NBK153601/>
- DiVincenzo, C., Elzinga, C. D., Medeiros, A. C., Karbassi, I., Jones, J. R., Evans, M. C., Braastad, C. D., Bishop, C. M., Jaremkov, M., Wang, Z., Liaquat, K., Hoffman, C. A., York, M. D., Batish, S. D., Lupski, J. R., & Higgins, J. J. (2014). The allelic spectrum of Charcot-Marie-Tooth disease in over 17,000 individuals with neuropathy. *Mol Genet Genomic Med*, 2(6), 522-529. <https://doi.org/10.1002/mgg3.106>
- Eichler, F. (2023, 05/08/2023). *Hereditary sensory and autonomic neuropathies*. <https://www.uptodate.com/contents/hereditary-sensory-and-autonomic-neuropathies>

Genetic Testing for Diagnosis of Inherited Peripheral Neuropathies

AHS – M2072

- England, J. D., Gronseth, G. S., Franklin, G., Carter, G. T., Kinsella, L. J., Cohen, J. A., Asbury, A. K., Szigeti, K., Lupski, J. R., Latov, N., Lewis, R. A., Low, P. A., Fisher, M. A., Herrmann, D. N., Howard, J. F., Lauria, G., Miller, R. G., Polydefkis, M., & Sumner, A. J. (2009). Practice Parameter: Evaluation of distal symmetric polyneuropathy: Role of laboratory and genetic testing (an evidence-based review). *Neurology*, 72(2), 185.
<https://doi.org/10.1212/01.wnl.0000336370.51010.a1>
- GeneDx. (2023). Hereditary Neuropathy Panel. <https://www.genedx.com/tests/detail/hereditary-neuropathy-panel-800>
- Hamid Azzedine, E. L., and Mustafa A Salih. (2015). Charcot-Marie-Tooth Neuropathy Type 4C. <https://www.ncbi.nlm.nih.gov/books/NBK1340/>
- Invitae. (2023). *Invitae Comprehensive Neuropathies Panel*. https://www.invitae.com/en/physician/tests/03200/#info-panel-assay_information
- Kang, P. (2022, 01-20-2022). *Charcot-Marie-Tooth disease: Genetics, clinical features, and diagnosis*. <https://www.uptodate.com/contents/charcot-marie-tooth-disease-genetics-clinical-features-and-diagnosis>
- Kang, P. (2023a). *Charcot-Marie-Tooth disease: Genetics, clinical features, and diagnosis*. <https://www.uptodate.com/contents/charcot-marie-tooth-disease-genetics-clinical-features-and-diagnosis>
- Kang, P. (2023b). *Charcot-Marie-Tooth disease: Management and prognosis*. <https://www.uptodate.com/contents/charcot-marie-tooth-disease-management-and-prognosis>
- Kang, P. (2023c). *Overview of hereditary neuropathies*. Retrieved 05/08/2023 from <https://www.uptodate.com/contents/overview-of-hereditary-neuropathies>
- Kim, J.-W., Kim, Hee-Jin. (2013). Charcot-Marie-Tooth Neuropathy X Type 5. <https://www.ncbi.nlm.nih.gov/books/NBK1876/>
- Li, J. (2013). Charcot-Marie-Tooth Neuropathy Type 4J. <https://www.ncbi.nlm.nih.gov/books/NBK169431/>
- MNG Laboratories. (2022). Charcot-Marie-Tooth Disease, Axonal (NGS Panel and Copy Number Analysis + mtDNA). <https://mnglabs.com/tests/NGS345A/charcot-marie-tooth-disease-axonal-ngs-panel-and-copy-number-analysis-mtdna>
- NINDS. (2007). Charcot-Marie-Tooth Disease Fact Sheet. https://www.ninds.nih.gov/Disorders/Patient-Caregiver-Education/Fact-Sheets/Charcot-Marie-Tooth-Disease-Fact-Sheet#3092_5
- NINDS. (2023). *Charcot-Marie-Tooth Disease*. <https://www.ninds.nih.gov/health-information/disorders/charcot-marie-tooth-disease>
- Pareyson, D., Saveri, P., & Pisciotta, C. (2017). New developments in Charcot-Marie-Tooth neuropathy and related diseases. *Curr Opin Neurol*, 30(5), 471-480.
<https://doi.org/10.1097/wco.0000000000000474>
- Peter De Jonghe, M., PhD and Albena K Jordanova, PhD. (2011). Charcot-Marie-Tooth Neuropathy Type 2E/1F. <https://www.ncbi.nlm.nih.gov/books/NBK1187/>

Genetic Testing for Diagnosis of Inherited Peripheral Neuropathies AHS – M2072

- Prevention Genetics. (2023). Charcot-Marie-Tooth (CMT) - Comprehensive Panel.
<https://www.preventiongenetics.com/testInfo?val=Charcot-Marie-Tooth+%28CMT%29+-+Comprehensive+Panel>
- Rudnik-Schöneborn, S., Auer-Grumbach, M., & Senderek, J. (2020). Charcot-Marie-Tooth disease and hereditary motor neuropathies – Update 2020. *Medizinische Genetik*, 32(3), 207-219.
<https://doi.org/doi:10.1515/medgen-2020-2038>
- Rudnik-Schöneborn, S., Tolle, D., Senderek, J., Eggermann, K., Elbracht, M., Kornak, U., von der Hagen, M., Kirschner, J., Leube, B., Müller-Felber, W., Schara, U., von Au, K., Wiczorek, D., Bussmann, C., & Zerres, K. (2016). Diagnostic algorithms in Charcot-Marie-Tooth neuropathies: experiences from a German genetic laboratory on the basis of 1206 index patients. *Clin Genet*, 89(1), 34-43. <https://doi.org/10.1111/cge.12594>
- Saporta, A. S., Sottile, S. L., Miller, L. J., Feely, S. M., Siskind, C. E., & Shy, M. E. (2011). Charcot-Marie-Tooth disease subtypes and genetic testing strategies. *Ann Neurol*, 69(1), 22-33.
<https://doi.org/10.1002/ana.22166>
- Schindler, A. (2014). TRPV4-Associated Disorders. <https://www.ncbi.nlm.nih.gov/books/NBK201366/>
- UpToDate. (2023). *Patient education: Charcot-Marie-Tooth disease (The Basics)*.
<https://www.uptodate.com/contents/charcot-marie-tooth-disease-the-basics>
- Vaeth, S., Christensen, R., Duno, M., Lildballe, D. L., Thorsen, K., Vissing, J., Svenstrup, K., Hertz, J. M., Andersen, H., & Jensen, U. B. (2019). Genetic analysis of Charcot-Marie-Tooth disease in Denmark and the implementation of a next generation sequencing platform. *Eur J Med Genet*, 62(1), 1-8. <https://doi.org/10.1016/j.ejmg.2018.04.003>
- Yalcintepe, S., Gurkan, H., Dogan, I. G., Demir, S., Sag, S. O., Kabayegit, Z. M., Atli, E. I., Atli, E., Eker, D., & Temel, S. G. (2021). The Importance of Multiple Gene Analysis for Diagnosis and Differential Diagnosis in Charcot Marie Tooth Disease. *Turk Neurosurg*, 31(6), 888-895.
<https://doi.org/10.5137/1019-5149.jtn.33661-21.3>
- Züchner, S. (2013). Charcot-Marie-Tooth Neuropathy Type 2A.
<https://www.ncbi.nlm.nih.gov/books/NBK1511/>
- Specialty Matched Consultant Advisory Panel review 7/2019
- Medical Director review 7/2019
- Specialty Matched Consultant Advisory Panel review 7/2020
- Medical Director review 7/2020
- Specialty Matched Consultant Advisory Panel review 7/2021
- Medical Director review 7/2021
- Medical Director review 7/2022
- Medical Director review 7/2023

Genetic Testing for Diagnosis of Inherited Peripheral Neuropathies

AHS – M2072

Policy Implementation/Update Information

- 1/1/2019 BCBSNC will provide coverage for genetic testing for diagnosis of inherited peripheral neuropathies when it is determined to be medically necessary because criteria and guidelines are met. Medical Director review 1/1/2019. Policy noticed 1/1/2019 for effective date 4/1/2019. (jd)
- 8/13/2019 Specialty Matched Consultant Advisory Panel review 7/2019. Medical Director review 7/2019 (jd)
- 9/10/2019 Reviewed by Avalon 2nd Quarter 2019 CAB. Related Policies added to Description section. When Covered section revised as follows: Item 1, added “or other inherited peripheral neuropathies”; item 2, added “If results indicate a demyelinating neuropathy, then first test for the most commonly identified CMT subtype, CMT1A (PMP22 duplication.”, and removed items a. and b. related to specific values for velocity testing in nerve conduction and specific cascade testing; added item 6 for Genetic testing for Hereditary Motor Neuropathy (HMN) (BSCL2 gene), and added “Note” which refers to policy, General Genetic Testing, Germline Disorders AHS – M2145 for all other uncommon hereditary peripheral neuropathy gene testing. Removed the following statement from the When Not Covered section: “Genetic testing for all other inherited peripheral neuropathies is considered investigational”. Code table removed from the Billing/Coding section and reimbursement statement added. Policy guidelines and references updated. Medical Director review. (jd)
- 10/29/19 Wording in the Policy, When Covered, and/or Not Covered section(s) changed from Medical Necessity to Reimbursement language, where needed. (hb)
- 7/28/20 Reviewed by Avalon 2nd Quarter 2020 CAB. Policy guidelines and references updated. Specialty Matched Consultant Advisory Panel review 7/2020. Medical Director review 7/2020. (jd)
- 8/24/21 Reviewed by Avalon 2nd Quarter 2021 CAB. The following statement was added to the When Not Covered section: “Genetic testing for CMT in asymptomatic individuals is considered investigational.” Background, policy guidelines, and references updated. Specialty Matched Consultant Advisory Panel review 7/2021. Medical Director review 7/2021. (jd)
- 9/13/22 Reviewed by Avalon 2nd Quarter 2022 CAB. Updated background, guidelines, and references. Billing/Coding section updated. Coverage criteria edited for clarity. No change to policy statement. Medical Director review 7/2022. (tm)
- 8/15/23 Reviewed by Avalon 2nd Quarter 2023 CAB. Updated Description, Policy Guidelines, and References. The following updates were made to the When Covered section: Former coverage criteria 2 split and expanded into new coverage criteria 1 and 2 for full clarity regarding which genes should be ordered first based on prevalence of cause. Recommendation for genetic counseling removed as coverage criteria and moved into the Policy Description. Removed former coverage criteria 4 (prenatal screening addressed in M2179) and coverage criteria 5 regarding peripheral nerve biopsy. previous Note 1 removed and replaced with new Note 1. Medical Director review 7/2023. (tm)

Medical policy is not an authorization, certification, explanation of benefits or a contract. Benefits and eligibility are determined before medical guidelines and payment guidelines are applied. Benefits are determined by the group contract and subscriber certificate that is in effect at the time services are rendered. This document is solely provided for informational

Genetic Testing for Diagnosis of Inherited Peripheral Neuropathies

AHS – M2072

purposes only and is based on research of current medical literature and review of common medical practices in the treatment and diagnosis of disease. Medical practices and knowledge are constantly changing and BCBSNC reserves the right to review and revise its medical policies periodically.