

# MY 2025 Blue Cross NC Provider Reference Guide

This version of the MY 2025 Blue Cross NC Provider Reference Guide is based on the preliminary HEDIS Specifications. Final HEDIS Specifications are not released by NCQA until end of March 2025. Blue Cross NC will update the MY 2025 Provider Reference Guide at that time if any changes are needed.

## Requirements for Meeting Clinical Goals - Last Updated March 2025



This reference guide is not intended to be a complete guide for all HEDIS® measures and requirements.

For additional details and specifications for HEDIS® measures, please go to https://www.ncqa.org/hedis/measures or contact Blue Cross NC Quality Management at qualitymanagement@bcbsnc.com or 919-765-4809.

MEASURE ABBREVIATION	MEASURE DESCRIPTION	PAGE
	Important Information	3
	Helpful Links and Resources	3
AAB	Avoidance of Antibiotic Treatment for Acute Bronchitis / Bronchiolitis	4
ADD-E	Follow-Up Care for Children Prescribed ADHD Medication	4
AIS-E	Adult Immunization Status	5
AMR	Asthma Medication Ratio	6
APM-E	Metabolic Monitoring for Children and Adolescents on Antipsychotics	6
APP	Use of First-Line Psychosocial Care of Children and Adolescents on Antipsychotics	7
BCS-E	Breast Cancer Screening	7
BPC-E	Blood Pressure Control for Patients with Hypertension	8
BPD	Blood Pressure Control for Patients with Diabetes	9
CBP	Controlling High Blood Pressure	10
CCS-E	Cervical Cancer Screening	11
CHL	Chlamydia Screening	12
CIS-E	Childhood Immunization Status	13
COA	Care for Older Adults	14
COL-E	Colorectal Cancer Screening	15
COU	Risk of Continued Opioid Use	16
CWP	Appropriate Testing for Pharyngitis	16
DAE	Use of High-Risk Medications in Older Adults	17
DBM-E	Documented Assessment After Mammogram	18
DDE	Potentially Harmful Drug-Disease Interactions in Older Adults	19
EED	Eye Exam for Patients with Diabetes	20
FMA-E	Follow-Up After Abnormal Mammogram Assessment	21
FMC	Follow-Up After Emergency Department Visit for People with High-Risk Chronic Conditions	22
FUA	Follow-Up After Emergency Department Visit for Substance Use	23
FUH	Follow-Up after Hospitalization for Mental Illness	24
FUI	Follow-Up After High-Intensity Care for Substance Use Disorder	25
FUM	Follow-Up After Emergency Department Visit for Mental Illness	26
GSD	Glycemic Status Assessment for Patients with Diabetes	27
GSD	Glycemic Status Assessment for Patients with Diabetes	28
HDO	Use of Opioids at High Dosage	29
IET	Initiation and Engagement of Substance Use Disorder Treatment	29
IMA-E	Immunizations for Adolescents	30
KED	Kidney Health Evaluation for Patients with Diabetes	31
LBP	Use of Imaging Studies for Low Back Pain	32
OMW	Osteoporosis Management in Women Who Had a Fracture	33
OSW	Osteoporosis Screening in Older Women	34
PBH	Persistence of Beta-Blocker Treatment after a Heart Attack	34
PCE	Pharmacotherapy Management of COPD Exacerbation	35
PCR	Plan All-Cause Readmissions	35
POD	Pharmacotherapy for Opioid Use Disorder	36
PPC	Prenatal and Postpartum Care	36
PRS-E	Prenatal Immunization Status	
SAA	Adherence to Antipsychotic Medications for Individuals with Schizophrenia	37

#### **TABLE OF CONTENTS**

## Requirements for Meeting Clinical Goals - Last Updated March 2025



This reference guide is not intended to be a complete guide for all HEDIS® measures and requirements.

For additional details and specifications for HEDIS® measures, please go to https://www.ncqa.org/hedis/measures or contact Blue Cross NC Quality Management at qualitymanagement@bcbsnc.com or 919-765-4809.

MEASURE ABBREVIATION	MEASURE DESCRIPTION	PAGE
SNS-E	Social Need Screening and Intervention	39
SPC	Statin Therapy for Patients With Cardiovascular Disease	40
SPD	Statin Therapy for Patients With Diabetes	41
TRC	Transitions of Care – Medication Reconciliation Post-Discharge	42
TRC	Transitions of Care – Notification of Inpatient Admission	43
TRC	Transitions of Care – Receipt of Discharge Information	43
TRC	Transitions of Care – Patient Engagement After Inpatient Discharge	44
UOP	Use of Opioids from Multiple Providers	45
URI	Appropriate Treatment for Upper Respiratory Infection	45
W30	Well-Child Visits in the first 30 Months of Life	46
WCC	Weight Assessment and Counseling for Nutrition and Physical Activity for Children / Adolescents – BMI Percentile	46
WCC	Weight Assessment and Counseling for Nutrition and Physical Activity for Children / Adolescents – Counseling for Nutrition	47
WCC	Weight Assessment and Counseling for Nutrition and Physical Activity for Children / Adolescents – Counseling for Physical Activity	48
WCV	Child and Adolescent Well-Care Visits	48
Medicare Part D	Medication Adherence for Cholesterol (Statins)	49
Medicare Part D	Medication Adherence for Hypertension (RAS Antagonists)	50
Appendix 1	BMI Coding Chart	51
Appendix 2	Measures that can accept medical records for care gap closure	51
Appendix 3	Measures used in overall Medicare Star Ratings	52
Appendix 4	CAHPS and Medicare HOS Survey information	53
Appendix 5	Medicare Medications with \$0 copay	54
	ADD-E	56
	CIS	57
	FMC	58
	FUA	59-61
	FUH	62,63
	FUI	64,65
	FUM	66,67
	IET	68-70
	PPC	71-75

#### **TABLE OF CONTENTS**

## **Important Information**



#### A comprehensive list of medication and NDC codes can be found on www.ncga.org

#### IMPORTANT EXCLUSION INFORMATION

- All exclusions are required.
- All deceased members are required to be excluded, even if there was compliance before death.

#### IMPORTANT DEFINITIONS

- Measurement year Considered 1/1 to 12/31 unless otherwise stated within measure specifications.
- Direct Reference Code A code included directly within the measure specifications with no associated value set or value set name. Note: Some Direct Reference Codes took the place of previous value sets that were retired due to only having one code.
- Administrative Measure Transaction data or other administrative data are used to identify the eligible population and numerator.
- Hybrid Measure Organizations look for numerator compliance in both administrative and medical record data.
- IPSD Index Prescription Start Date
- IESD Index Episode Start Date

#### RACE AND ETHNICITY STRATIFICATION INCLUDED MEASURES:

- AIS-E Adult Immunization Status
- AMR Asthma Medication Ratio
- BCS-E Breast Cancer Screening
- CBP Controlling High Blood Pressure
- CCS-E Cervical Cancer Screening
- CIS-E Childhood Immunization Status Combo 10 Indicator
- COL-E Colorectal Cancer Screening
- EED Eye Exam for Patients with Diabetes
- FUA Follow-up After Emergency Department Visit for Substance Use
- FUH Follow-Up After Hospitalization for Mental Illness
- FUM Follow-up After Emergency Department Visit for Mental Illness

- GSD Glycemic Status Assessment in Patients with Diabetes
- IET Initiation and Engagement of Substance Use Disorder Treatment
- IMA-E Immunizations for Adolescents
- KED Kidney Health Evaluation for Patients with Diabetes
- PDS-E Postpartum Depression Screening and Follow-Up
- PND-E Prenatal Depression Screening and Follow-Up
- POD Pharmacotherapy for Opioid Use Disorder
- PPC Prenatal and Postpartum Care
- PRS-E Prenatal Immunization Status
- W30 Well-Child Visits in the First 30 Months of Life
- WCV Child and Adolescent Well-Care Visits

## Helpful Links and Resources



Sign up for Provider E-briefs:

https://www.BlueCrossNC.com/Providers/Provider-News

Provider Blueline<sup>™</sup>:

**1-800-214-4844** (toll free)



**Network Management:** 1-800-777-1643 (toll free)

919-765-4349 (fax)

ProviderUpdates@bcbsnc.com (email)

RISK ADJUSTMENT CONTACTS					
Prospective Risk	Prospective				
Adjustment Program	Adjustment Cha				

Prospective Risk **Adjustment Chart Requests** 

**Davina Bowden** 

Team Leader, Clinical Teams Davina.Bowden@bcbsnc.com

919-765-3200

**George Taylor** 

Team Lead, Operations George.Taylor@bcbsnc.com

919-765-2750





HEDIS® QUALITY MEASURE	CLINICAL GOAL	CRITERIA TO MEET THE GOAL
AAB – Avoidance of Antibiotic Treatment for Acute Bronchitis / Bronchiolitis	Members treated for acute bronchitis / bronchiolitis should <b>NOT</b> be prescribed antibiotics unless there are	Claims:  Numerator compliance for this measure is based on pharmacy claims.
Administrative Measure  The percentage of episodes for	co-morbid conditions or competing diagnoses that require antibiotic therapy.	Medical Record documentation not applicable.  Required Exclusion:
members 3 months of age and older with a diagnosis of acute	require antibiotic therapy.	Members in hospice or using hospice services anytime during the measurement year.
bronchitis / bronchiolitis that did <b>NOT</b> result in an antibiotic	PRODUCT LINE	Note: To avoid inappropriate members being included into
The Intake Period: 7/1 of the year prior to measurement year to 6/30 of the measurement year	Commercial, Marketplace, Medicare and Medicaid	the eligible population, submit a claim with all appropriate diagnosis codes including any competing conditions (i.e. cellulitis, pharyngitis) and any co-morbid condition diagnoses. Telehealth visits are allowed for this measure.

# ADD-E

HEDIS® QUALITY MEASURE	CLINICAL GOAL	CRITERIA TO MEET THE GOAL				
ADD-E – Follow-Up Care for Children Prescribed ADHD Medication ECDS Measure The percentage of children newly	Claims:  • Submit a claim for a follow-up visit with a prescribing provider and the appropriate CPT and ICD-10 ADHD codes within 30 days of staring the medication and then at least two additional visits with a provider in the 9 months following the 30 day visit.					
prescribed ADHD medication who had at least 3 follow-up	Continuation and	Medical reco	rd doc	umentation not ap	plicable.	
visits within a 10 month period;	Maintenance Phase:	Required Exc	lusion	s:		
one needs to be within 30 days of when the medication was dispensed.  The Intake Period: 12 month window starting 3/1 of the year	The percentage of members 6 – 12 years of age with a prescription dispensed for ADHD medication, who remained on the medication for at least 210 days and who, in addition to the visit in the Initiation Phase, had at least two follow-up visits with a practitioner within 270 days (9 months) after the Initiation Phase ended. Only one of the two visits may be an e-visit or virtual check-in.	<ul> <li>Members in hospice or using hospice services anytime during the measurement year.</li> <li>Members with a diagnosis of narcolepsy any time during their history through December 31 of the measurement year.</li> </ul>				
prior to the measurement year and ending the last calendar day		HEDIS Value Set Name	Code Set	Code	Comments	
of February of the measurement year.		Visit Setting Unspecified	CPT	90791, 90792, 90832, 90833, 90834, 90836, 90837, 90838, 90839, 90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221, 99222, 99223, 99231, 99232, 99233, 99238, 99239, 99252, 99253, 99254,	With one of the following:  Outpatient POS  POS code 52  POS code 53  Teleheatlh POS code	
	PRODUCT LINE			99255		
	Commercial and Medicaid	Remaining H	HEDIS	Value Set codes car	n be found in Appendix 7.	



HEDIS® QUALITY MEASURE	CLINICAL GOAL	CRITERIA	A TO I	MEET THE GOAL			
AIS-E – Adult Immunization Status	Annual routine influenza vaccination; tetanus, diphtheria, and acellular	Claims: Submit a claim for all immuniz	zations	given.			
Status	pertussis (Tdap) and/or	Report all immunizations to the North Carolina Immunization Registry.					
ECDS Measure	tetanus and diphtheria (Td) vaccine; herpes zoster;	Required Exclusions:					
The percentage of members 19 years of	pneumococcal and hepatitis B vaccination for adults at various ages.	time during the measuremer	nt perio				
age and older who are up to date on	Members 19 years of age	Members who die any time of	uring	the measurement period.			
recommended routine	and older:	Notes:					
vaccines for influenza, tetanus, and diphtheria (Td) or tetanus,	Influenza: 1 influenza vaccine on or between July 1 of the year prior to	Additional numerator complia     All immunization numerators before or during the measure	s: Anap	phylaxis due to the vaccine at any time			
diphtheria, and acellular pertussis (Tdap), zoster, pneumococcal and hepatitis B.	the measurement period and June 30 of the measurement period.  Td/Tdap: At least one TD vaccine OR one TDAP	Hepatitis B: A history of hepa surface antigen, hepatitis B s B core antigen test, any time	ntis B il urface before	Iness; or had a positive hepatitis B antibody, or total antibody to hepatitis or during the measurement period. ohtheria, tetanus, or pertussis vaccine.			
	vaccine between 9 years prior to the start of the measurement period	HEDIS Value Set Name	Code				
	and the end of the measurement period.	Adult Immunization Value Set	CVX	135, 140, 141, 144, 150, 153, 155, 158, 166, 168, 171, 185, 186, 197, 205			
	Members 19-59 years of age: • Hepatitis B: Any of the following:	Adult Influenza Vaccine Procedure	CPT	90630, 90653, 90654, 90656, 90658, 90661, 90662, 90673, 90674, 90682, 90686, 90688, 90689, 90694, 90756			
	At least 3 doses of the childhood hepatitis	Influenza Virus LAIV Immunization	CVX	111, 149			
	vaccine.  ‡ Received a hepatitis B	Influenza Virus LAIV Vaccine Procedure	CPT	90672			
	vaccine series on or after their 19th birthday,	Td Immunization	CVX	09, 113, 138, 139			
	before or during the measurement period.	Td Immunization	CPT	90714, 90715			
	Members 50 years of age	Herpes Zoster Recombinant Vaccine Procedure	CPT	90750			
	<ul><li>and older:</li><li>Zoster: 2 doses of the</li></ul>	Adult Pneumococcal Immunization	CVX	33, 109, 133, 152, 215, 216			
	herpes zoster recombinant	Adult Pneumococcal Vaccine Procedure	CPT	90670, 90671, 90677, 90732			
	vaccine at least 28 days apart, on Oct 1, 2017,	Adult Pneumococcal Vaccine Procedure	HCPCS				
	through the end of the	Hepatitis B Immunization	CVX	08, 44, 45, 51, 110 ,146 198			
	measurement period.	Hepatitis B Vaccine Procedure	CPT	90697, 90723, 90740, 90744, 90747, 90748			
	Members 65 years and older:	Hepatitis B Vaccine Procedure	HCPCS				
	Pneumococcal: At least one dose of an adult pneumococcal vaccine on	Direct Reference Code (Newborn Hepatitis B Vaccination)	ICD-10- PCS	3E0234Z			
	or after their 19th birthday, before or during the	Adult Hepatitis B Immunization (3 dose)	CVX	43, 44, 45, 104, 220			
	measurement period.	Adult Hepatitis B Vaccine Procedure (2 dose)	CPT	90739, 90740, 90743, 90744, 90746, 90747, 90759			
	PRODUCT LINE	Direct Reference Code (Hepatitis B Vaccination 2 Dose)	CVX				
	Commercial, Medicaid and Medicare	Hepatitis B	ICD10CM	B16.0, B16.1, B16.2, B16.9, B17.0, B18.0, B18.1, B19.10, B19.11			





HEDIS® QUALITY MEASURE	CLINICAL GOAL	CRITERIA TO MEET THE GOAL
AMR – Asthma Medication Ratio	Members with persistent asthma	Claims:
Administrative Measure	will receive asthma controller prescriptions that account for at least 50% of their total asthma	<ul> <li>Numerator compliance for this measure is based on pharmacy claims.</li> </ul>
The percentage of members 5 – 64 years of age who were	medications. (Controller medications versus rescue	Medical record documentation not applicable.
identified as having persistent	medications.)	Required Exclusion:
asthma and had a ratio of controller medications to total		Members in hospice or using hospice services anytime during the measurement year.
asthma medications of 0.50 or	PRODUCT LINE	Members who had a diagnosis that requires a different
greater during the measurement year.	Commercial, Marketplace	treatment approach than members with asthma.
	and Medicaid	Members who had no asthma controller or reliever medications dispensed during the measurement year.

## APM-E

HEDIS® QUALITY MEASURE	CLINICAL GOAL	CRITERIA TO MEET THE GOAL				
APM-E – Metabolic Monitoring for Children and Adolescents on Antipsychotics	Children and adolescents who had two or more antipsychotic medication dispensing events during the measurement year	Claims:  Blood Glucose Testing Rate:  Submit a claim for at least one test for blood glucose or HbA1c				
The percentage of children and adolescents 1 – 17 years of age who had two or more antipsychotic prescriptions and had metabolic testing.  Three rates are reported:	will have metabolic testing.	or HbA1c.  Cholesterol Testing Rate:  Submit a claim for at least one test for LDL-C or cholesterol.  Blood Glucose and Cholesterol Testing Rate:  Submit claims for both of the following during the measurement year on the same or different dates of service:  At least one test for blood glucose or HbA1c.  At least one test for LDL-C or cholesterol.				
Percentage of children and adolescents on antipsychotics		Medical record Required Exclusion		entation not applicable.		
<ul><li>who received:</li><li>Blood glucose testing.</li><li>Cholesterol testing.</li></ul>	PRODUCT LINE  Commercial and Medicaid	Members in hospice or using hospice services any during the measurement year.				
Blood glucose and cholesterol testing.	Commercial and Wedicald	HEDIS Value Set Name	Code Set	Code	Comments	
		Glucose Lab Test	СРТ	80047, 80048, 80050, 80053, 80069, 82947, 82950, 82951		
		HbA1c Lab Test	CPT	83036, 83037		
		HbA1c Lab Test	LOINC	17856-6, 4548-4, 4549-2, 96595-4, 17855-8		
		HbA1c Test Result or Finding	CPT – CAT-II	3044F, 3046F, 3051F, 3052F	Without a 1P, 2P, 3P, or 8P modifier, or POS code 81	
		LDL-C Lab Test	CPT	80061, 83700, 83701, 83704, 83721		
		LDL-C Test Result or Finding	CPT – CAT-II	3048F, 3049F, 3050F	Without a 1P, 2P, 3P, or 8P modifier, or POS code 81	
		Cholesterol Lab Test	CPT	82465, 83718, 83722, 84478		





HEDIS® QUALITY MEASURE	CLINICAL GOAL	CRITERIA TO MEET THE GOAL									
APP – Use of First-Line Psychosocial Care for Children and Adolescents on	Children and adolescents aged 1 – 17 years of age will have	<ul> <li>Claims:</li> <li>Submit claim for psychosocial care 4 months prior to the IPSD through the 30 days after the IPSD.</li> </ul>					Submit claim for psychosocial care 4 months prior to the IPSD through the 30				
Antipsychotics  Administrative Measure  The percentage of children and adolescents 1 – 17 years of age who had a new prescription for an antipsychotic medication and had documentation	psychosocial care as first-line treatment prior to receiving new prescription for antipsychotic medication.	<ul> <li>Required Exclusions:</li> <li>Exclude members who met any of the following during the measurement year:</li> <li>Members in hospice or using hospice services any time during the measurement year.</li> <li>One acute inpatient encounter or 2 visits in an outpatient, intensive outpatient, or partial hospitalization with diagnosis of schizophrenia, schizoaffective disorder, bipolar disorder, other psychotic disorder, autism, or other developmental disorder during the measurement year.</li> <li>An outpatient visit for Behavioral Health.</li> <li>A community mental health center visit.</li> </ul>									
of psychosocial care as first-line treatment.	PRODUCT LINE Commercial,	<ul> <li>Electroconvulsive therapy.</li> <li>An observation visit, telehealth visit, telephone visit, e-visit, or virtual check-in with POS value set.</li> </ul>					• An observation visit, telehealth visit, telephone visit, e-visit, or virtual check				
	Medicaid	HEDIS Value Code Set Name Code									
		Psychosocial Care         CPT 90832, 90833, 90834, 90836, 90837, 90838, 90839, 90840, 90845, 90846, 90849, 90853, 90875, 90876, 90880									
	Psychosocial HCPCS G0176, G0177, G0409, G0410, G0411, H0004, H0035, H0 H0039, H0040, H2000, H2001, H2011, H2012, H2013, H2019, H2020, S0201, S9480, S9484, S9485										

# BCS-E

Cancer Screening the ag years	ge of 50 and 74 will have one or mammograms		ammograr					
	st every 2 years.	<ul> <li>Submit a claim for mammogram any time on or between October 1 two years prior to the measurement period and 12/31 of the measurement year.</li> <li>Required Exclusions:</li> <li>Bilateral mastectomy.</li> <li>Unilateral mastectomy with a bilateral modifier from the same procedure.</li> </ul>						
There is a 27 month look back period from 10/1 two years prior	ODUCT LINE nercial,	eral mastectomy found in clinical data with bilateral modifier from same lure.  are members 66 years of age and older as of 12/31 of the measurement tho are in an I-SNP or living long-term in an institution any time during the rement year.  ers 66 years of age and older as of 12/31 of the measurement year with and advanced illness.  ers receiving palliative care or hospice services during the measurement						
to 12/31 of the current year.	ledicald	Members who had gender-affirming chest surgery with a diagnosis of gendysphoria any time during the member's history through the end of the measurement period.      HEDIS Value Set Name						



HEDIS® QUALITY MEASURE	CLINICAL GOAL	CRITERIA TO MEET THE GOAL				
BPC-E – Blood Pressure Control for Patients with Hypertension  ECDS Measure  Measure Description: The percentage of members 18 – 85 years of age who had a diagnosis of hypertension (HTN) and whose most recent blood pressure (BP) was <140/90 mm Hg during the measurement period.  Identify members who had:  • At least two outpatient visits, telephone visits, e-visits or virtual check-ins on different dates of service with a diagnosis of hypertension on or between January 1 of the year prior to the measurement period and June 30 of the measurement period.  • At least one outpatient visit, telephone visit, e-visit or virtual check-in with a diagnosis of hypertension and at least one dispensed antihypertensive medication on or between January 1 of the year prior to the measurement period and June 30 of the measurement period.	Treatment of adults who have hypertension to a standard blood pressure target (<140/90 mm Hg) to reduce the risk of all-cause and cardiovascular mortality.  PRODUCT LINE  Commercial, Medicaid and Medicare	Claims:  Submit a clair results of the  The member is <140/90 mm measurement  Note: BPs mus of HTN. BPs talvisit do not coulous a hosp period.  Members with measurement of Members with disease (ESRI prior to the late) Members with nephrectomy, member's his measurement of Members with measurement of Members with nephrectomy.  Members with nephrectomy member's his measurement of Members with nephrectomy.  Members with nephrectomy member of of measurement of Members and the measurement of Members 81 years two indictional during the measurement of Members with measurement of Memb	m with the BP at each is numerate in Hg. The reach is numerate in Hg, if the term of the period, or the period. It is a constant is a constant in Hg. The reach is a constant in Hg. The r	2 apprroutin or commember re is not if the or after cute in any tire. The cute in any tire in part of age frailty is a commember of age.	ropriate CPTII codes to report e office visit. Inpliant if the representative BP er is not compliant if the BP of BP reading during the reading is incomplete.  In the date of second diagnosis patient setting or during an ED  palliative care services or elect me during the measurement  attient admission during the the indicates end-stage renal g the member's history on or surement period.  In tindicates ESRD: dialysis, lant any time during the to the last day of the  pregnancy any time during the finage and older as of 12/31 of the in an I-SNP or living long- during the measurement year. and older as of 12/31 of the and advanced illness. Tolder as of the 12/31 with at with different dates of service	





HEDIS® QUALITY MEASURE	CLINICAL GOAL	CRITERIA TO MEET THE GOAL					
BPD – Blood Pressure Control for Patients with Diabetes  Hybrid Measure	Members with diabetes will have blood pressure control of < 140/90 mm Hg.	Submit a claim with the 2 appropriate CPTII codes to report results of the BP at each office visit.					
Trybita Weasure		Medical record documentation*:					
Members 18 – 75 years of age with diabetes (Types 1 and 2)		<ul> <li>A visit note identifying the most recent BP reading (&lt; 140/90mm Hg) during the measurement year.</li> <li>If multiple BP measurements occur on the same date, or are noted in the chart on the same date, use the lowest systolic and lowest diastolic reading.</li> </ul>					
whose BP was adequately controlled (< 140/90) during the measurement year.							
		<b>Note:</b> BP readings taken by the member with a digital cuff and documented in the member's medical record are eligible for use in reporting.					
		Required Exclu	ısions:				
		Members red	ceiving pall	iative o	care or hospice services during		
		the measure	•	oore o	f age and older as of 12/31		
	PRODUCT LINE	of the measu	rement yea	ar who	are in an I-SNP or living long-		
	Commercial, Medicaid and Medicare	<ul> <li>term in an institution any time during the measurement year</li> <li>Members 66 of age and older as of 12/31 of the measurement year with frailty and advanced illness.</li> <li>*Medical record documentation is only accepted for gap closure during the annual HEDIS Medical Record Review.</li> </ul>					
		HEDIS Value Set Name	Code Set	Code	Comments*		
		Systolic and Diastolic Result	CPT-CAT-II	3074F	Most recent systolic blood pressure < 130 mm Hg (Systolic Compliant)		
		Systolic and Diastolic Result	CPT-CAT-II	3075F	Most recent systolic blood pressure 130-139 mm Hg (Systolic Compliant)		
		Systolic and Diastolic Result	CPT-CAT-II		≥ 140 mm Hg (Systolic Non-Compliant)		
		Systolic and Diastolic Result	CPT-CAT-II	3078F	Most recent diastolic blood pressure < 80 mm Hg (Diastolic Compliant)		
		Systolic and Diastolic Result	CPT-CAT-II	3079F	Most recent diastolic blood pressure 80-89 mm Hg (Diastolic Compliant)		
		Systolic and Diastolic Result	CPT-CAT-II	3080F	Most recent diastolic blood pressure ≥ 90 mm Hg (Diastolic Non-Compliant)		
		*(Without a 1P	, 2P, 3P, or	8P mo	difier, or POS code 81)		





		1				
HEDIS® QUALITY MEASURE	CLINICAL GOAL		CRITERIA	TO M	EET THE GOAL	
CBP – Controlling High Blood Pressure	Control of high BP < 140/90 is extremely important to prevent additional cardiovascular	<ul> <li>Claims:</li> <li>Submit a claim with the 2 appropriate CPTII codes to report results of the BP at each routine office visit.</li> </ul>				
Hybrid Measure  The percentage of members	conditions.	Note: BPs must occur on or after the date of second diagnosis of HTN. BPs taken in an acute inpatient setting or during an ED visit do not count.				
18 – 85 years of age who had a		Medical record documentation*:				
diagnosis of hypertension (HTN) and whose BP was controlled						
during the measurement year.  • BP goal < 140/90 mm Hg			n Hg) durin	g the n	st recent BP reading neasurement year on or after tension.	
Identification and a bad at					occur on the same date, or are	
Identify members who had at least two outpatient visits,		noted in the and lowest d			e date, use the lowest systolic	
telephone visits, e-visits or				_	member with a digital cuff and	
virtual check-ins on different					nedical record are eligible for	
dates of service with a diagnosis of hypertension on or between		use in reportin	ıg.		-	
January 1 of the year prior to	PRODUCT LINE	Required Exclu	usions:			
the measurement year and June 30 of the measurement year.	Commercial, DSNP, Marketplace, Medicare and Medicaid	<ul> <li>Members with evidence of end-stage rena nephrectomy or kidney transplant any time member's history on or prior to December measurement year.</li> <li>Members with a diagnosis of pregnancy and</li> </ul>	lant any time during the to December 31 of the			
		<ul> <li>Members receiving palliative care or hospice services during the measurement year.</li> <li>Medicare members 66 years of age and older as of 12/31 of the measurement year who are in an I-SNP or living long-term in an institution any time during the measurement year</li> </ul>				
					and older as of 12/31 of the	
			•		and advanced illness.	
		<ul> <li>Members 81 years of age and older as of 12/31 of the measurement year with at least two indications of frailty on different dates of service.</li> </ul>				
			cal record documentation is <b>only</b> accepted for ga e during the annual HEDIS Medical Record Revie			
		HEDIS Value Set Name	Code Set	Code	Comments*	
		Systolic and Diastolic Result	CPT-CAT-II	3074F	Most recent systolic blood pressure < 130 mm Hg (Systolic Compliant)	
		Systolic and Diastolic Result	CPT-CAT-II	3075F	130-139 mm Hg (Systolic Compliant)	
		Systolic and Diastolic Result	CPT-CAT-II	3077F	Most recent systolic blood pressure ≥ 140 mm Hg (Systolic Non-Compliant)	
		Systolic and Diastolic Result	CPT-CAT-II	3078F	Most recent diastolic blood pressure < 80 mm Hg (Diastolic Compliant)	
		Systolic and Diastolic Result	CPT-CAT-II	3079F	Most recent diastolic blood pressure 80-89 mm Hg (Diastolic Compliant)	
		Systolic and Diastolic Result	CPT-CAT-II	3080F	Most recent diastolic blood pressure ≥ 90 mm Hg (Diastolic Non-Compliant)	

\*(Without a 1P, 2P, 3P, or 8P modifier, or POS code 81)



HEDIS® QUALITY MEASURE	CLINICAL GOAL	C	RITERI <i>A</i>	A TO MEET THE GO	DAL
CCS-E - Cervical Cancer Screening ECDS Measure	Members 21 – 64 will have a cervical cytology (Pap smear) every 3 years.	Claims: Submit a claim for procedures completed with appropriate codes. Submit a claim using the appropriate code for history of exclusion.			
The percentage of members 21 – 64 years of age who were screened for cervical cancer.  The measurement year is 1/1 to 12/31.	Members 30 – 64 years of age who were ned for cervical cancer.  Members 30 – 64 years of age will have primary HPV testing every 5 years.  Members age 30 – 64 will have cervical cytology with HPV co-testing every 5 years.  Note: HPV reflex testing does not count for the 5 years.  Members 30 – 64 years of age will have will have cervical cytology with HPV co-testing every 5 years.		d results 5 years. on of "tol rectomy exclusions: with no ence of co gh Decen eiving pa nent year	residual cervix, cervi ervix anytime during nber 31 of the measu lliative care or hospic	ete" abdominal / neet for  ical agenesis or the member's rement year. ce services during
		the patient's h	Code	Code	Comments
		Set Name Direct Reference Code	Set LOINC	76689-9 (Sex assigned at birth), LA2-8 (Male)	Required exclusion when used together at any time in patient's history
	PRODUCT LINE Commercial, Marketplace	Cervical Cytology Lab Test	CPT	CPT 88141, 88142, 88143, 88147, 88148, 88150, 88152, 88153, 88164, 88165, 88166, 88167, 88174, 88175	
	and Medicaid	Cervical HCPCS G0123, G0124, G0141, Cytology G0143, G0144, G0145,	G0123, G0124, G0141, G0143, G0144, G0145, G0147, G0148, P3000,		
		High Risk HPV Lab Test	CPT	87624, 87625	
		High Risk HPV Lab Test	HCPCS	G0476	
		Absence of Cervix Diagnosis	ICD10M	Q51.5, Z90.710, Z90.712	Required Exclusion
		Hysterectomy With No Residual Cervix	СРТ	57530, 57531, 57540, 57545, 57550, 57555, 57556, 58150, 58152, 58200, 58210, 58240, 58260, 58262, 58263, 58267, 58270, 58275, 58280, 58285, 58290, 58291, 58292, 58293, 58294, 58548, 58550, 58552, 58553, 58554, 58570, 58571, 58572, 58573, 58575, 58951, 58953, 58954, 58956, 59135	Required Exclusion





HEDIS® QUALITY MEASURE	CLINICAL GOAL	CI	RITERIA	A TO MEET THE	GOAL
CHL – Chlamydia Screening  Administrative Measure  The percentage of members 16 – 24 years of age who were recommended for routine chlamydia screening, were identified as sexually active and had at least one test for chlamydia during the measurement year.	Annual screening for chlamydia for all sexually active members 16 – 24 years or age who were recommended for screening.  Obtain a urine sample or obtain a direct sample (i.e. cervix, urethra, vagina) for chlamydia culture yearly.	Claims: Submit a claim for chlamydia screening with appropriate codes.  Medical record documentation: A visit note indicating date of service and lab results showing chlamydia screening was performed.  Note: The two methods to determine sexual activity are pharmacy data (contraceptives) and claims encounter for sexual activity, pregnancy test or pregnancy diagnosis.			I lab results rmed. al activity are s encounter for
measurement year.		Required Exclus	ions:		
	PRODUCT LINE	Members in h during the me		or using hospice se	rvices anytime
	Commercial, Marketplace and Medicaid	_		ssigned male at bir	th.
		HEDIS Value Set Name	Code Set	Code	Comments
		Chlamydia Tests	CPT	87110, 87270, 87320, 87490, 87491, 87492, 87810	
		Direct Reference Code	LOINC	76689-9 (Sex assigned at birth), LA2-8 (Male)	Required exclusion when used together at any time in patient's history







#### **HEDIS® QUALITY MEASURE CLINICAL GOAL** CRITERIA TO MEET THE GOAL **COA - Care for Older Adults** Medication Review - Either of Claims: the following meets criteria: • Submit a claim for services completed with appropriate **Administrative Measure** codes. · Both of the following during the same visit during the The percentage of adults 66 measurement year where Medical record documentation\*: years and older who had each the provider type is a • Medication review conducted by a prescribing practitioner of the following during the prescribing practitioner or or clinical pharmacist during the measurement year and the measurement year: clinical pharmacist: presence of a medication list in the medical record. · Medication review. At least one medication Medication list, signed and dated during the measurement Functional status review. year by the appropriate practitioner type. assessment. The presence of a • Functional status assessment during the measurement year. medication list in the medical record. Required Exclusion: Transitional care management Members in hospice or using hospice services anytime services during the during the measurement year. measurement year. \*Medical record documentation is **only** accepted for gap Exclude services provided in an closure during the annual HEDIS Medical Record Review. acute inpatient setting. **HEDIS Value** Code Functional Status Assessment -Code Comments Set Name Set At least one functional status Medication List CPT-CAT-II Without a 1P, 2P, 3P, or assessment during the 1159F 8P modifier. Must be measurement year. Exclude used with a Medication services provided in an acute Review code during the inpatient setting. same visit Medication List **HCPCS** G8427 Must be used with a Medication Review code during the same visit Medication **CPT** 90863, 99483, Must be used with a Review 99605, 99606 Medication List code during the same visit Medication CPT-CAT-II 1160F Without a 1P, 2P, 3P, or 8P modifier. Must be Review used with a Medication List code during the same visit **Transitional Care** CPT 99495, 99496 PRODUCT LINE Management Services **DSNP** and MMP CPT-CAT-II Without a 1P, 2P, Functional Status 1170F 3P, or 8P modifier Assessment **Functional Status HCPCS** G0438, G0439 Assessment **Functional Status** CPT 99483 Assessment



## HEDIS® QUALITY MEASURE

## COL-E – Colorectal Cancer Screening

#### **ECDS Measure**

The percentage of members 45 – 75 years of age who had appropriate screening for colorectal cancer.

#### **CLINICAL GOAL**

The member will have one of the following screening tests during the indicated period:

- Fecal occult blood test (FOBT or FIT) between 1/1 and 12/31 of the measurement year.\*\*
- Colonoscopy performed between 1/1 nine (9) years prior to the measurement year and 12/31 of the measurement year.
- Flexible sigmoidoscopy performed between 1/1 four (4) years prior to the measurement year and 12/31 of the measurement year.
- CT colonography performed between 1/1 four (4) years prior to the measurement year and 12/31 of the measurement year.
- Stool DNA (sDNA) with FIT test performed between 1/1 two (2) years prior to the measurement year and 12/31 of the measurement year.

Note: For Stool DNA (sDNA) with FIT test, please verify with customer service the patient's availability of coverage for the test.

\*\* FOBT tests performed on a sample collected from a digital rectal exam do not meet the measure requirements.

#### PRODUCT LINE

Commercial, DSNP, Marketplace, Medicaid and Medicare

#### **CRITERIA TO MEET THE GOAL**

- Submit a claim(s) for any of the following:3 Fecal Occult Blood (FOBT) or 1 Fecal Immunochemical (FIT) test.
- Sigmoidoscopy.
- · CT colonography.
- Stool DNA (sDNA) with FIT test.

#### **Medical Record Documentation of:**

- FOBT (3) done during the measurement year.
- FIT (1) done during the measurement year.
- Colonoscopy done within 9 years prior to the measurement year and 12/31 of the measurement year.
- Flexible sigmoidoscopy done within 4 years prior to the measurement year and 12/31 of the measurement year.
- CT colonography done within 4 years prior to the measurement year and 12/31 of the measurement year.
- FIT-DNA test done within 2 years prior to the measurement year and 12/31 of the measurement year.
- Documentation of exclusion: colorectal cancer or a total colectomy at any time during the member's history through 12/31 of the measurement year.

#### **Required Exclusions:**

- Members receiving palliative care or hospice services during the measurement year.
- Medicare members 66 years of age and older as of 12/31 of the measurement year who are in an I-SNP or living long-term in an institution any time during the measurement year.
- Members 66 of age and older as of 12/31 of the measurement year with frailty and advanced illness.
- Members who had colorectal cancer or a total colectomy any time during the member's history through December 31 of the measurement year (no longer optional).

HEDIS Value Set Name	Code Set	Code	Comments
Colonoscopy	CPT	44388, 44389, 44390, 44391, 44392, 44394, 44401, 44402, 44403, 44404, 44405, 44406, 44407, 44408, 45378, 45379, 45380, 45381, 45382, 45384, 45385, 45386, 45388, 45389, 45390, 45391, 45392, 45393, 45398	
Colonoscopy	HCPCS	G0105, G0121	
CT Colonography	CPT	74261, 74262, 74263	
Flexible Sigmoidoscopy	CPT	45330, 45331, 45332, 45333, 45334, 45335, 45337, 45338, 45340, 45341, 45342, 45346, 45347, 45349, 45350	
Flexible Sigmoidoscopy	HCPCS	G0104	
FOBT Lab Test	CPT	82270, 82274	
FOBT Lab Test	HCPCS	G0328	
sDNA FIT Lab Test	CPT	81528	
Total Colectomy	CPT	44150, 44151, 44152, 44153, 44155, 44156, 44157, 44158, 44210, 44211, 44212	Required Exclusion



HEDIS® QUALITY MEASURE	CLINICAL GOAL	CRITERIA TO MEET THE GOAL
COU – Risk of Continued	Identify members who have a	Claims:
Opioid Use	risk of continued opioid use.	Numerator compliance for this measure is based on pharmacy claims.
Administrative Measure		
The percentage of members 18 years of age and older who have a new episode of opioid use that puts them at risk for continued		≥ 15 Days Covered: Members who had 15 or more calendar days covered by an opioid medication during the 30-day period beginning on the IPSD through 29 days after the IPSD meet numerator compliance.
opioid use. Two rates are reported:  1. The percentage of members with at least 15 days of		≥ 31 Days Covered: Members who had 31 or more calendar days covered by an opioid medication during the 62-day period beginning on the IPSD through 61 days after the IPSD meet numerator compliance.
prescription opioids in a 30-day period.	PRODUCT LINE	Medical record documentation not applicable.
2. The percentage of members with at least 31 days of	Commercial, Medicaid and Medicare	Required Exclusions:
prescription opioids in a 62-day period.		Members who met at least one of the following at any time 1 year prior to the IPSD through 61 days after the IPSD:
		Cancer
<b>Note</b> : A lower rate indicates better performance.		Sickle cell disease
better performance.		Palliative care
Intake Period 11/1 of the year		<ul> <li>Members in hospice or using hospice services any time during the measurement year.</li> </ul>
prior to the measurement year to 10/31 of the measurement year.		<ul> <li>Members who died any time during the measurement year.</li> </ul>
10/01 of the measurement year.		

# **CWP**

HEDIS® QUALITY MEASURE	CLINICAL GOAL		CRITERIA TO MEET THE GOAL	
CWP – Appropriate Testing for Pharyngitis  Administrative Measure  The percentage of members 3 years of age and older who were diagnosed with pharyngitis, dispensed an antibiotic and received a Group A streptococcus (strep) test for the episode in the 7 day period from 3 days prior to the episode date through 3 days after the episode date.	Members 3 years of age and older diagnosed with pharyngitis / tonsillitis must receive a strep test prior to receiving a prescription for antibiotics.	Claims:  Submit a claim for a group A step test with approper Submit a claim for all additional competing diagnor would require antibiotic therapy.  Medical record documentation not applicable.  Required Exclusion:  Members in hospice or using hospice services any during the measurement year.  Note: Other family members with strep and parental are not exclusions to this measure.		umentation not applicable.  : ice or using hospice services anytime arement year.  members with strep and parental refusal
The Intake Period: 7/1 of the prior year to 6/30 of the	PRODUCT LINE Commercial, Marketplace,	HEDIS Value Set Name	Code Set	Code
measurement year.	Medicare and Medicaid	Group A Strep Tests	CPT	87070, 87071, 87081, 87430, 87650, 87651, 87652, 87880



HEDIS® QUALITY MEASURE	CLINICAL GOAL	CRITERIA TO MEET THE GOAL
DAE – Use of High-Risk Medications in Older Adults Administrative Measure	High-risk medications will <b>NOT</b> be prescribed unless absolutely necessary and when appropriate diagnoses are present.	Claims:  Numerator compliance for this measure is based on pharmacy claims.
The percentage of Medicare members 67 years of age and older who had at least two dispensing events for the same high-risk medication.		Required Exclusions:  • Members receiving palliative care or hospice services during the measurement year.
Measure includes:  The percentage of Medicare members 67 years of age and older who had at least two dispensing events for high-risk medications to avoid from the same drug class.  The percentage of Medicare members 67 years of age and older who had at least the dispensing events for		
two dispensing events for high-risk medications to avoid from the same drug class, except for appropriate diagnoses.	PRODUCT LINE  DSNP and Medicare	
The measure reflects potentially inappropriate medication use in older adults, both for medications where any use is inappropriate (Rate 1) and for medications where use under all but specific indications is potentially inappropriate (Rate 2).		
<b>Note</b> : A lower rate represents better performance.		



HEDIS® QUALITY MEASURE	CLINICAL GOAL	CRITERIA TO MEET THE GOAL
DBM-E – Documented Assessment After Mammogram ECDS Measure	Episodes of mammograms will receive a Breast Imaging Reporting and Data System (BI-RADS) scoring category on or within 14 days after the episode date (15 days total).	Mammogram episode is assigned BI-RADS score on or within 14 days after the episode date (15 days total).  Notes: The denominator for this measure is based on episodes, not on members.
Measure Description:	episode date (15 days total).	not on members.
The percentage of episodes of mammograms documented in the form of a BI-RADS assessment within 14 days of the mammogram for members 40 – 74 years of age.		Required Exclusions:  • Members who use hospice services or elect to use a hospice benefit any time during the measurement period
Intake Period:  December 18 of the prior measurement period to December 17 of the measurement period. The intake period is used to capture the episode date.		
	PRODUCT LINE	
	Commercial, Medicaid and Medicare	

DBM-E





HEDIS® QUALITY MEASURE	CLINICAL GOAL	CRITERIA TO MEET THE GOAL
DDE – Potentially Harmful Drug-Disease Interactions in Older Adults Administrative Measure	Potentially harmful prescriptions will <b>NOT</b> be prescribed unless absolutely necessary and when an appropriate diagnosis is present.	Claims:  Numerator compliance for this measure is based on pharmacy claims.  Medical record documentation not applicable.
The percentage of Medicare members 67 years of age and older who have evidence of an underlying disease, condition, or health concern and who were dispensed an ambulatory prescription for a potentially harmful medication, concurrent with or after the diagnosis.  Three rates are reported:  Drug-Disease Interactions: A history of falls and a prescription for anticholinergic agents, antiepileptics, antipsychotics, benzodiazepines, non-benzodiazepine hypnotics or antidepressants (SSRIs, tricyclic antidepressants and SNRIs).		<ul> <li>Required Exclusions:</li> <li>For all three rates: Members receiving palliative care or hospice services during the measurement year.</li> <li>For Rate 1: Diagnosis of psychosis, schizophrenia, schizoaffective disorder, bipolar disorder, major depressive disorder, or seizure disorder on or between January 1 of the year prior to the measurement year and December 1 of the measurement year.</li> <li>For Rate 2: Diagnosis of psychosis, schizophrenia, schizoaffective disorder or bipolar disorder, between January 1 of the year prior to the measurement year and December 1 of the measurement year.</li> </ul>
Drug-Disease Interactions:  Dementia and a prescription for antipsychotics, benzodiazepines, non-benzodiazepine hypnotics, tricyclic antidepressants, or anticholinergic agents.  Drug-Disease Interactions: Chronic kidney disease and prescription for Cox-2 selective NSAIDs or nonaspirin NSAIDs.  Measurement period 1/1 of the year prior to the measurement year and 12/1 of the measurement year.  Note:  • A lower rate indicates better performance for all rates.  • Members with more than one disease or condition may appear in the measure multiple times.	DSNP and Medicare	

#### DDE





HEDIS® QUALITY MEASURE	CLINICAL GOAL		CI	RITERIA TO MEET	THE GOAL	
EED – Eye Exam for Patients with Diabetes  Administrative Measure  Members 18 – 75 years of age with diabetes (Types 1 and 2) who had a retinal eye exam.  Member will have:  • A retinal or dilated eye exam by an eye care professional (optometrist or ophthalmologist) in the measurement year.  • A negative retinal or dilated eye	<ul> <li>A retinal or dilated eye exam by an eye care professional (optometrist or ophthalmologist) in the measurement year.</li> <li>A <u>negative</u> retinal or dilated eye exam (negative for retinopathy) by an eye care professional in the <b>year prior</b> to the</li> </ul>	<ul> <li>Eye exam from year prior to measurement year MUST be negative for numerator compliance.</li> <li>Claims: <ul> <li>If you have or reviewed a report form the patient's ophthalmologist or optomet submit a claim with appropriate CPT II code.</li> </ul> </li> <li>Medical record documentation of one of the following: <ul> <li>A note or letter indicating that an ophthalmoscopic exam was completed by eye care professional, the date when the procedure was performed, and the rest and one of the following: <ul> <li>Evidence that results were read by an eye care professional.</li> <li>Evidence results were read by a qualified reading center that operates under the direction of a medical director who is a retinal specialist.</li> </ul> </li> <li>Evidence results were read by a system that provides an artificial intelligence interpretation.</li> <li>Bilateral eye enucleation or acquired absence of both eyes.</li> <li>Negative retinal or dilated exam by an eye care professional in the year prior</li> </ul> </li> </ul>				
	<ul> <li>the measurement year.</li> <li>Required Exclusions:</li> <li>Members receiving palliative care or hospice services during the measurement year.</li> <li>Medicare members 66 years of age and older as of 12/31 of the measurement year who are in an I-SNP or living long-term in an institution any time during the measurement year.</li> <li>Members 66 of age and older as of 12/31 of the measurement year with frailty and advanced illness.</li> <li>Bilateral eye enucleation anytime during the member's history through December 31 of the measurement year.</li> </ul>					
		HEDIS Value Set Name	Code Set	Code	Comments	
		Retinal Eye Exams  Retinal Eye Exams	CPT	92002, 92004, 92012, 92014, 92018, 92019, 92134, 92201, 92202, 92227, 92228, 92230, 92235, 92250, 99203, 99204, 99205, 99213, 99214, 99215, 99242, 99243, 99244, 99245 S0620, S0621, S3000	Billed by an eye care professional (optometrist or ophthalmologist) during the measurement year OR during the year prior to the measurement year, with a diagnosis of diabetes without complications  Billed by an eye care professional (optometrist or ophthalmologist) during the	
		Diabetes Mellitus Without Complications	ICD10M	E10.9, E11.9, E13.9	measurement year OR during the year prior to the measurement year, with a diagnosis of diabetes without complications  To be used with appropriate visit code for numerator compliance	
		Direct Reference Code (formally Automated Eye Exam)	CPT	92229	Billed by any provider type during the measurement year	
		Eye Exam With Evidence of Retinopathy	CPT-CAT-II	2022F, 2024F, 2026F	Without a 1P, 2P, 3P, or 8P modifier and during the measurement year	
		Eye Exam Without Evidence of Retinopathy	CPT-CAT-II	2023F, 2025F, 2033F	Without a 1P, 2P, 3P, or 8P modifier, and during the measurement year or the year prior to the measurement year	
		Direct Reference Code (formally Diabetic Retinal Screening Negative In Prior Year)	CPT-CAT-II	3072F	Without a 1P, 2P, 3P, or 8P modifier, and during the measurement year	
		Retinal Imaging	CPT	92227, 92228	During the measurement year	



HEDIS® QUALITY MEASURE	CLINICAL GOAL	CRITERIA TO MEET THE GOAL
FMA-E – Follow-up After Abnormal Mammogram Assessment  ECDS Measure  Measure Description: The percentage of episodes for members 40 – 74 years of age with inconclusive or high-risk BI-RADS assessments who received appropriate follow-up within 90 days of the assessment.  Intake Period: October 3 of the year prior to the measurement period to October 2 of the measurement period. The intake period is used to capture the episode date.	CLINICAL GOAL  Appropriate and timely follow up for members whose breast cancer screening was classified as High-Risk (Category 4 or 5) or Incomplete (Category 0) with the Breast Imaging Reporting and Data System (BI-RADS) reporting system. Follow up includes:  • Category 0 – Incomplete: Received an additional screening (mammogram or ultrasound) on or within 90 days after the episode date (91 days total).  • Category 4 – Suspicious  OR  Category 5 – Highly Suggestive of Malignancy: Received a breast biopsy on or within 90 days after the episode date (91 days total).  PRODUCT LINE  Commercial, Medicaid and Medicare	Claims:  • Submit a claim for services completed with appropriate codes.  Notes:  • A higher rate indicates better performance  • The denominator of this measure is based on episodes, not on members.  Required Exclusions:  • Members who use hospice services or elect to use a hospice benefit any time during the measurement period.



HEDIS® QUALITY MEASURE	CLINICAL GOAL	CRITERIA TO MEET THE GOAL					
FMC – Follow-Up After Emergency Department Visit for People with Multiple High-Risk Chronic Conditions	There will be a follow-up service within 7 days after the ED visit (8 days total). ED visits resulting in an inpatient stay are excluded.	Visits that will meet follow-up criteria:  • Outpatient					
Administrative Measure	Eligible chronic conditions:						
The percentage of emergency department (ED) visits for members 18 years of age and older who have multiple highrisk chronic conditions who had a follow-up service within 7 days of the ED visit.  The denominator for this measure is based on ED visits, not on members. All eligible ED visits are included on or between 1/1 – 12/1 of the measurement year.	COPD and Asthma, Alzheimer's disease and related disorders, Chronic kidney disease, Depression, Heart Failure, Acute myocardial infarction, Atrial Fibrillation, Stroke and Transient Ischemic Attack	Telephone E-visit or virtua Telehealth Transitional ca Case managen Complex care Outpatient or Intensive outpa Community m Electroconvuls Substance use Substance abu Medical record of	е				
				поп пот аррисавіе.			
	PRODUCT LINE	Required Exclusion:  • Members in hospice or using hospice services anytime during the measurement year.					
	Medicare	HEDIS Value Set Name	Code Set	Code	Comments		
		Transitional Care Management Services	CPT	99495, 99496			
		Case Management Encounter	CPT	99366			
		Case Management Encounter	HCPCS	T1016, T1017, T2022, T2023			
		Complex Care Management Services	CPT	99439, 99487, 99489, 99490, 99491			
		Complex Care Management Services	HCPCS	G0506			
		Remaining HED	ilS Value Se	et codes can be four	nd in Appendix 7.		



HEDIS® QUALITY MEASURE	CLINICAL GOAL	CRITERIA TO MEET THE GOAL				
FUA – Follow-Up After Emergency Department Visit for Substance Use  Administrative Measure  The percentage of emergency department (ED) visits among members age 13 years and older with a principal diagnosis of Substance Use Disorder (SUD), or any diagnosis of drug overdose, for which there was follow-up.	7-Day Follow-up: Members who had an ED visit for substance use disorder or drug overdose will have a follow-up visit or have a pharmacotherapy dispensing event within 7 days of ED visit.  30-Day Follow-up: Members who had an ED visit for substance use disorder or drug overdose will have a follow-up visit or have a pharmacotherapy dispensing	for a follow days after of Visits, when or events that • Outpatient • Intensive o • Non-reside • Communit • Peer support	laim, w y-up ser dischard paired at will n utpatie intial su y menta	ith the appropriate rvice within 7 days ge.  with appropriate defined follow-up crite on the counter or particular the cou	provider and coding, after discharge and 30 liagnosis / provider, eria: rtial hospitalization atment facility	
Two rates are reported: 7-Day Follow-Up: The percentage of ED visits for which the member received follow-up within 7 days of the ED visits (8 total days).  30-Day Follow-Up: The percentage of ED visits for	event within 30 days of ED visit.	surveillanc  • Behavioral	abuse ( e. health	disorder service, co screening or asses	-	
which the member received	PRODUCT LINE Commercial, Medicaid	Medical record documentation not applicable.				
follow-up within 30 days of the ED visits (31 total days).  The denominator for this measure is based on ED visits, not on members. All eligible ED visits are included on or between 1/1 – 12/1 of the measurement year.	Commercial, Medicaid and Medicare	during the  HEDIS Value Set Name  Visit Setting Unspecified	Code Set	Code  90791, 90792, 90832, 90833, 90834, 90836, 90837, 90838, 90839, 90840, 90845, 90847, 90849, 90853, 90876, 99221, 99222, 99223, 99231, 99232, 99233, 99252, 99253, 99254, 99255	Comments  With either a mental health provider OR a diagnosis of SUD, substance use, or drug overdose AND one of the following:  Outpatient POS code POS code 52 POS code 53 Non-residential substance abuse treatment facility POS code Telehealth POS code	



FUH					<b>+</b> ♥NC
HEDIS® QUALITY MEASURE	CLINICAL GOAL		CRITE	RIA TO MEET TI	HE GOAL
FUH – Follow-Up After Hospitalization for Mental Illness  Administrative Measure  The percentage of discharges for members 6 years of age and older who were hospitalized for a principle diagnosis of mental illness, or any diagnosis of intentional self-harm, and had a mental health follow-up service.  Two rates are reported: 7-Day Follow-Up: The percentage of discharges for which the member received follow-up within 7 days after discharge. 30-Day Follow-Up: The percentage of discharges for which the member received follow-up within 30 days after discharge.	7-Day Follow-up: Members who were hospitalized for select mental illnesses or intentional self-harm diagnoses will have a mental health follow-up service within 7 days after discharge.  30-Day Follow-up: Members who were hospitalized for select mental illnesses or intentional self-harm diagnoses will have a mental health follow-up service within 30 days after discharge.  Note: Follow up with a PCP does not meet the measure. The visit must be with a mental health practitioner. Do not include services that occur on the date of discharge.	Claims:  Submit a claim, with the appropriate provider and conformation for a follow-up service within 7 days after discharge days after discharge.  Visits, when paired with appropriate diagnosis / provided that will meet follow-up criteria:  Outpatient Intensive outpatient encounter or partial hospitalization.  Community mental health center visit  Electroconvulsive therapy			
The denominator for this measure is based on discharges,		HEDIS Value Set Name	Code Set	Code	Comments
not on members. All discharges are included on or between 1/1 to 12/1 of the measurement year.	PRODUCT LINE  Commercial, DSNP, Marketplace, Medicare and Medicaid	Visit Setting Unspecified	CPT	90791, 90792, 90832, 90833, 90834, 90836, 90837, 90838, 90839, 90840, 90845, 90875, 90876, 99221, 99222, 99223, 99231, 99232, 99233, 99254, 99255	With one of the following:  Outpatient POS code with a mental health provider  Mental health disorder diagnosis  POS code 52  POS code 53  POS code 56  Telehealth POS code and with a mental health provider
		Partial Hospitalization or Intensive Outpatient  Remaining F	HCPCS	G0410, G0411, H0035, H2001, H2012, S0201, S9480, S9484, S9485	Electroconvulsive     Therapy  be found in Appendix 7.



HEDIS® QUALITY MEASURE	CLINICAL GOAL	CRITERIA TO MEET THE GOAL
FUI- Follow-Up After High-Intensity Care for Substance Use Disorder  Administrative Measure  The percentage of acute inpatient hospitalizations, residential treatment or withdrawal management visits for a diagnosis of substance use disorder among members 13 years of age and older that result in a follow-up visit or service for substance use disorder.  Two rates are reported: 7-Day Follow-Up: The percentage of visits or discharges for which the member received follow-up for substance use disorder with in the 7 days after the visit or discharge.  30-Day Follow-Up: The percentage of visits or discharges for which the member received follow-up for substance use disorder with in the 30 days after the visit or discharge.  The denominator for this measure is based on episodes, not on members. Include all episodes on or between 1/1 to12/1 of the measurement year.	7-Day Follow-up:  Members will have a follow-up visit or event with any practitioner for a principle diagnosis of substance use disorder within the 7 days after an episode for substance use disorder.  30-Day Follow-up:  Members will have a follow-up visit or event with any practitioner for a principle diagnosis of substance use disorder within the 30 days after an episode for substance use disorder.  Note: Do not include visits that occur on the date of the denominator episode.  PRODUCT LINE  Commercial, Medicaid and Medicare	Claims:  Submit a claim, with the appropriate coding, for a follow-up service within 7 days after discharge and 30 days after discharge or visit.  Visits, when paired with appropriate diagnosis / provider, or events that will meet follow-up criteria:  Acute or nonacute inpatient admission or residential behavioral health stay on the D/C claim  Outpatient  Intensive outpatient encounter or partial hospitalization  Non-residential substance abuse treatment facility  Community mental health center  Telehealth  Substance use disorder service  Opicial treatment service that bills monthly or weekly  Residential behavioral health treatment  Telephone  E-visit or virtual check-in  Pharmacotherapy dispensing event  Acute or nonacute inpatient admission or residential behavioral health stay  Note: Follow-up does not include withdrawal management.  Required Exclusion:  Members in hospice or using hospice services any time during the measurement year.  HEDIS Value Set codes can be found in Appendix 7.



Administrative Measure  The percentage of emergency department (ED) visits for members 6 years of age and older with a principal diagnosis of mental illness or any diagnosis of intentional self-harm, and had a mental health follow-up service.  The percentage of emergency department (ED) visits for members 6 years of age and older with a principal diagnosis of mental illness or any diagnosis of intentional self-harm, and had a mental health follow-up service.  The percentage of emergency department all illnesses or intentional self-harm diagnoses will have a mental health follow-up service within 7 days after ED visit and 30 days after ED visit.  Visits, when paired with appropriate diagnosis provider, that will meet follow-up criteria:  Outpatient  Intensive outpatient encounter or partial hospitalization  Community mental health center	HEDIS® QUALITY MEASURE	CLINICAL GOAL	CRITERIA TO MEET THE GOAL
The percentage of ED visits for which the member received follow-up within 7 days of the ED visit (8 total days)  The percentage of ED visits for which the member received follow-up within 30 days after the ED visit.  The percentage of ED visits for which the member received follow-up within 30 days of the ED visit (31 total days)  The denominator for this measure is based on ED visits, not on members.  All eligible ED visits are included on or between 1/1 – 12/1 of the measurement year.  The percentage of ED visits for which the within 30 days after the ED visit.  The percentage of ED visits for which the within 30 days after the ED visit.  The denominator for this measure is based on ED visits, not on members.  All eligible ED visits are included on or between 1/1 – 12/1 of the measurement year.  The denominator for this measure is mental health follow-up service within 30 days after the ED visit.  Telehealth  Telephone  E-visit or virtual check-in  Behavioral healthcare setting  Peer support services  Psychiatric residential treatment  Medical record documentation not applicable.  Required Exclusion:  Members in hospice or using hospice service anytime during the measurement year.	FUM – Follow-Up After Emergency Department Visit for Mental Illness  Administrative Measure  The percentage of emergency department (ED) visits for members 6 years of age and older with a principal diagnosis of mental illness or any diagnosis of intentional self-harm, and had a mental health follow-up service.  Two rates are reported: The percentage of ED visits for which the member received follow-up within 7 days of the ED visit (8 total days)  The percentage of ED visits for which the member received follow-up within 30 days of the ED visit (31 total days)  The denominator for this measure is based on ED visits, not on members.  All eligible ED visits are included on or between 1/1 – 12/1 of the measurement	7-Day Follow-up:  Members who had an ED visit for a principle diagnosis of mental illnesses or intentional self-harm diagnoses will have mental health follow-up servic within 7 days after the ED visit for a principle diagnosis of mental illnesses or intentional self-harm diagnoses will have mental health follow-up servic within 30 days after the ED visit for a principle diagnosis of mental illnesses or intentional self-harm diagnoses will have mental health follow-up servic within 30 days after the ED visit for a principle diagnosis of mental illnesses or intentional self-harm diagnoses will have mental health follow-up servic within 30 days after the ED visit for a principle diagnosis of mental health follow-up servic within 30 days after the ED visit for a principle diagnosis of mental illnesses or intentional self-harm diagnoses will have mental health follow-up servic within 30 days after the ED visit for a principle diagnosis of mental illnesses or intentional self-harm diagnoses will have mental health follow-up servic within 30 days after the ED visit for a principle diagnosis of mental illnesses or intentional self-harm diagnoses will have mental health follow-up servic within 30 days after the ED visit for a principle diagnosis of mental illnesses or intentional self-harm diagnoses will have mental health follow-up servic within 30 days after the ED visit for a principle diagnosis of mental illnesses or intentional self-harm diagnoses will have mental health follow-up servic within 30 days after the ED visit for a principle diagnosis of mental illnesses or intentional self-harm diagnoses will have mental health follow-up servic within 30 days after the ED visit for a principle diagnosis of mental illnesses or intentional self-harm diagnoses will have mental health follow-up servic within 30 days after the ED visit for a principle diagnosis of mental illnesses or intentional self-harm diagnoses will have mental health follow-up servic within 30 days after the ED visit for a principle diagnosis of mental	Claims:  Submit a claim, with the appropriate coding, for a follow-up service within 7 days after ED visit and 30 days after ED visit.  Visits, when paired with appropriate diagnosis / provider, that will meet follow-up criteria:  Outpatient  Intensive outpatient encounter or partial hospitalization  Community mental health center  Electroconvulsive therapy  Telehealth  Telephone  E-visit or virtual check-in  Behavioral healthcare setting  Peer support services  Psychiatric residential treatment  Medical record documentation not applicable.  Required Exclusion:  Members in hospice or using hospice services

**GSD** 



#### **GSD - Glycemic Status Assessment for Patients with Diabetes**

**Hybrid Measure** – The percentage of members 18 – 75 years of age with diabetes (types 1 and 2) whose most recent glycemic status (hemoglobin A1c [HbA1c] or glucose management indicator [GMI] was at the following levels dring the measurement year:

• Glycemic Status < 8.0%

• Glycemic Status > 9.0%

HEDIS® QUALITY MEASURE	CLINICAL GOAL		CRITER	IA TO MEET T	THE GOAL		
Glycemic Status < 8.0%. Identify the most recent glycemic status assessment (HbA1c or GMI) during the measurement year. The member is numerator compliant if the most recent glycemic status assessment has a result of < 8.0%. The member is not numerator compliant if the result of the most recent glycemic status assessment is ≥ 8.0% or is missing a result, or if a glycemic status assessment was not done during the measurement year.	Members glycemic status (HbA1c or GMI test) remains < 8% during the measurement year.  PRODUCT LINE  Commercial, Marketplace, Medicaid and Medicare	Medical Record     Date and value measuremen     If there are measuremen     Members record measuremen     Medicare measuremen institution an	I Document ie of most re t year. ultiple glyce ne lowest re sions: eiving pallie t year. mbers 66 ye t year who y time durir years or age	ecent HbA1C or emic status assessants.  ative care or hose are of age and are in an I-SNP on the measurer e and older as o	GMI test result of the same date of spice services during the older as of 12/31 of the or living long-term in an		
		HEDIS Value Code Code Comments					
		HbA1c Lab Test	CPT	83036, 83037	Most recent and with CPT-CAT-II code (without a 1P, 2P, 3P, or 8P modifier, or POS code 81).		
		HbA1c Lab Test	LOINC	17855-8, 17856-6, 4548-4, 4549-2, 96595-4	Most recent and with CPT-CAT-II code (without a 1P, 2P, 3P, or 8P modifier, or POS code 81).		
		HbA1c Testing Result or Finding	CPT-CAT-II	3046F	Most recent hemoglobin A1c level Greater Than 9.0% (Numerator Non-Compliant).		
		HbA1c Testing Result or Finding	CPT-CAT-II	3052F	Most recent hemoglobin A1c (HbA1c) level Greater Than or Equal to 8.0% and Less Than or Equal to 9.0% (Numerator Non-Compliant).		
		HbA1c Testing Result or Finding	CPT-CAT-II	3044F	Most recent hemoglobin A1c (HbA1c) level Less Than 7.0% (Numerator Compliant).		
		HbA1c Testing Result or Finding	CPT-CAT-II	3051F	Most recent hemoglobin A1c (HbA1c) level Greater Than or Equal to 7.0% and Less Than 8.0% (Numerator Compliant).		
		Direct Reference Code (Glucose Management Indicator)	LOINC	97506-0	The CPT-CAT-II Codes listed above only apply to the HbA1c Lab Test, not the Glucose Management Indicator.		

**GSD** 



#### **GSD - Glycemic Status Assessment for Patients with Diabetes**

**Hybrid Measure** – The percentage of members 18 – 75 years of age with diabetes (types 1 and 2) whose most recent glycemic status (hemoglobin A1c [HbA1c] or glucose management indicator [GMI] was at the following levels dring the measurement year:

• Glycemic Status < 8.0%

• Glycemic Status > 9.0%

Measurement year 1/1 to 12/31 of the measurement year.

HEDIS® QUALITY MEASURE	CLINICAL GOAL		CRITERIA.	TO MEET TH	IE GOAL
Glycemic Status > 9.0%. Identify the most recent glycemic status assessment (HbA1c or GMI) during the measurement year. The member is numerator compliant if the most recent glycemic status assessment has a result of > 9.0% or is missing a result, or if a glycemic status assessment was not done during the measurement year. The member is not numerator compliant if the result of the most recent glycemic status assessment is ≤ 9.0%.	Members glycemic status (HbA1c or GMI test) remains ≤ 9% during the measurement year.  PRODUCT LINE  Commercial, Marketplace, Medicaid and Medicare	measurement y  If there are mult service, use the Required Exclusion  Members receive measurement y  Medicare members measurement y institution any terms.	cocumentation of most received ar. liple glycem lowest resurces. It wing palliative ar. liple see the composition of the correction of the	ion of: ent HbA1C or G ic status assess ilts.  ve care or hosp rs of age and o e in an I-SNP or the measurem and older as of	MI test result of the sments on the same date of pice services during the lder as of 12/31 of the r living long-term in an
Note: This is an inverse measure rate. A lower re-		HEDIS Value Set Name	Code Set	Code	Comments
ported rate indicates better performance for this indi- cator (i.e., low rates of Gly- cemic Status > 9% indicate		HbA1c Lab Test	CPT	83036, 83037	Most recent and with CPT-CAT-II code (without a 1P, 2P, 3P, or 8P modifier).
better care). However, on reporting given to Providers Blue Cross NC chooses		HbA1c Lab Test	LOINC	17855-8, 17856-6, 4548-4, 4549-2, 96595-4	Most recent and with CPT-CAT-II code (without a 1P, 2P, 3P, or 8P modifier).
to reflect this rate as the actual clinical goal of ≤ 9% to align reporting of both Glycemic Status rates and		HbA1c Test Result or Finding	CPT-CAT-II	3046F	Most recent hemoglobin A1c level Greater Than 9.0% (Numerator Compliant).
reduce confusion of the clinical goal.		HbA1c Testing Result or Finding	CPT-CAT-II	3052F	Most recent hemoglobin A1c (HbA1c) level Greater Than or Equal to 8.0% and Less Than or Equal to 9.0% (Numerator Non-Compliant).
		HbA1c Testing Result or Finding	CPT-CAT-II	3044F	Most recent hemoglobin A1c (HbA1c) level Less Than 7.0% (Numerator Non-Compliant).
		HbA1c Testing Result or Finding	CPT-CAT-II	3051F	Most recent hemoglobin A1c (HbA1c) level Greater Than or Equal to 7.0% and Less Than 8.0% (Numerator Non-Compliant).
		Direct Reference Code (Glucose Management Indicator)	LOINC	97506-0	The CPT-CAT-II Codes listed above only apply to the HbA1c Lab Test, not the Glucose Management Indicator.





HEDIS® QUALITY MEASURE	CLINICAL GOAL	CRITERIA TO MEET THE GOAL
HDO – Use of Opioids at High Dosage	Assesses potentially high-risk opioid analgesic prescribing practices.	Members whose average MME was ≥ 90 meet the numerator criteria.
Administrative Measure		Medical record documentation not applicable.
The percentage of members 18 years of age and older who received prescription opioids at a high dosage (average morphine milligram equivalent dose [MME] ≥ 90) for ≥ 15 days during the measurement year.  Note: A lower rate indicates better performance.	PRODUCT LINE  Commercial, Medicaid and Medicare	Required Exclusions:  Cancer Sickle cell disease Palliative care Members in hospice or using hospice services.

# **IET**

HEDIS® QUALITY MEASURE	CLINICAL GOAL	CRITERIA TO MEET THE GOAL
IET- Initiation and Engagement of Substance Use Disorder Treatment	Initiation phase – Members will have a follow-up visit within 14 days of the SUD diagnosis.	Claims: Submit a claim, with the appropriate coding, for follow-up services within the appropriate time frame.
Administrative Measure  New Substance Use Disorder (SUD) episodes that result in treatment initiation and engagement.  Ages 13 and older as of the SUD episode date.  Two rates are reported: Initiation of SUD Treatment: The percentage of new SUD episodes that result in treatment initiation through inpatient SUD admission, outpatient visit, intensive outpatient encounter, partial hospitalization, telehealth visit or medication treatment within 14 days.	Continuation phase – Members will have at least two of the following on the day after the initiation encounter through 34 days after the initiation event:  • Engagement visit.  • Engagement medication treatment event.  Note: If a member is noncompliant with Initiation within 14 days the member is then non-compliant for both Initiation and Engagement.	Visits, when paired with appropriate diagnosis/coding, or events that will meet follow-up criteria:  • An acute or nonacute inpatient admission with diagnosis of alcohol abuse and dependence, opioid abuse and dependence and other drug abuse and dependence.  • An outpatient visit.  • Intensive outpatient encounter or partial hospitalization.  • A non-residential substance abuse treatment facility visit.  • A community mental health center visit.  • A telehealth visit.  • Substance use disorder service.  • Telephone visit.  • E-visit or virtual check-in.  • An acute or nonacute inpatient.  • Inpatient stay for medically managed withdrawal.  Either of the following meets criteria for a medication treatment event:
Engagement of SUD Treatment: The percentage of new SUD	PRODUCT LINE	<ul> <li>Alcohol use disorder and alcohol use disorder medication treatment dispensing event.</li> <li>Opioid use disorder medication treatment dispensing event.</li> </ul>
episodes that have evidence of treatment engagement within 34 days of initiation.	Commercial, Marketplace, Medicare and Medicaid	Medical record documentation not applicable.  Required Exclusion:
The Intake Period: 11/15 of the year prior to the measurement year to 11/14 of the measurement year.		Members in hospice or using hospice services anytime during the measurement year.  HEDIS Value Set codes can be found in Appendix 7.

#### HDO – IET

IMA-E



HEDIS® QUALITY MEASURE	CLINICAL GOAL	CRITERIA TO MEET THE GOAL				
IMA-E – Immunizations for Adolescents  ECDS Measure  The percentage of adolescents 13 years of age who had the vaccinations listed under the Clinical Goal column.	By the 13th birthday, members will have received:  1 meningococcal vaccine on or between 10th and 13th birthdays.  1 Tdap vaccine on or between 10th and 13th birthdays.  2 or 3 HPV vaccines with different dates of service on or between the member's 9th and 13th birthdays.  If reporting only 2 vaccines, there must be at least 146 days between the first and second dose of the HPV vaccine.  Combo 1 = Numerator compliant for both meningococcal and Tdap.  Combo 2 = Numerator compliant for all.	<ul> <li>Report all Immunizations Registry.</li> <li>Required Exclusion:</li> <li>Members in hospice or unduring the measurement</li> <li>Medical record documenta</li> <li>A visit note indicating the the date of the immunization</li> <li>A certificate of immunization health care provider or a and types of immunization</li> <li>Note:         <ul> <li>A member is numerator construction</li> <li>Anaphylaxis due to the repertussis or HPV vaccine 13th birthday meets critered in the total construction</li> <li>Encephalitis due to the total construction</li> </ul> </li> </ul>	for all vaccinations members receive. unizations to the North Carolina Immunizatio  on: spice or using hospice services anytime surement year.  ocumentation of: icating the name of the specific antigen and immunization. immunization prepared by an authorized vider or agency, including the specific dates imunizations administered.			
		HEDIS Value Set Name	Code Set	Code		
		Meningococcal Immunization	CVX	32, 108, 114, 136, 147, 167, 203, 316		
	PRODUCT LINE	Meningococcal Vaccine Procedure	СРТ	90619, 90733, 90734, 90623		
	Commercial, Marketplace and Medicaid	Direct Reference Code (formally Tdap Immunization)	CVX	115		
		Tdap Vaccine Procedure	CPT	90715		
		HPV Immunization	CVX	62, 118, 137, 165		
		HPV Vaccine Procedure	CPT	90649, 90650, 90651		





HEDIS® QUALITY MEASURE	CLINICAL GOAL	CRITERIA TO MEET THE GOAL					
KED – Kidney Health Evaluation for Patients with Diabetes  Administrative Measure  Members 18 – 85 years of age with diabetes (Type 1 and Type 2) who received a kidney health evaluation during the measurement year.	<ul> <li>Claims:</li> <li>Submit a claim for eGFR AND both a quantitative urine albumin test and a urine creatinine test OR a urine albumin creatinine ratio lab test.</li> <li>Medical record documentation not applicable.</li> <li>Required Exclusions:</li> <li>Members who had dialysis any time during the member's history on or prior to December 31 of the measurement year.</li> <li>ESRD</li> <li>Members receiving palliative care or hospice services during the measurement year.</li> <li>Medicare members 66-80 years of age as of 12/31 of the measurement year who are in an I-SNP or living longterm in an institution any time during the measurement year.</li> <li>Members 66 years of age and older as of 12/31 of the measurement year with frailty and advanced illness.</li> <li>Members 81 years of age and older as of 12/31 of the measurement year with frailty during the measurement year.</li> </ul>						
	PRODUCT LINE  Commercial, Marketplace, Medicare and Medicaid	HEDIS Value Set Name	Code Set	Code	Comments		
Commercia		Estimated Glomerular Filtration Rate Lab Test	CPT	80047, 80048, 80050, 80053, 80069, 82565			
		Quantitative Urine Albumin Lab Test	CPT	82043	With urine creatinine test (with service dates four days or less apart)		
		Urine Creatinine Lab Test	CPT	82570	With quantitative urine albumin test (with service dates four days or less apart)		
		Urine Albumin Creatinine Ratio Lab Test	LOINC	13705-9, 14958-3, 14959-1, 30000-4, 44292-1, 59159-4, 76401-9, 77253-3, 77254-1, 89998-9, 9318-7			

## LBP



HEDIS® QUALITY MEASURE	CLINICAL GOAL	CRITERIA TO MEET THE GOAL						
			UNITEN	IA TO WILLT THE GOAL				
LBP – Use of Imaging Studies for Low Back Pain	Members 18 – 75 years will <b>NOT</b> receive imaging studies within 28 days of the initial diagnosis	<ul> <li>Claims:</li> <li>Submit a claim with a code for low back pain with first symptom of low back pain.</li> </ul>						
Administrative Measure	of low back pain. This includes a plain X-ray.			entation not applicable.				
The percentage of members  18 – 75 years of age with a principal diagnosis of low back pain who did <b>NOT</b> have an imaging study (plain X-ray,	<b>Note</b> : Consider referral for physical therapy evaluation before X-rays are ordered.	Required Exclusions*:  • Cancer or a history of cancer  • Fragility fracture  • HIV						
MRI, CT scan) within 28 days of the diagnosis.		Hospice     IV drug abus						
The Intake Period: 1/1 to 12/3 of the measurement year.		<ul><li>Lumbar surg</li><li>Major organ</li><li>Neurologic i</li></ul>	transpla					
This measure is reported as an		Osteoporosi						
inverted rate. A higher score indicates appropriate treatment		<ul><li>Palliative car</li><li>Prolonged u</li></ul>		rticosteroids				
of low back pain.		Recent traur						
	PRODUCT LINE	Spinal infect     Spandylana						
	Commercial, Marketplace, Medicaid and Medicare	Spondylopathy     * Please see specification for the time frame for an exclusion						
		HEDIS Value Set Name	Code Set	Code	Comments			
		Imaging Study	CPT	72020, 72052, 72100, 72110, 72114, 72120, 72131, 72132, 72133, 72141, 72142, 72146, 72147, 72148, 72149, 72156, 72158, 72200, 72202, 72220, 72040, 72050, 72070, 72072, 72074, 72080, 72081, 72082, 72083, 72084, 72125, 72126, 72127, 72128, 72129, 72130, 72157	Must be used along with a diagnosis of uncomplicated low back pain			
		Uncomplicated Low Back Pain	ICD10CM	M47.26, M47.27, M47.28, M47.816, M47.816, M47.817, M47.818, M47.896, M47.897, M47.898, M48.061, M48.07, M48.08, M51.16, M51.17, M51.26, M51.27, M51.36, M51.37, M51.86, M51.87, M53.2X6, M53.2X7, M53.2X8, M53.3, M53.86, M53.87, M53.88, M54.16, M54.17, M54.18, M54.30, M54.31, M54.32, M54.40, M54.41, M54.42, M54.5, M54.50, M54.51, M54.59, M54.89, M54.9, M99.03, M99.04, M99.23, M99.33, M99.43, M99.53, M99.63, M99.73, M99.83, M99.84, S33.100A, S33.100D, S33.110S, S33.120A, S33.120D, S33.120S, S33.120A, S33.120D, S33.120S, S33.140A, S33.140D, S33.140S, S33.140A, S33.140D, S33.140S, S33.140A, S33.140D, S33.140S, S33.140A, S33.140D, S33.140S, S33.5XXA, S33.6XXA, S33.8XXA, S33.9XXA, S39.002A, S39.002D, S39.002S, S39.012A, S39.092D, S39.092S, S39.82XA, S39.82XD, S39.92XS, S39.92XA, S39.92XD, S39.92XS, S39.92XA, S39.92XD, S39.92XS	Must be used along with an imaging study			





HEDIS® QUALITY MEASURE	CLINICAL GOAL	CRITE	RIA TO N	MEET THE GOAL
OMW – Osteoporosis Management in Women Who Had a Fracture Administrative Measure	Members 67 – 85 years of age, who had a fracture, will have a BMD test or a prescription for a drug to treat osteoporosis within 6 months of the date of fracture.		ance for me	ng within 6 months of a fracture. edication to treat osteoporosis is
The percentage of women 67 – 85 years of age who suffered a fracture and had either a bone mineral density (BMD) test or prescription for a drug to treat osteoporosis in the six months after the fracture.  The Intake Period: 7/1 of the prior year to 6/30 of the measurement year.	Note: Consider BMD testing every 2 years in this age group.  PRODUCT LINE  DSNP and Medicare	<ul> <li>Medical Record documentation:</li> <li>A visit note with evidence of BMD test in the appropriate timeframe.</li> <li>A visit note with evidence of fill or dispense date of medication given within the appropriate timeframe.</li> <li>Required Exclusions:</li> <li>Members receiving palliative care or hospice services during the measurement year.</li> <li>Medicare members 67 years of age and older as of 12/31 of the measurement year who are in an I-SNP or living long-term in an institution any time during the measurement year.</li> <li>Members 67 – 80 years of age and older as of 12/31 of the measurement year with frailty and advanced illness.</li> <li>Members 81 years of age and older as of 12/31 of the measurement year with frailty.</li> </ul>		
		HEDIS Value Set Name	Code Set	Code
		Bone Mineral Density Tests	CPT	76977, 77078, 77080, 77081, 77085, 77086
		Long-Acting Osteoporosis Medications	HCPCS	J0897, J1740, J3489
		Osteoporosis Medication Therapy	HCPCS	J0897, J1740, J3110, J3111, J3489





## PBH

HEDIS® QUALITY MEASURE	CLINICAL GOAL	CRITI	ERIA TO MEET THE GO	AL
PBH – Persistence of Beta-Blocker Treatment After a Heart Attack	Members 18 years of age and older with a new diagnosis of AMI will remain on	Claims:  Numerator compliance for this measure is based on pharmacy claims.  At least 135 days of treatment with beta-blockers during the 180-day measurement interval.  Medical record documentation not applicable.		
Administrative Measure  The percentage of members 18 years of age and older who were hospitalized and discharged with a diagnosis of AMI and who remained on beta-blocker treatment for six months after discharge.  The Intake Period: 7/1 of the prior year to 6/30 of the measurement year.  beta-blocker treatment for six months after the hospital discharge.  Note: Consider 90 day supply or refills x 6 if appropriate.		Required Exclusions:  Asthma Hypotension Heartblock > 1 degree Sinus bradycardia COPD Chronic respiratory due to inhaled fumes / vapors Intolerance or allergy to beta-blocker therapy Members in hospice or using hospice services anytime during the measurement year. Medicare members 66 years of age and older as of 12/31 of the measurement year who are in an I-SNP or living long-term in an institution any time during the measurement year. Members 66 – 80 years of age and older as of 12/31 of the measurement year with frailty and advanced illness. Members 81 years of age and older as of 12/31 of the measurement year with frailty. Member with a medication dispensing event that indicates contraindication to		
	Commercial, DSNP, Medicare and Medicaid	continuous enrollment perio Required Asthma Medication  Description  Bronchodilator combinations  Inhaled corticosteroids	Prescription  Budesonide-formoterol  Fluticasone-vilanterol  Beclomethasone	Fluticasone-salmeterol Formoterol-mometasone Flunisolide
			Budesonide Ciclesonide	Fluticasone  Mometasone



HEDIS® QUALITY MEASURE	CLINICAL GOAL	CRITERIA TO MEET THE GOAL
PCE – Pharmacotherapy Management of COPD Exacerbation	Member will have a dispensed prescription for an appropriate systemic corticosteroid on or 14 days after the discharge date	Claims:     Numerator compliance for this measure is based on pharmacy claims.
Administrative Measure	and a bronchodilator on or 30 days after the discharge date.	Medical record documentation not applicable.
The percentage of COPD exacerbations for members 40 years of age and older who had an acute inpatient discharge or ED visit and were dispensed appropriate medications.		Required Exclusion:  Members in hospice or using hospice services anytime during the measurement year.  Note: Measure is based on acute inpatient discharges and ED visits, not on members – denominator can include multiple
Two rates are reported:	PRODUCT LINE	events for the same member.
<ol> <li>Systemic corticosteroid dispensed within 14 days of discharge date.</li> <li>Bronchodilator dispensed on or within 30 days of discharge date.</li> <li>The Intake Period: 1/1 to 11/30 of the measurement year.</li> </ol>	Commercial, DSNP, Medicare and Medicaid	

# PCR

HEDIS® QUALITY MEASURE	CLINICAL GOAL	CRITERIA TO MEET THE GOAL	
PCR - Plan All-Cause Readmissions For members 18 years	Members will not have unnecessary / avoidable readmissions.	This is a risk adjusted measure that calculates 30 day readmission rates utilizing the following components:  Number of discharges – denominator  Number of 30-day readmissions – numerator	
of age and older, the number of acute inpatient and	PRODUCT LINE	<ul> <li>Rate = (numerator / expected number of readmissions)</li> <li>Global rate</li> <li>Expected readmission probability is calculated with each discharge</li> </ul>	
observation stays during the measurement year that were followed by	Commercial, DSNP, Marketplace, Medicare and Medicaid	Medical record documentation is not applicable.  Required Exclusion:  • Members in hospice are excluded from the eligible population.	
an unplanned acute readmission for any diagnosis within 30 days and the predicted probability of an acute readmission.  Note: For Commercial and Medicaid, report only members 18 – 64	Note: Per General Guideline Members With Dual Enrollment, members with dual commercial and Medicaid enrollment may only be reported in the commercial product line. Members with dual Medicaid / Medicare	<ul> <li>Members in nospice are excluded from the eligible population.</li> <li>Exclude acute hospitalizations for the following reasons:</li> <li>Pregnancy</li> <li>A principle diagnosis for a perinatal condition originating in the perinatal period, on the discharge claim.</li> <li>Planned admission; chemotherapy, rehabilitation, an organ transplant or a potentially planned procedure without an acute diagnosis.</li> <li>Outliers:         <ul> <li>MA member with 4 or more IHS.</li> <li>Commercial member with 3 or more IHS.</li> <li>The member died during stay.</li> </ul> </li> </ul>	
years of age.  Event / diagnosis date:	enrollment"dual eligible" and with Medicare- Medicaid (MMP)	IHS (Index Hospital Stay): An acute inpatient or observation stay with a discharge on or between January 1 and December 1 of the measurement year, as identified in the denominator.	
1/1 to 12/1 of the measurement year.	enrollment may only be reported in the Medicare product line.	Tips for success:  Ensure proper HCC coding on all members. Having members coded to the highest specificity will help capture the complexity of a member.	

## POD



HEDIS® QUALITY MEASURE	CLINICAL GOAL	CRITERIA TO MEET THE GOAL
POD – Pharmacotherapy for Opioid Use Disorder  Administrative Measure  The percentage of Opioid Use Disorder (OUD) pharmacotherapy events that lasted at least 180 days among members 16 years of age and older with a diagnosis of OUD and a new	Members will have OUD pharmacotherapy for 180 or more days without a gap in treatment of 8 or more consecutive days.	Numerator compliance for this measure is based on pharmacy claims.      Medical record documentation is not applicable.  Required Exclusion:      Members in hospice or using hospice services any time during the measurement year.
OUD pharmacotherapy event.  The Intake Period: 7/1 of the prior year to 6/30 of the measurement year.	PRODUCT LINE  Commercial, Medicaid and Medicare	

## PPC

HEDIS® QUALITY MEASURE	CLINICAL GOAL	CRITERIA TO MEET THE GOAL
PPC – Prenatal and Postpartum Care  Hybrid Measure  Timeliness of Prenatal Care The percentage of deliveries that received a prenatal visit in the first trimester.  Postpartum Care The percentage of deliveries that had a postpartum visit on or between 7 and 84 days after delivery.	Members will receive a prenatal visit in the first trimester of pregnancy (or within 42 days of enrollment in Blue Cross NC plan).  Members will receive a postpartum visit with their provider between day 7 and day 84 postpartum.	Claims:  Submit claim for prenatal and postpartum visit with appropriate coding.  For Global Billing – You must submit an additional claim with the dates of the prenatal and postpartum visits. Refer to Corporate Reimbursement Policy.  Medical record documentation:  Date service rendered  Service rendered  EDC or LMP  Date of Delivery  Required Exclusion:
The Intake Period: 10/8 of the prior year to 10/7 of the measurement year.	PRODUCT LINE  Commercial, Marketplace and Medicaid	Members in hospice or using hospice services anytime during the measurement year.  HEDIS Value Set codes can be found in Appendix 7.



HEDIS® QUALITY MEASURE	CLINICAL GOAL	CRITEF	RIA TO	MEET THE GOAL	
PRS-E – Prenatal Immunization Status  ECDS Measure  The percentage of deliveries in the measurement period in which members had received influenza and tetanus, diphtheria toxoids and acellular pertussis (Tdap) vaccinations.  The denominator for this measure is based on deliveries, not on members.  Intake Period: 28 days prior to the delivery date through the delivery date.	Members who have deliveries will receive influenza, tetanus, diphtheria toxoids and acellular vaccinations.  PRODUCT LINE  Commercial and Medicaid	Influenza:  • Deliveries where members received an adult influenza vaccine on or between July 1 of the year prior to the measurement period and the delivery date, or  • Deliveries where members had anaphylaxis due to the influenza vaccine on or before the delivery date.  Tdap:  • Deliveries where members received at least one Tdap vaccine during the pregnancy, or  • Deliveries where members had any of the following:  - Anaphylaxis due to the diphtheria, tetanus or pertussis vaccine on or before the delivery date.  - Encephalitis due to the diphtheria, tetanus, or pertussis vaccine on or before the delivery date.  - Required Exclusions:  • Deliveries that occurred at less than 37 weeks gestation.  • Deliveries in which members were in hospice or using hospice services any time during the measurement year.			
		HEDIS Value Set Name	Code Set	Code	
		Adult Influenza Immunization	CVX	88, 135, 140, 141, 144, 150, 153, 155, 158, 166, 168, 171, 185, 186, 197, 205	
		Adult Influenza Vaccine Procedure	CPT	90630, 90653, 90654, 90656, 90658, 90661, 90662, 90673, 90674, 90682, 90686, 90688, 90689, 90694, 90756	
		Direct Reference Code (Formally Tdap Immunization)	CVX	115	
		Tdap Vaccine Procedure	CPT	90715	





	CRITERIA TO MEET THE GOAL
Antipsychotic Medications for Individuals with Schizophrenia  Administrative Measure  The percentage of members 18 years of age and older during the measurement year with schizophrenia or schizoaffective disorder who were dispensed and remained on an antipsychotic medication for at least 80% of their treatment period.  Adherence defined as:  A proportion of days covered (PDC) at 80% or over for antipsychotic medication(s) during the measurement period.  • Matherence defined as:  A proportion of days covered (PDC) at 80% or over for antipsychotic medication(s) during the measurement period.  • Matherence defined as:  A proportion of days covered (PDC) at 80% or over for antipsychotic medication(s) during the measurement period.	Claims:  Numerator compliance for this measure is based on pharmacy claims.  Medical record documentation is not applicable.  Required Exclusions:  Exclude members who met any of the following during the measurement year:  A diagnosis of dementia  Members in hospice or using hospice services any time during the measurement year.  Medicare members 66 years of age and older as of 12/31 of the measurement year who are in an I-SNP or living longterm in an institution any time during the measurement year.  Members 66 years of age and older as of 12/31 of the measurement year with frailty and advanced illness.  Members 81 years of age and older as of December 31 of the measurement year with at least two indications of frailty.  Did not have at least two antipsychotic medication dispensing events.



HEDIS® QUALITY MEASURE	CLINICAL GOAL	CRITERIA TO MEET THE GOAL			GOAL	
SNS-E – Social Need Screening and Intervention  ECDS Measure  The percentage of members who were screened, using prespecified instruments, at least once during the	Complete food, housing, and transportation screenings on all members; provide an appropriate intervention which includes assistance, assessment, counseling, coordination, education, evaluation of eligibility, provision or referral for a positive screening in any	An intervention corresponding to the type of need identified on or up to 30 days after the date of the first positive screening during the measurement period.  Interventions may include any of the following categories:				
measurement period for unmet food, housing and transportation needs, and received a corresponding intervention if they screened positive.  • Food Screening. The percentage of members who were screened for food	category.	screening numer specific screening Required Exclusion Members who benefit any time Members receithe measurements	erators. ng codi sions: o use ho me duri eiving p nent yea	Please ask your One of the measurement alliative care or ho	MC for a list of elect to use a hospice nt period. spice services during	
insecurity.  • Food Intervention. The	PRODUCT LINE Commercial, Medicaid	of the measu	rement	year who are in an	I-SNP or living long- e measurement year.	
percentage of members who received a corresponding intervention within 30 days	and Medicare	HEDIS Value Set Name	Code Set	Code	Comments	
(1 month) of screening positive for food insecurity.	nonth) of screening positive food insecurity.  using Screening. The centage of members who re screened for housing ability, homelessness or using inadequacy.  using Intervention. The centage of members who revention within 30 days (1 mith) of screening positive housing instability,	Food Insecurity Procedures	CPT	96156, 96160, 96161, 97802, 97803, 97804	Numerator compliance for Food Intervention	
Housing Screening. The percentage of members who		Food Insecurity Procedures	HCPCS	S5170, S9470	Numerator compliance for Food Intervention	
<ul> <li>instability, homelessness or housing inadequacy.</li> <li>Housing Intervention. The percentage of members who received a corresponding intervention within 30 days (1)</li> </ul>		Homelessness / Housing Instability / Inadequate Housing Procedures	СРТ	96156, 96160, 96161	Numerator compliance for Housing Intervention	
month) of screening positive for housing instability, homelessness or housing		Transportation Insecurity Procedures	CPT	96156, 96160, 96161	Numerator compliance for Transportation Intervention	
inadequacy.  • Transportation Screening. The percentage of members who were screened for transportation insecurity.  • Transportation Intervention. The percentage of members who received a corresponding intervention within 30 days (1 month) of screening positive for transportation insecurity.						





HEDIS® QUALITY MEASURE	CLINICAL GOAL	CRITERIA TO MEET THE GOAL
SPC – Statin Therapy for Patients With Cardiovascular Disease  Administrative Measure  The percentage of males 21 – 75 years of age and females 40 – 75 years of age during the measurement year, who were identified as having clinical atherosclerotic cardiovascular disease (ASCVD) and met the following criteria:  Two rates are reported:  Received Statin Therapy:  Members who were dispensed at least one high-intensity or moderate-intensity statin medication during the measurement year.  Statin Adherence 80%:  Members who remained on a high-intensity or moderate-intensity statin medication for at least 80% of the treatment period.	Members identified as having clinical atherosclerotic cardiovascular disease (ASCVD) will be prescribed and then maintain 80% adherence on a statin medication.  PRODUCT LINE  Commercial, Medicaid and Medicare	Claims:  Numerator compliance for this measure is based on pharmacy claims.  Medical record documentation not applicable.  Required Exclusions:  Pregnancy diagnosis in the measurement year or the year prior.  IVF – in the measurement year or the year prior.  ESRD in the measurement year or the year prior to the measurement year.  Dialysis during the measurement year or year prior to the measurement year.  Dispensed at least one prescription for clomiphene in the measurement year or the year prior.  Cirrhosis in the measurement year or the year prior.  Myalgia, myositis, myopathy or rhabdomyolysis during the measurement year.  Myalgia or rhabdomyolysis caused by a statin.  Members receiving palliative care or hospice services during the measurement year.  Medicare members 66 years of age and older as of 12/31 of the measurement year who are in an I-SNP or living longterm in an institution any time during the measurement year.  Members 66 years of age and older as of 12/31 of the measurement year who are in an I-SNP or living longterm in an institution any time during the measurement year.  Members 66 years of age and older as of 12/31 of the measurement year with frailty and advanced illness.





SPD – Statin Therapy for Patients With Diabetes  Administrative Measure  The percentage of members 40 – 75 years of age during the measurement year with diabetes but do not have clinical atherosclerotic cardiovascular disease (ASCVD) who met the following criteria:  Two rates are reported: Received Statin Therapy:  Members who were dispensed at least one statin medication of any intensity during the measurement year.  PRODUCT LINE  Claims:  Numerator compliance for this measure is to Medical record documentation not applicable.  Members whodo not have atherosclerotic cardiovascular disease (ASCVD) will be prescribed and then maintain 80% adherence on a statin medication.  Hedical record documentation not applicable.  Members with at least one of the following measurement year: MI, CABG, PCI, or other of Members with diagnosis of pregnancy during year prior.  Dispensed at least one prescription for clored year or the year prior.  ESRD diagnosis or dialysis during the measurement year or to Myalgia, myositis, myopathy or rhabdomyod year.  Members in palliative care or using hospice measurement year.  Medical record documentation not applicable.  Members with at least one of the following measurement year: MI, CABG, PCI, or other of Members with diagnosis of pregnancy during year prior.  ESRD diagnosis or dialysis during the measurement year or to Myalgia or rhabdomyolysis caused by a state of Members in palliative care or using hospice measurement year.  Medical record documentation not applicable.	
Administrative Measure  The percentage of members 40 – 75 years of age during the measurement year with diabetes but do not have clinical atherosclerotic cardiovascular disease (ASCVD) who met the following criteria:  Two rates are reported: Received Statin Therapy:  Members who were dispensed at least one statin medication of any intensity during the measurement year intensity during the measurement year.  who do not have atherosclerotic cardiovascular disease (ASCVD) will be prescribed and then maintain 80% adherence on a statin medication.  Medical record documentation not applicable Required Exclusions:  • Members with at least one of the following measurement year: MI, CABG, PCI, or other with the least one prescription for clomy year or the year prior.  • Dispensed at least one prescription for clomy year or the year prior.  • ESRD diagnosis or dialysis during the measurement year or the year prior.  • Myalgia or rhabdomyolysis caused by a statin medication of any intensity during the measurement year.  • Myalgia or rhabdomyolysis caused by a statin medication of any intensity during the measurement year.  • Myalgia or rhabdomyolysis caused by a statin medication of any intensity during the measurement year.  • Medicare members 66 years of age and old	is based on pharmacy claims.
The percentage of members 40 – 75 years of age during the measurement year with diabetes but do not have clinical atherosclerotic cardiovascular disease (ASCVD) who met the following criteria:  Two rates are reported:  Received Statin Therapy:  Members who were dispensed at least one statin medication of any intensity during the measurement year.  Teach of age during the measurement year with diagnosis of pregnancy during the measurement year prior.  Members with diagnosis of pregnancy during year prior.  Dispensed at least one prescription for clorn year or the year prior.  ESRD diagnosis or dialysis during the measurement year or the year prior.  Myalgia, myositis, myopathy or rhabdomyof year.  Myalgia or rhabdomyolysis caused by a statin medication of any intensity during the measurement year.  Medicare members 66 years of age and old.	
Statin Adherence 80%:  Members who remained on a statin medication of any intensity for at least 80% of the treatment period.  Members 66 years of age and older as of 12 frailty and advanced illness.  IVF during the measurement year or year properties.  Members 66 years of age and older as of 12 frailty and advanced illness.  IVF during the measurement year or year properties who had at least one encounter with measurement year or the year prior.	ing during the year prior to ther revascularization. during the measurement year or the clomiphene during the measurement neasurement year or the year prior. or the year prior. myolysis during the measurement a statin. pice anytime during the older as of 12/31 of the measurement term in an institution any time during of 12/31 of the measurement year with the prior to the measurement year.





#### TRC - Transitions of Care

Hybrid Measure – Members 18 years and older as of December 31 of the measurement year who had each of the following:

Medication Reconciliation Post-Discharge, Notification of Inpatient Admission, Receipt of Discharge Information, and Patient Engagement After Inpatient Admission.

Note: The denominator for this measure is based on discharges, not on members.



HEDIS® QUALITY MEASURE	CLINICAL GOAL	CRITERIA TO MEET THE GOAL
TRC – Notification of Inpatient Admission  Hybrid Measure  Documentation of receipt of notification of inpatient admission on the day of admission or through 2 days after the admission (3 total days).  Eligible Population Event – An acute or non-acute inpatient discharge on or between 1/1 and 12/1 of the measurement year.	Documentation in the outpatient medical record must include evidence of receipt of notification of inpatient admission on the day of admission through 2 days after admission. The documentation must include evidence of the date when the documentation was received.  PRODUCT LINE  DSNP and Medicare	<ul> <li>Medical record documentation, any of the following meet criteria:</li> <li>Communication between inpatient providers or staff and the member's PCP or ongoing care provider.</li> <li>Communication about admission between emergency department and the member's PCP or ongoing care provider.</li> <li>Communication about admission to the member's PCP or ongoing care provider: <ul> <li>Through a health information exchange; an automated admission, or discharge and transfer (ADT) alert system.</li> <li>Through a shared electronic medical record (EMR) system.</li> <li>From the member's health plan.</li> </ul> </li> <li>Indication: <ul> <li>That the member's PCP or ongoing care provider admitted the member to the hospital.</li> <li>That a specialist admitted the member to the hospital and notified the member's PCP or ongoing care provider.</li> <li>That the PCP or ongoing care provider placed orders for tests and treatments any time during the member's inpatient stay.</li> </ul> </li> <li>Documentation that the PCP or ongoing care provider performed a preadmission exam or received communication about a planned inpatient admission.</li> <li>Required Exclusion:</li> <li>Members in hospice or hospice services anytime during the measurement year.</li> </ul>

## TRC

<b>HEDIS® QUALITY MEASURE</b>	CLINICAL GOAL	CRITERIA TO MEET THE GOAL
TRC – Receipt of Discharge Information  Hybrid Measure  Documentation of receipt of discharge information on the day of discharge through 2 days after the discharge (3 total days).  Eligible Population Event – An acute or non-acute inpatient discharge on or between 1/1 and 12/1 of the measurement year.	Documentation in the outpatient medical record must include evidence of receipt of discharge information on the day of discharge through 2 days after the discharge with evidence of the date when the documentation was received.  PRODUCT LINE  DSNP and Medicare	<ul> <li>Medical record documentation: The discharge information must include all of the following: <ul> <li>The practitioner responsible for the member's care during the inpatient stay.</li> <li>Procedures or treatment provided.</li> <li>Diagnoses at discharge.</li> <li>Current medication list.</li> <li>Testing results, or documentation of pending tests or no tests pending.</li> <li>Instructions for patient care post-discharge.</li> </ul> </li> <li>Required Exclusion: <ul> <li>Members in hospice or hospice services anytime during the measurement year.</li> </ul> </li> </ul>

#### TRC - TRC



HEDIS® QUALITY MEASURE	CLINICAL GOAL	CRITERIA TO MEET THE GOAL			
TRC – Patient Engagement After Inpatient Discharge	Patient engagement occurs within 30 days after discharge.	Claims: • Submit a claim for patient engagement after discharge.			
Hybrid Measure		Medical record documentation*, any of the following meet criteria:			
Documentation of patient engagement within 30 days after discharge. Do not include		<ul><li>An outpatient v</li><li>A telephone vis</li></ul>		uding office visits and home visits.	
patient engagement that occurred on the date of discharge.		<ul> <li>A synchronous telehealth visit where real-tin occurred between the member and provider video communication.</li> </ul>			
Eligible Population Event – An acute or non-acute inpatient			ction, w	ck-in (asynchronous telehealth where hich was not real-time, occurred and provider).	
discharge on or between 1/1	PRODUCT LINE	RODUCT LINE			
and 12/1 of the measurement year.	DSNP and Medicare	Required Exclusi     Members in homeasurements	spice or	hospice services anytime during the	
		HEDIS Value Set Name	Code Set	Code	
		Outpatient and Telehealth	CPT	98966, 98967, 98968, 99441, 99442, 99443, 98970, 98971, 98972, 99421, 99422, 99423, 99457, 99458, 98980, 98981, 99202, 99203, 99204, 99205, 99211, 99212, 99213, 99214, 99215, 99242, 99243, 99244, 99245, 99341, 99342, 99344, 99345, 99347, 99348, 99389, 99381, 99382, 99383, 99384, 99385, 99386, 99387, 99391, 99392, 99393, 99394, 99395, 99396, 99397, 99401, 99402, 99403, 99404, 99411, 99412, 99429, 99455, 99456, 99483	
		Outpatient and Telehealth	HCPCS	G0402, G0438, G0439, G0463, G0071, G2010, G2012, G2250, G2251, G2252, T1015	
		Transitional Care Management Services	CPT	99495, 99496	
				entation is <b>only</b> accepted for gap al HEDIS Medical Record Review.	



## URI

This measure is reported as an inverted rate. A higher score indicates appropriate URI treatment.  Administrative Measure  The percentage of episodes for members 3 months of age and older with a diagnosis of upper respiratory infection (URI) that did NOT result in an antibiotic dispensing event.  The Intake Period: 7/1 of the prior year to 6/30 of the measurement year.  The Intake Period: 7/1 of the prior year to 6/30 of the measurement year.  The Intake Period: 7/1 of the prior year to 6/30 of the measurement year.  The Intake Period: 7/1 of the prior year to 6/30 of the measurement year.  The Intake Period: 7/1 of the prior year to 6/30 of the measurement year.  The Intake Period: 7/1 of the prior year to 6/30 of the measurement year.	HEDIS® QUALITY MEASURE	CLINICAL GOAL	CRITERIA TO MEET THE GOAL
Commercial, Marketplace, Medicare and Medicaid	URI – Appropriate Treatment for Upper Respiratory Infection This measure is reported as an inverted rate. A higher score indicates appropriate URI treatment.  Administrative Measure The percentage of episodes for members 3 months of age and older with a diagnosis of upper respiratory infection (URI) that did NOT result in an antibiotic dispensing event.  The Intake Period: 7/1 of the prior year to 6/30 of the	Antibiotics will NOT be prescribed to members who are diagnosed with URI only.  If there is another diagnosis which requires antibiotic treatment, include the coding information on the claim.  PRODUCT LINE  Commercial, Marketplace,	Claims:  Submit a claim for all additional competing diagnoses requiring antibiotic therapy on or within 3 days after the date of claim for URI.  Medical record documentation not applicable.  Required Exclusion:  Members in hospice or using



HEDIS® QUALITY MEASURE	CLINICAL GOAL	CRITERIA TO MEET THE GOAL					
W30 – Well-Child Visits in the First 30 Months of Life (Revised Measure Name from W15)	Children should have 6 or more well-child visits	Claims:  • Submit a claim for all member visits with proper coding for the visit servi  Medical record documentation not applicable.					
Administrative Measure  The percentage of members who had the following number of well-	during the first 15 months of life.  Required Exclusion:  • Members in hospice or using hospice services anytime during measurement year.  Children who  Note: Telehealth well visits no longer count for numerator complia						
child visits with a PCP during the last 15 months.	turn 30 months old during the measurement	HEDIS Value Set Name	Code Set	Code	Comments		
Two rates are reported: Well-Child Visits in the First 15 Months:	year should have 2 or more well- child visits	Well Care	СРТ	99381, 99382, 99383, 99384, 99385, 99391, 99392, 99393, 99394, 99395, 99461			
Children who turned 15 months old during the measurement year:	between 15 and 30 months.	Well Care	HCPCS	G0438, G0439, S0302, S0610, S0612, S0613			
Six or more well-child visits.      Well-Child Visits for Age 15	PRODUCT LINE	Encounter for Well Care	ICD10CM	Z00.00, Z00.01, Z00.110, Z00.111, Z00.121, Z00.129, Z00.2, Z00.3, Z01.411, Z01.419, Z02.5, Z02.84,	Do not include laboratory claims (claims with POS code 81)		
Months – 30 Months: Children who turned 30 months old during the measurement year:  Two or more well-child visits.	Commercial, Marketplace and Medicaid	Z76.1, Z76.2					

## WCC

## WCC – Weight Assessment and Counseling for Nutrition and Physical Activity for Children and Adolescents Hybrid Measure The percentage of members 3 – 17 years of age who had an outpatient visit with a PCP or OB/GYN and had the following during the measurement year (1/1 to 12/31):

HEDIS® QUALITY MEASURE	CLINICAL GOAL	CRITERIA TO MEET THE GOAL					
WCC – BMI Percentile	Members ages 3 – 17 will be	Claims:  • Submit a claim including the appropriate code					
BMI percentile during the measurement year.	assessed for height, weight, and BMI percentile during the measurement year.	sed for t, weight, MI percentile g the  Medical record documentation of:  Height, Weight and BMI percentile or BMI percentile plotted on age-growth chart during the measurement year. The height, weight and BMI percentile must be from the same data source.					
	PRODUCT LINE  Commercial, Marketplace and Medicaid	Required Exclusions:  • Members in hospice or using hospice services anytime during the measurement year.  • Members who have a diagnosis of pregnancy.  Note:  Member-collected biometric values (height, weight, BMI percentile) are eligible for use in reporting. They must be collected by a PCP or specialist, the information must be recorded, dated and maintained in the member's legal health record.					
		HEDIS® Value Set Name	Code Set	Code	Comment		
		BMI Percentile	ICD10M	Z68.51, Z68.52, Z68.53, Z68.54	Do not include laboratory claims (claims with POS code 81)		
		BMI Percentile	LOINC	59574-4, 59575-1, 59576-9	Do not include laboratory claims (claims with POS code 81)		





HEDIS® QUALITY MEASURE	<b>CLINICAL GOAL</b>	C	RITERI	A TO MEET THE (	GOAL	
WCC – Counseling or Nutrition  Documentation of counseling or nutrition or referral for nutrition education during he measurement year.	Members ages 3 – 17 will be counseled on nutrition during the measurement year.	Claims:  Submit a claim with the appropriate code.  Medical record documentation of:  A note indicating the date of service and at least one of the follow  Discussion of current nutrition behaviors.  Checklist indicating nutrition was addressed.  Educational materials on nutrition given to the member during to face visits.  Anticipatory guidance for nutrition.  Counseling or referral for nutrition education.  Weight or obesity counseling.  Referral to WIC.				
	PRODUCT LINE  Commercial, Marketplace and Medicaid					
		HEDIS Value Set Name	Code Set	Code	Comment	
		Nutrition Counseling	CPT	97802, 97803, 97804		
		Nutrition Counseling	HCPCS	G0270, G0271, G0447, S9449, S9452, S9470		
		Direct Reference Code (Formally BMI Percentile)	ICD10M	Z71.3	Do not include laboratory claims (claims with POS code 81)	

WCC





HEDIS® QUALITY MEASURE	CLINICAL GOAL	CRITERIA TO MEET THE GOAL				
WCC – Counseling for Physical Activity  Documentation of counseling for physical activity or referral for physical activity during the measurement year.	Members ages 3 – 17 will be counseled on physical activity during the measurement year.	I Medical record documentation of.				
	PRODUCT LINE  Commercial, Marketplace and Medicaid					
		HEDIS Value Set Name	Code Set	Code	Comment	
		Physical Activity Counseling	HCPCS	G0447, S9451		
		Encounter for Physical Activity Counseling	ICD10CM	Z02.5, Z71.82	Do not include laboratory claims (claims with POS code 81).	

## WCV

HEDIS® QUALITY MEASURE	CLINICAL GOAL		CRITE	RIA TO MEET THE G	iOAL .
WCV – Child and Adolescent Well-Care Visits  Administrative Measure  The percentage of members 3 – 21 years of age who had at least one comprehensive well-care visit with a PCP or an OB/GYN practitioner during the	Members 3 – 21 years of age will have at least one well-care visit during the measurement year. The visit must occur with a PCP or an OB/GYN practitioner but the practitioner does not have to be the practitioner assigned to the member.	Claims: Submit a claim for all member visits with proper codi the visit service.  Medical record documentation not applicable.  Required Exclusion: Members in hospice or using hospice services anytim during the measurement year.  Note: Telehealth well visits no longer count for numera compliance.			<b>ble</b> . vices anytime
measurement year.  Measure includes 3 age		HEDIS Value	e Code Set	Code	Comments
stratifications and total rate: • 3 – 11 years • 12 – 17 years		Well Care	CPT	99381, 99382, 99383, 99384, 99385, 99391, 99392, 99393, 99394, 99395, 99461	
• 18 – 21 years	PRODUCT LINE  Commercial, Marketplace and Medicaid	Well Care	HCPCS	G0438, G0439, S0302, S0610, S0612, S0613	
	and Medicald	Encounter for Well Care	ICD10CM	Z00.00, Z00.01, Z00.110, Z00.111, Z00.121, Z00.129, Z00.2, Z00.3, Z01.411, Z01.419, Z02.5, Z02.84, Z76.1, Z76.2	Do not include laboratory claims (claims with POS code 81)
		Refer to the AAP's Bright Futures website for more information about well-child visits (https://brightfutures.aap.org/materials-and-tools/guidelines-and-pocket-guide/).			

## Medicare Part D Pharmacy



MEDICARE PART D PHARMACY MEASURES	CLINICAL GOAL	CRITERIA TO MEET THE GOAL
Medication Adherence for Cholesterol (Statins)  Administrative Measure  The percentage of Medicare Part D beneficiaries 18 years or older who adhere to their prescribed drug therapy for statin cholesterol medications during the measurement period.	CONSIDER 90-day supply of medication. Medication synchronization. Cost savings by using preferred retail or mail order pharmacy.  EDUCATE member regarding medication compliance and risk factors.  ASSESS compliance and remove barriers to compliance.	<ul> <li>Claims:</li> <li>Information for this measure is based on claims data for prescription drugs filled by the members through December 31 of the measurement year.</li> <li>There is no reporting required from the provider.</li> <li>Medical record documentation not applicable.</li> <li>Adherence defined as: A proportion of days covered (PDC) at 80% or over for statin cholesterol medication(s) during the measurement period.</li> <li>Required Exclusions:</li> <li>Hospice enrollment.</li> <li>ESRD diagnosis.</li> <li>Dialysis coverage dates.</li> </ul>

MEDICARE PART D PHARMACY MEASURES	CLINICAL GOAL	CRITERIA TO MEET THE GOAL
Medication Adherence for Hypertension (RAS antagonists)  Administrative Measure  The percentage of Medicare Part D beneficiaries 18 years or older who adhere to their prescribed drug therapy for Renin Angiotensin System (RAS) antagonists: Angiotensin Converting Enzyme Inhibitor (ACEI), Angiotensin Receptor Blocker (ARB), or direct renin inhibitor medications.	CONSIDER 90-day supply of medication. Medication synchronization. Cost savings by using preferred retail or mail order pharmacy.  EDUCATE member regarding medication compliance and risk factors.  ASSESS compliance and remove barriers to compliance.	<ul> <li>Claims:</li> <li>Information for this measure is based on claims data for prescription drugs filled by the members through December 31 of the measurement year.</li> <li>There is no reporting required from the provider.</li> <li>Medical record documentation not applicable.</li> <li>Adherence defined as: A proportion of days covered (PDC) at 80% or higher for RAS antagonist medications during the measurement period.</li> <li>Required Exclusions:</li> <li>Hospice enrollment.</li> <li>ESRD diagnosis or dialysis coverage dates.</li> <li>One or more prescriptions for sacubitril / valsartan.</li> </ul>

#### **Medicare Part D Pharmacy**

## Medicare Part D Pharmacy



• ESRD diagnosis or dialysis coverage dates.

MEDICARE PART D PHARMACY MEASURES	CLINICAL GOAL	CRITERIA TO MEET THE GOAL
Medication Adherence for Diabetes Medications  Administrative Measure	CONSIDER 90-day supply of medication. Medication synchronization.	Claims:  • Information for this measure is based on claims data for prescription drugs filled by the members through December 31 of the
The percentage of Medicare Part D beneficiaries 18 years or older who adhere to their prescribed drug therapy across classes of diabetes	Cost savings by using preferred retail or mail order pharmacy.  An alternate generic regimen for therapeutic coverage of brand name medications for greater affordability, 1or in the even of a drug shortage.	measurement year.  There is no reporting required from the provider.  Medical record documentation not applicable.
medications: biguanides, sulfonylureas, thiazolidinediones, DiPeptidyl Peptidase (DDP)-4 inhibitors, GLP-1 receptor agonists, meglitinides, and sodium glucose cotransporter 2 (SGLT2) inhibitors.	EDUCATE member regarding medication compliance and risk factors.  ASSESS compliance and remove barriers to compliance.	Adherence defined as: A proportion of days covered (PDC) at 80% or higher across the classes of diabetes medications.  Required Exclusions:  One or more prescriptions for insulin.  Hospice enrollment.

Statin Use in Persons with Diabetes (SUPD)  Administrative Measure  The percentage of Medicare Part D beneficiaries between 40 and 75 years old who received at least two diabetes medication fills and also received a statin medication during the measurement period.  EDUCATE member regarding medication compliance and risk factors.  ASSESS compliance and remove barriers to compliance.  ASSESS compliance.  EVALUATE Measurement period.  ASSESS Compliance and remove barriers to compliance.  EVENT MEMBER SOLUCION (Code Set Submitted each measurement year)  EVENT MEMBER SOLUCION (Code Set Submitted each measurement year)  EVENT MEMBER SOLUCION (Code Set Submitted each measurement year)  EVENT MEMBER SOLUCION (Code Set Submitted each measurement year)  EVENT MEMBER SOLUCION (Code Set Submitted each measurement year)  EVENT MEMBER SOLUCION (Code Set Submitted each measurement year)  EVENT MEMBER SOLUCION (Code Set Submitted each measurement year)  EVENT MEMBER SOLUCION (Code Set Submitted each measurement year)  EVENT MEMBER SOLUCION (Code Set Submitted each measurement year)  EVENT MEMBER SOLUCION (Code Set Submitted each measurement year)  EVENT MEMBER SOLUCION (Code Set Submitted each measurement year)  EVENT MEMBER SOLUCION (Code Set Submitted each measurement year)  EVENT MEMBER SOLUCION (Code Set Submitted each measurement year)  EVENT MEMBER SOLUCION (Code Set Submitted each measurement year)  EVENT MEMBER SOLUCION (Code Set Submitted each measurement year)  EVENT MEMBER SOLUCION (Code Set Submitted each measurement year)  EVENT MEMBER SOLUCION (Code Set Submitted each measurement year)  EVENT MEMBER SOLUCION (Code Set Submitted each measurement year)  EVENT MEMBER SOLUCION (Code Set Submitted each measurement year)  EVENT MEMBER SOLUCION (Code Set Submitted each measurement year)  EVENT MEMBER SOLUCION (Code Set Submitted each measurement year)  EVENT MEMBER SOLUCION (Code Set Submitted each measurement year)  EVENT MEMBER SOLUCION (Code Set Submitted each measurement year)  EVENT MEMBER SOLUCION (Code Set Submit	MEDICARE PART D PHARMACY MEASURES	CLINICAL GOAL	CRITERIA TO MEET THE GOAL				
Medicare Part D beneficiaries between 40 and 75 years old who received at least two diabetes medication fills and also received a statin medication during the measurement period.  EDUCATE member regarding medication compliance and risk factors.  ASSESS compliance and remove barriers to compliance.  ASSESS compliance.  ASSESS compliance and remove barriers to compliance.  Medical record documentation not applicable.  Required Exclusions: Any of the following during the measurement year:  Hospice enrollment.  ESRD diagnosis or dialysis coverage dates  Rhabdomyolysis and myopathy  Pregnancy  Lactation and fertility  Cirrhosis  Pre-Diabetes  Polycystic Ovary Syndrome  Note: The ICD-10 diagnosis code must be submitted each measurement year for exclusion from the measure.  Exclusion*  Cost savings by using preferred retail or mail order pharmacy.  Hospice enrollment.  ESRD diagnosis or dialysis coverage dates  Rhabdomyolysis and myopathy  Pregnancy  Lactation and fertility  Cirrhosis  Pre-Diabetes  Polycystic Ovary Syndrome  Note: The ICD-10 diagnosis code must be submitted each measurement year for exclusion from the measure.  Exclusion*  Code Set  Code  Exclusion*  Code Set  Code  Code  Cirrhosis  Currently pregnant, breast- feeding, or of child-bearing age  Polycystic ovary syndrome (PCOS)  ICD10CM	with Diabetes (SUPD)	90-day supply of	<ul> <li>Information for this measure is based on claims data for prescription drugs</li> </ul>				
Medicare Part D beneficiaries between 40 and 75 years old who received at least two diabetes medication fills and also received a statin medication during the measurement period.  EDUCATE member regarding medication compliance and risk factors.  ASSESS compliance and remove barriers to compliance.  **Robinson**  Cost savings by using preferred retail or mail order pharmacy.  **EDUCATE** member regarding medication compliance and risk factors.  **ASSESS** compliance and remove barriers to compliance.  **Pre-Diabetes** **Polycystic Ovary Syndrome**  Note: The ICD-10 diagnosis code must be submitted each measurement year for exclusion from the measure.  **Exclusion***  Code Set**  Code**  Rhabdomyolysis or myopathy  Pre-diabetes*  Cirrhosis*	Administrative Measure	Medication	There is no reporting required from the provider.				
Medicare Part D beneficiaries between 40 and 75 years old who received at least two diabetes medication fills and also received a statin medication during the measurement period.  EDUCATE member regarding medication compliance and risk factors.  ASSESS compliance and remove barriers to compliance.  Exclusion*  Code Set Rhabdomyolysis or myopathy  Pre-diabetes  Code Rhabdomyolysis or myopathy  Diction Ry3.03, Ry3.09  Cirrhosis  Code Rhabdomyolysis or myopathy  Pre-diabetes  Cirrhosis  Code Rhabdomyolysis or myopathy  Diction Ry3.03, Ry3.09  Cirrhosis  Currently pregnant, breast- feeding, or of child-bearing age  Polycystic ovary syndrome (PCOS)  ESRD  ICD10CM   E28.2  ESRD  ICD10CM   I12.0, I13.11, I13.2, N18.5, N18.6, N19, Z91.15, Z99.2	The percentage of	synchronization.	Medical record documentar	tion not a	pplicable.		
Exclusion*   Code Set   Code	beneficiaries between 40 and 75 years old who received at least two diabetes medication fills and also received a statin medication during the	using preferred retail or mail order pharmacy.  EDUCATE member regarding medication compliance and risk factors.  ASSESS compliance and	Any of the following during the measurement year:  • Hospice enrollment.  • ESRD diagnosis or dialysis coverage dates  • Rhabdomyolysis and myopathy  • Pregnancy  • Lactation and fertility  • Cirrhosis  • Pre-Diabetes  • Polycystic Ovary Syndrome  Note: The ICD-10 diagnosis code must be submitted each measurement year				
Pre-diabetes         ICD10CM         R73.03, R73.09           Cirrhosis         ICD10CM         K70.30, K70.31, K71.7, K74.3, K74.4, K74.5, K74.60, K74.69           Currently pregnant, breast-feeding, or of child-bearing age         ICD10CM         091.03, 092.03, Z39.1, 000.111, etc.           Polycystic ovary syndrome (PCOS)         ICD10CM         E28.2           ESRD         ICD10CM         I12.0, I13.11, I13.2, N18.5, N18.6, N19, Z91.15, Z99.2			Exclusion*	Code Set	Code		
Cirrhosis         ICD10CM         K70.30, K70.31, K71.7, K74.3, K74.4, K74.5, K74.60, K74.69           Currently pregnant, breast-feeding, or of child-bearing age         ICD10CM         091.03, 092.03, Z39.1, 000.111, etc.           Polycystic ovary syndrome (PCOS)         ICD10CM         E28.2           ESRD         ICD10CM         I12.0, I13.11, I13.2, N18.5, N18.6, N19, Z91.15, Z99.2			Rhabdomyolysis or myopathy	ICD10CM	G72.0, G72.89, G72.9, M60.80, M60.9, M62.82, etc.		
Currently pregnant, breast-feeding, or of child-bearing age  Polycystic ovary syndrome (PCOS)   ICD10CM   E28.2  ESRD   ICD10CM   I12.0, I13.11, I13.2, N18.5, N18.6, N19, Z91.15, Z99.2			Pre-diabetes	ICD10CM	R73.03, R73.09		
feeding, or of child-bearing age			Cirrhosis	ICD10CM	K70.30, K70.31, K71.7, K74.3, K74.4, K74.5, K74.60, K74.69		
ESRD ICD10CM I12.0, I13.11, I13.2, N18.5, N18.6, N19, Z91.15, Z99.2				ICD10CM	091.03, 092.03, Z39.1, 000.111, etc.		
			Polycystic ovary syndrome (PCOS)	ICD10CM	E28.2		
*Please contact your QMC for a complete list of exclusion codes.			ESRD	ICD10CM	I12.0, I13.11, I13.2, N18.5, N18.6, N19, Z91.15, Z99.2		
			*Please contact your QMC for a	complete li	st of exclusion codes.		

#### **Medicare Part D Pharmacy**



## **APPENDIX 1**

#### **BMI CODE CHART**

# ICD-10-CM CODES FOR WCC AGES 3 – 17 PER HEDIS SPECIFICATIONS Z68.51 Body mass index (BMI) pediatric, less than 5th percentile for age Z68.52 Body mass index (BMI) pediatric, 5th percentile to less than 85th percentile for age Z68.53 Body mass index (BMI) pediatric, 85th percentile to less than 95th percentile for age

Z68.54 Body mass index (BMI) pediatric, greater than or equal to 95th percentile for age

### **APPENDIX 2**

#### MEASURES USED IN OVERALL MEDICARE STAR RATINGS

The Centers for Medicare and Medicaid Services (CMS) uses a five-star quality rating system to promote improvement in quality. The following weighted measures are utilized when calculating an overall Medicare Star Rating:

MEASURE	WEIGHT
Medication Adherence for Cholesterol (Statins)	3
Medication Adherence for Hypertension (RAS Antagonists)	3
Medication Adherence for Diabetes Medications	3
GSD – Glycemic Status Assessment for Patients with Diabetes > 9.0%	3
CBP – Controlling High Blood Pressure	3
PCR – Plan All-Cause Readmissions	3
OMW – Osteoporosis Management	1
COL-E – Colorectal Cancer Screening	1
FMC – Follow-Up After Emergency Department Visit for People with Multiple High-Risk Chronic Conditions	1
EED – Eye Exam for Patients with Diabetes	1
TRC – Transitions of Care	1
SPC – Statin Therapy for Patients with Cardiovascular Disease	1
SUPD – Statin Use in Persons with Diabetes	1
BCS-E -Breast Cancer Screening	1
KED - Kidney Health Evaluation for Patients with Diabetes	1



#### **APPENDIX 3**

#### CAHPS AND MEDICARE HOS INFORMATION

#### CAHPS (Consumer Assessment of Healthcare Providers and Systems) Survey: Evaluating the Patient's Experience

The CAHPS® survey evaluates the consumer's perspective of the quality of health services provided by Commercial, Medicare Advantage and Part D programs.

#### Below are some tips to addressing the CAHPS-specific patient experience factors:

- Be familiar with the questions patients are being asked on the member survey.
- Blue Cross and Blue Shield of North Carolina (Blue Cross NC) has changed the benefit for payment of the Medicare Advantage Annual Wellness Visit from once every 12 months to once per calendar year. This reconfiguration took effect January 1, 2021 and will remain in place until further notice.
- Encourage patients to get a flu shot each year.
- Understand patients' costs for prescribed drugs based on their plan formulary.
- Follow up with patients promptly regarding test results, regardless of whether the results require additional care.
- Prioritize timely appointment scheduling, for both routine and specialty care.

#### Medicare Health Outcomes Survey (HOS) 101: For Providers

The HOS member survey assesses the physical and mental health of patients over a two year period. Responses to this survey contribute to the plan's overall Star Rating. Many of the survey questions ask the patient if they have had conversations with their doctor or nurse about activity level, falls, and bladder control.

#### **Best Practices for Providers:**

- Be familiar with the questions patients are being asked on the member survey.
- Place reminders in your EMR to speak with patients at each visit regarding these topics:
  - Suggest that they use a cane or walker if needed.
  - Lying or standing blood pressure.
  - Exercise or physical therapy program recommendations.
  - Vision or hearing test (if necessary).
- Advise patients on ways to manage the leaking of urine, including bladder training exercises, medication and surgery.



#### **MEDICARE MEDICATIONS WITH \$0 COPAY\***

#### 2025 Formulary Tier 6 Select Care Drugs – GENERIC DRUGS ONLY

\*\$0 copay at Preferred Retail Pharmacies and the following Preferred Mail Order Pharmacies:
Walgreens Mail Service and Express Scripts Pharmacy
(List of Preferred Pharmacies available at BlueCrossNC.com/Find-a-Drug-or-Pharmacy.)

	DRUG CLASS						
Angiotensin – Converting Enzyme Inhibitors (ACE-I)	Angiotensin – Receptor Blockers (ARB)	Renin Inhibitors	Statins				
Benazepril Benazepril / Amlodipine Benazepril / HCTZ Captopril Enalapril tablet Enalapril / HCTZ Fosinopril Fosinopril / HCTZ Lisinopril Lisinopril / HCTZ Moexipril Perindopril Quinapril Quinapril Trandolapril	Candesartan Candesartan / HCTZ Irbesartan Irbesartan / HCTZ Losartan Losartan / HCTZ Olmesartan Olmesartan / Amlodipine Olmesartan / Amlodipine / HCTZ Olmesartan / HCTZ Telmisartan Telmisartan / Amlodipine Telmisartan / HCTZ Valsartan Valsartan / Amlodipine Valsartan / Amlodipine Valsartan / Amlodipine Valsartan / Amlodipine	Aliskiren	Atorvastatin Atorvastatin / Amlodipine Fluvastatin (20 mg, 40 mg) Lovastatin Pravastatin Rosuvastatin Simvastatin Simvastatin				
	COMMON USE						
BLOOD PRESSURE	BLOOD PRESSURE	BLOOD PRESSURE	CHOLESTEROL				

DRUG CLASS				
Sulfonylureas	Thiazolidinediones (TZDs)	Meglitinides	Biguanides	
Glimepiride Glimepiride / Pioglitazone Glipizide Glipizide ER Glipizide XL Glipizide / Metformin Glyburide Glyburide Micronized Glyburide / Metformin	Pioglitazone Pioglitazone / Glimepiride Pioglitazone / Metformin	Nateglinide Repaglinide	Metformin tablet Metformin ER tablet** Metformin / Glipizide Metformin / Glyburide Metformin / Pioglitazone  ** Metformin ER does not include generic Fortamet (Metformin ER osmotic release tablet) or Glumetza (Metformin ER modified release tablet).	
	COMMON USE			
DIABETES	DIABETES	DIABETES	DIABETES	

### **APPENDIX 5**



#### **ELECTRONIC CLINICAL DATA SYSTEMS (ECDS) MEASURES**

#### **Electronic Clinical Data Systems (ECDS) Measures**

- A structured method to collect and report electronic clinical data for HEDIS quality measurement and for quality improvement.
- HEDIS reporting standard for health plans collecting and submitting measures to NCQA. This
  reporting standard defines the data sources and types of structured data acceptable for use for
  a measure.
  - **‡** Data collection:
    - Personal Health Record (PHR) / Electronic Health Record (EHR)
    - Clinical Registry / Health Information Exchange (HIE)
    - Case management system
    - Admin / enrollment
  - ‡ Data must:
    - Use standard layouts
    - Meet the technical specifications
    - Must be accessible by the care team upon request
    - Elements reported according to data source

NCQA developed ECDS to encourage health information exchange, the secure sharing of patient medical information electronically.

## Appendix 6 Measure Value Set Codes (Continued)





HEDIS VALUE SET NAME	CODE SET	CODE	COMMENTS
BH Outpatient	СРТ	98960, 98961, 98962, 99078, 99202, 99203, 99204, 99205, 99211, 99212, 99213, 99214, 99215, 99242, 99244, 99245, 99341, 99342, 99344, 99345, 99347, 99348, 99349, 99350, 99381, 99382, 99383, 99384, 99385, 99386, 99387, 99391, 99392, 99393, 99394, 99395, 99396, 99397, 99401, 99402, 99403, 99404, 99411, 99412, 99483, 99492, 99493, 99494, 99510	
BH Outpatient	HCPCS	G0155, G0176, G0177, G0409, G0463, G0512, H0002, H0004, H0031, H0034, H0036, H0037, H0039, H0040, H2000, H2010, H2011, H2013, H2014, H2015, H2016, H2017, H2018, H2019, H2020, T1015	
Health and Behavior Assessment or Intervention	СРТ	96156, 96158, 96159, 96164, 96165, 96167, 96168, 96170, 96171	
Partial Hospitalization or Intensive Outpatient	HCPCS	G0410, G0411, H0035, H2001, H2012, S0201, S9480, S9484, S9485	
Telephone Visits	CPT	98966, 98967, 98968, 99441, 99442, 99443	
Online Assessment	CPT	98970, 98971, 98972, 99421, 99422, 99423, 99457, 99458, 98980, 98981	Only Numerator Compliance for Rate 2 (C&M Phase), and only one of the two visits (during the 31 – 300 days after the IPSD) may be an e-visit or virtual check-in.
Online Assessment	HCPCS	G0071, G2010, G2012, G2250, G2251, G2252	Only Numerator Compliance for Rate 2 (C&M Phase), and only one of the two visits (during the 31 – 300 days after the IPSD) may be an e-visit or virtual check-in.
Direct Reference Code	POS	52, 53	With a Visit Setting Unspecified code
Telehealth POS	POS	2, 10	With a Visit Setting Unspecified code
Outpatient POS	POS	03, 05, 07, 09, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 22, 33, 49, 50, 71, 72	With a Visit Setting Unspecified code

## CIS (continued)



HEDIS VALUE SET NAME	CODE SET	CODE
Hepatitis B Immunization	CVX	8, 44, 45, 51, 110, 146, 198
Hepatitis B Vaccine Procedure	CPT	90697, 90723, 90740, 90744, 90747, 90748
Hepatitis B Vaccine Procedure	HCPCS	G0010
Direct Reference Code (Newborn Hepatitis B Vaccine)	1CD-10-PCS	3E0234Z
Haemophilus Influenzae Type B (HiB) Immunization	CVX	17, 46, 47, 48, 49, 50, 51, 120, 146, 148, 198
Haemophilus Influenzae Type B (HiB) Vaccine Procedure	CPT	90644, 90647, 90648, 90697, 90698, 90748
Inactivated Polio Vaccine (IPV) Immunization	CVX	10, 89, 110, 120, 146
Inactivated Polio Vaccine (IPV) Procedure	CPT	90697, 90698, 90713, 90723
Influenza Immunization	CVX	88, 140, 141, 150, 153, 155, 158, 161, 171, 186
Influenza Vaccine Procedure	CPT	90655, 90657, 90661, 90673, 90674, 90685, 90686, 90687, 90688, 90689, 90756
Influenza Virus LAIV Immunization	CVX	111, 149
Influenza Virus LAIV Vaccine Procedure	CPT	90660, 90672
Measles	ICD10CM	B05.0, B05.1, B05.2, B05.3, B05.4, B05.81, B05.89, B05.9
Measles, Mumps and Rubella (MMR) Immunization	CVX	3, 94
Measles, Mumps and Rubella (MMR) Vaccine Procedure	CPT	90707, 90710
Mumps	ICD10CM	B26.0, B26.1, B26.2, B26.3, B26.81, B26.82, B26.83, B26.84, B26.85, B26.89, B26.9
Rubella	ICD10CM	B06.00, B06.01, B06.02, B06.09, B06.81, B06.82, B06.89, B06.9
Pneumococcal Conjugate Immunization	CVX	109, 133, 152, 215
Pneumococcal Conjugate Vaccine Procedure	CPT	90670, 90671
Pneumococcal Conjugate Vaccine Procedure	HCPCS	G0009
Direct Reference Code (formally Rotavirus) (2 Dose Schedule) Immunization	CVX	119
Rotavirus (3 Dose Schedule) Immunization	CVX	116, 122
Rotavirus Vaccine (2 Dose Schedule) Procedure	CPT	90681
Rotavirus Vaccine (3 Dose Schedule) Procedure	CPT	90680
Varicella Zoster (VZV) Immunization	CVX	21, 94
Varicella Zoster (VZV) Vaccine Procedure	CPT	90710, 90716
Varicella Zoster	ICD10CM	B01.0, B01.11, B01.12, B01.2, B01.81, B01.89, B01.9, B02.0, B02.1, B02.21, B02.22, B02.23, B02.24, B02.29, B02.30, B02.31, B02.32, B02.33, B02.34, B02.39, B02.7, B02.8, B02.9





HEDIS VALUE SET NAME	CODE SET	CODE	COMMENTS
Visit Setting Unspecified	CPT	90791, 90792, 90832, 90833, 90834, 90836, 90837, 90838, 90839, 90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221, 99222, 99223, 99231, 99232, 99233, 99238, 99239, 99252, 99253, 99254, 99255	With one of the following:  Outpatient POS code  POS code 52  POS code 53  Telehealth POS code
BH Outpatient	CPT	98960, 98961, 98962, 99078, 99202, 99203, 99204, 99205, 99211, 99212, 99213, 99214, 99215, 99242, 99243, 99244, 99245, 99341, 99342, 99344, 99345, 99347, 99348, 99349, 99350, 99381, 99382, 99383, 99384, 99385, 99386, 99387, 99391, 99392, 99393, 99394, 99395, 99396, 99397, 99401, 99402, 99403, 99404, 99411, 99412, 99483, 99492, 99493, 99494, 99510	
BH Outpatient	HCPCS	G0155, G0176, G0177, G0409, G0463, G0512, H0002, H0004, H0031, H0034, H0036, H0037, H0039, H0040, H2000, H2010, H2011, H2013, H2014, H2015, H2016, H2017, H2018, H2019, H2020, T1015	
Partial Hospitalization or Intensive Outpatient	HCPCS	G0410, G0411, H0035, H2001, H2012, S0201, S9480, S9484, S9485	
Electroconvulsive Therapy	СРТ	90870	with Outpatient POS code OR POS code 24 OR POS code 52 OR POS code 53
Substance Use Disorder Services	CPT	99408, 99409	
Substance Use Disorder Services	HCPCS	G0396, G0397, G0443, H0001, H0005, H0007, H0015, H0016, H0022, H0047, H0050, H2035, H2036, T1006, T1012	
Outpatient POS	POS	03, 05, 07, 09, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 22, 33, 49, 50, 71, 72	With a Visit Setting Unspecified code <b>OR</b> Electroconvulsive Therapy
Direct Reference Code	POS	52, 53	With a Visit Setting Unspecified code <b>OR</b> Electroconvulsive Therapy
Direct Reference Code	POS	24	With Electroconvulsive Therapy
Telehealth POS	POS	2, 10	With a Visit Setting Unspecified code
Substance Abuse Counseling and Surveillance	ICD10CM	Z71.41, Z71.51	Do not include laboratory claims (claims with POS code 81)
Outpatient and Telehealth	СРТ	98966, 98967, 98968, 98970, 98971, 98972, 98980, 98981, 99202, 99203, 99204, 99205, 99211, 99212, 99213, 99214, 99215, 99241, 99242, 99243, 99244, 99245, 99341, 99342, 99343, 99344, 99345, 99347, 99348, 99349, 99350, 99381, 99382, 99383, 99384, 99385, 99386, 99387, 99391, 99392, 99393, 99394, 99395, 99396, 99397, 99401, 99402, 99403, 99404, 99411, 99412, 99421, 99422, 99423, 99429, 99441, 99442, 99443, 99455, 99456, 99457, 99458, 99483	
Outpatient and Telehealth	HCPCS	G0071, G0402, G0438, G0439, G0463, G2010, G2012, G2250, G2251, G2252, T1015	

## FUA (continued)



HEDIS VALUE SET NAME	CODE SET	CODE	COMMENTS
AOD Abuse and Dependence	ICD10CM	F10.10, F10.120, F10.121, F10.129, F10.130, F10.131, F10.132, F10.139, F10.14, F10.150, F10.151, F10.159, F10.180, F10.181, F10.182, F10.188, F10.19, F10.20, F10.220, F10.221, F10.229, F10.230, F10.231, F10.232, F10.239, F10.24, F10.250, F10.251, F10.259, F10.26, F10.27, F10.280, F10.281, F10.282, F10.288, F10.29, F11.10, F11.120, F11.121, F11.122, F11.129, F11.13, F11.14, F11.150, F11.151, F11.159, F11.181, F11.182, F11.188, F11.19, F11.20, F11.220, F11.221, F11.222, F11.229, F11.23, F11.24, F11.250, F11.251, F11.259, F11.281, F11.282, F11.288, F11.29, F12.10, F12.120, F12.121, F12.122, F12.129, F12.13, F12.150, F12.151, F12.159, F12.251, F12.259, F12.280, F12.281, F12.29, F13.130, F13.131, F13.131, F13.132, F13.331, F13.312, F13.331, F13.3131, F13.3132, F13.339, F13.344, F13.150, F13.151, F13.150, F13.151, F13.159, F13.180, F13.181, F13.182, F13.188, F13.19, F13.20, F13.220, F13.221, F13.229, F13.230, F13.231, F13.232, F13.239, F13.24, F13.250, F13.251, F13.259, F13.26, F13.27, F13.280, F13.281, F13.282, F13.288, F13.29, F14.10, F14.120, F14.121, F14.122, F14.129, F14.13, F14.14, F14.150, F14.151, F14.159, F14.180, F14.181, F14.182, F14.188, F14.19, F14.20, F14.221, F14.222, F14.229, F14.23, F14.24, F14.250, F14.251, F14.259, F14.280, F14.281, F14.282, F14.288, F14.29, F15.10, F15.120, F15.121, F15.122, F15.129, F15.13, F15.14, F15.150, F15.151, F15.159, F15.180, F15.181, F15.182, F15.188, F15.19, F15.20, F15.221, F15.222, F15.229, F16.20, F16.221, F16.229, F16.24, F16.250, F16.251, F16.259, F16.284, F16.250, F16.251, F16.259, F16.284, F16.250, F16.251, F16.259, F16.280, F16.283, F16.29, F18.17, F18.180, F18.184, F18.19, F18.284, F18.29, F18.27, F18.280, F18.284, F18.296, F18.297, F18.299, F18.27, F18.280, F18.284, F19.286, F19.297, F19.280, F19.281, F19.282, F19.288, F19.299, F19.20, F19.250, F19.251,	To be used with appropriate visit codes for numerator compliance
AOD Medication Treatment	HCPCS	G2069, G2070, G2072, G2073, H0020, H0033, J0570, J0571, J0572, J0573, J0574, J0575, J0577, J0578, J2315, Q9991, Q9992, S0109	
Substance Induced Disorders	ICD10CM	F10.920, F10.921, F10.929, F10.930, F10.931, F10.932, F10.939, F10.94, F10.950, F10.951, F10.959, F10.96, F10.97, F10.980, F10.981, F10.982, F10.988, F10.99, F11.90, F11.920, F11.921, F11.922, F11.929, F11.93, F11.94, F11.950, F11.951, F11.959, F11.981, F11.982, F11.988, F11.99, F12.90, F12.920, F12.921, F12.922, F12.929, F12.93, F12.950, F12.950, F12.951, F12.959, F12.980, F12.988, F12.99, F13.90, F13.920, F13.921, F13.929, F13.930, F13.931, F13.932, F13.939, F13.94, F13.950, F13.951, F13.959, F13.96, F13.97, F13.980, F13.981, F13.982, F13.988, F13.99, F14.90, F14.920, F14.921, F14.922, F14.929, F14.93, F14.94, F14.950, F14.951, F14.959, F14.980, F14.981, F14.982, F14.988, F14.99, F15.90, F15.920, F15.921, F15.922, F15.929, F15.93, F15.94, F15.950, F15.951, F15.959, F15.980, F15.981, F15.982, F15.988, F15.99, F16.90, F16.920, F16.921, F16.929, F16.94, F16.950, F16.951, F16.959, F16.983, F16.988, F16.99, F18.90, F18.920, F18.921, F18.929, F18.94, F18.950, F18.951, F18.959, F18.97, F18.980, F18.988, F18.99, F19.90, F19.920, F19.921, F19.922, F19.929, F19.930, F19.931, F19.932, F19.938, F19.94, F19.950, F19.951, F19.959, F19.96, F19.97, F19.980, F19.981, F19.982, F19.988, F19.99, F10.90	To be used with appropriate visit codes for numerator compliance
Unintentional Drug Overdose	ICD10CM	T40.0X1A, T40.0X1D, T40.0X1S, T40.0X4A, T40.0X4D, T40.0X4S, T40.1X1A, T40.1X1D, T40.1X1S, T40.1X4A, T40.1X4D, T40.1X4S, T40.2X1A, T40.2X1D, T40.2X1S, T40.2X4A, T40.2X4D, T40.2X4S, T40.3X1A, T40.3X1D, T40.3X1S, T40.3X4A, T40.3X4D, T40.3X4S, T40.411A, T40.411D, T40.411S, T40.414A, T40.414D, T40.414S, T40.421A, T40.421D, T40.421S, T40.424A, T40.424D, T40.424S, T40.421A, T40.491D, T40.491D, T40.491S, T40.494A, T40.494D, T40.494S, T40.5X1A, T40.5X1D, T40.5X1S, T40.5X4A, T40.5X4D, T40.5X4S, T40.601A, T40.601D, T40.601S, T40.604A, T40.604D, T40.604S, T40.691A, T40.691D, T40.691S, T40.694A, T40.694D, T40.604S, T40.711D, T40.711S, T40.714A, T40.714D, T40.714S, T40.721A, T40.721D, T40.721S, T40.724A, T40.724D, T40.724S, T40.8X1A, T40.8X1D, T40.8X1S, T40.8X4A, T40.8X4D, T40.891D, T40.901A, T40.904D, T40.904S, T40.991A, T40.991D, T40.991S, T40.994A, T40.994D, T40.904S, T41.0X1A, T41.0X1D, T41.0X1S, T41.0X4A, T41.0X4D, T41.0X4S, T41.1X1A, T41.1X1D	To be used with appropriate visit codes for numerator compliance





HEDIS VALUE SET NAME	CODE SET	CODE	COMMENTS
Unintentional Drug Overdose	ICD10CM	T41.1X1S, T41.1X4A, T41.1X4D, T41.1X4S, T41.201A, T41.201D, T41.201S, T41.204A, T41.204D, T41.204S, T41.291A, T41.291D, T41.291S, T41.294A, T41.294D, T41.294S, T41.3X1A, T41.3X1D, T41.3X1S, T41.3X4A, T41.3X4D, T41.3X4S, T41.41XA, T41.41XD, T41.41XS, T41.44XA, T41.44XD, T41.44XS, T41.5X1D, T41.5X1D, T41.5X1S, T41.5X4A, T41.5X4D, T41.5X4S, T42.3X1A, T42.3X1D, T42.3X1S, T42.3X4A, T42.3X4D, T42.3X4S, T42.4X1A, T42.4X1D, T42.4X1S, T42.4X4A, T42.4X4D, T42.4X4S, T43.601A, T43.601D, T43.601S, T43.604A, T43.604D, T43.604S, T43.631D, T43.631D, T43.634D, T43.634D, T43.634D, T43.634D, T43.634D, T43.634D, T43.631D, T43.644D, T43.644S, T43.691A, T43.691D, T43.691S, T43.694A, T43.694D, T43.694S, T51.0X1D, T51.0X1D, T51.0X1S, T51.0X4A, T51.0X4D, T51.0X4S, T43.651D, T43.651D, T43.654S, T43.654D, T43.654S	To be used with appropriate visit codes for numerator compliance
BH Outpatient	CPT	98960, 98961, 98962, 99078, 99202, 99203, 99204, 99205, 99211, 99212, 99213, 99214, 99215, 99242, 99243, 99244, 99245, 99341, 99342, 99344, 99345, 99347, 99348, 99349, 99350, 99381, 99382, 99383, 99384, 99385, 99386, 99387, 99391, 99392, 99393, 99394, 99395, 99396, 99397, 99401, 99402, 99403, 99404, 99411, 99412, 99483, 99492, 99493, 99494, 99510	With diagnosis of SUD, substance use, or drug overdose <u>OR</u> with a mental health provider
BH Outpatient	HCPCS	G0155, G0176, G0177, G0409, G0463, G0512, H0002, H0004, H0031, H0034, H0036, H0037, H0039, H0040, H2000, H2010, H2011, H2013, H2014, H2015, H2016, H2017, H2018, H2019, H2020, T1015	With diagnosis of SUD, substance use, or drug overdose <b>OR</b> with a mental health provider
Partial Hospitalization or Intensive Outpatient	HCPCS	G0410, G0411, H0035, H2001, H2012, S0201, S9480, S9484, S9485	With diagnosis of SUD, substance use, or drug overdose <b>OR</b> with a mental health provider
Peer Support Services	HCPCS	G0140, G0177, H0024, H0025, H0038, H0039, H0040, H0046, H2014, H2023, S9445, T1012, T1016	With diagnosis of SUD, substance use, or drug overdose
OUD Monthly Office Based Treatment	HCPCS	G2086, G2087	With diagnosis of SUD, substance use, or drug overdose
OUD Weekly Non-Drug Service	HCPCS	G2071, G2074, G2075, G2076, G2077, G2080	With diagnosis of SUD, substance use, or drug overdose
Telephone Visits	CPT	98966, 98967, 98968, 99441, 99442, 99443	With diagnosis of SUD, substance use, or drug overdose <u>OR</u> with a mental health provider
Online Assessments	CPT	98971, 98972, 99421, 99422, 99423, 99457, 99458, 98980, 98981	With diagnosis of SUD, substance use, or drug overdose <u>OR</u> with a mental health provider
Online Assessments	HCPCS	G0071, G2010, G2012, G2250, G2251, G2252	With diagnosis of SUD, substance use, or drug overdose <u>OR</u> with a mental health provider
Substance Use Disorder Services	CPT	99408, 99409	
Substance Use Disorder Services	HCPCS	G0396, G0397, G0443, H0001, H0005, H0007, H0015, H0016, H0022, H0047, H0050, H2035, H2036, T1006, T1012	
Behavioral Health Assessment	CPT	99408, 99409	
Behavioral Health Assessment	HCPCS	G0396, G0397, G0442, G2011, H0001, H0002, H0031, H0049	

## $FUA \ ({\sf continued})$



HEDIS VALUE SET NAME	CODE SET	CODE	COMMENTS
OUD Weekly Drug Treatment Service	HCPCS	G2067, G2068, G2069, G2070, G2072, G2073	
Substance Use Services	HCPCS	H0006, H0028	
Outpatient POS	POS	03, 05, 07, 09, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 22, 33, 49, 50, 71, 72	With a Visit Setting Unspecified code <u>AND</u> with either any diagnosis of SUD, substance use, or drug overdose <u>OR</u> with a mental health provider
Non-residential Substance Abuse Treatment Facility POS	POS	57, 58	With a Visit Setting Unspecified code <b>AND</b> with either any diagnosis of SUD, substance use, or drug overdose <b>OR</b> with a mental health provider
Direct Reference Code	POS	52, 53	With a Visit Setting Unspecified code <b>AND</b> with either any diagnosis of SUD, substance use, or drug overdose <b>OR</b> with a mental health provider
Telehealth POS	POS	2, 10	With a Visit Setting Unspecified code <u>AND</u> with either any diagnosis of SUD, substance use, or drug overdose <u>OR</u> with a mental health provider
Substance Abuse Counseling and Surveillance	ICD10CM	Z71.41, Z71.51	Do not include laboratory claims (claims with POS code 81)

## $FUH \ (\hbox{continued})$



HEDIS VALUE SET NAME	CODE SET	CODE	COMMENTS
BH Outpatient	CPT	98960, 98961, 98962, 99078, 99202, 99203, 99204, 99205, 99211, 99212, 99213, 99214, 99215, 99242, 99243, 99244, 99245, 99341, 99342, 99344, 99345, 99347, 99348, 99349, 99350, 99381, 99382, 99383, 99384, 99385, 99386, 99387, 99391, 99392, 99393, 99394, 99395, 99396, 99397, 99401, 99402, 99403, 99404, 99411, 99412, 99483, 99492, 99493, 99494, 99510	With one of the following:  • A mental health provider  • Mental health disorder diagnosis  • POS code 53
BH Outpatient	HCPCS	G0155, G0176, G0177, G0409, G0463, G0512, H0002, H0004, H0031, H0034, H0036, H0037, H0039, H0040, H2000, H2010, H2011, H2013, H2014, H2015, H2016, H2017, H2018, H2019, H2020, T1015	With one of the following:  • A mental health provider  • Mental health disorder diagnosis  • POS code 53
Transitional Care Management Services	CPT	99495, 99496	With a mental health provider <b>OR</b> with POS code 53
Electroconvulsive Therapy	CPT	90870	With one of the following:  • Oupatient POS code  • POS code 24  • POS code 52  • POS code 53
Psychiatric Collaborative Care Management	CPT	99492, 99493, 99494	
Psychiatric Collaborative Care Management	HCPCS	G0512	
Telephone Visits	СРТ	98966, 98967, 98968, 99441, 99442, 99443	With a mental health provider
Direct Reference Code	POS	52	With a Visit Setting Unspecified code <b>OR</b> Electroconvulsive Therapy
Direct Reference Code	POS	53	With a mental health provider and Visit Setting Unspecified code <u>OR</u> Electroconvulsive Therapy <u>OR</u> Transitional Care
Direct Reference Code	POS	24	With Electroconvulsive Therapy
Direct Reference Code	POS	56	With Visit Setting Unspecified code
Telehealth POS	POS	2, 10	With a mental health provider and Visit Setting Unspecified code
Outpatient POS	POS	03, 05, 07, 09, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 22, 33, 49, 50, 71, 72	With one of the following:  • Visit Setting Unspecified code with either a mental health provider <b>OR</b> mental health disorder diagnosis  • POS code 53

## $FUH-\hbox{Follow-Up After Hospitalization for Mental Illness (continued)}$



HEDIS VALUE SET NAME	CODE SET	CODE	COMMENTS
Mental Health Diagnosis	ICD10CM	F03.90, F03.91, F20.0, F20.1, F20.2, F20.3, F20.5, F20.81, F20.89, F20.9, F21, F22, F23, F24, F25.0, F25.1, F25.8, F25.9, F28, F29, F30.10, F30.11, F30.12, F30.13, F30.2, F30.3, F30.4, F30.8, F30.9, F31.0, F31.10, F31.11, F31.12, F31.13, F31.2, F31.30, F31.31, F31.32, F31.4, F31.5, F31.60, F31.61, F31.62, F31.63, F31.64, F31.70, F31.71, F31.72, F31.73, F31.74, F31.75, F31.76, F31.77,	With one of the following:  • A mental health provider  • Mental health disorder diagnosis  • POS code 53
Mental Health Diagnosis	ICD10CM	F31.78, F31.81, F31.89, F31.9, F32.0, F32.1, F32.2, F32.3, F32.4, F32.5, F32.81, F32.89, F32.9, F32.A, F33.0, F33.1, F33.2, F33.3, F33.40, F33.41, F33.42, F33.8, F33.9, F34.0, F34.1, F34.81, F34.89, F34.9, F39, F40.00, F40.01, F40.02, F40.10, F40.11, F40.210, F40.218, F40.220, F40.228, F40.230, F40.231, F40.232, F40.233, F40.240, F40.241, F40.242, F40.243, F40.248, F40.290, F40.291, F40.298, F40.8, F40.9, F41.0, F41.1, F41.3, F41.8, F41.9, F42.2, F42.3, F42.4, F42.8, F42.9, F43.0, F43.10, F43.11, F43.12, F43.20, F43.21, F43.22, F43.23, F43.24, F43.25, F43.29, F43.8, F43.9, F44.0, F44.1, F44.2, F44.4, F44.5, F44.6, F44.7, F44.81, F44.89, F44.9, F45.0, F45.1, F45.20, F45.21, F45.22, F45.29, F45.41, F45.42, F45.8, F45.9, F48.1, F48.2, F48.8, F48.9, F50.00, F50.01, F50.02, F50.2, F50.81, F50.82, F50.89, F50.9, F51.01, F51.02, F51.03, F51.04, F51.05, F51.09, F51.11, F51.12, F51.13, F51.19, F51.3, F51.4, F51.5, F51.8, F51.9, F52.0, F52.1, F52.21, F52.22, F52.31, F52.32, F52.4, F52.5, F52.6, F52.8, F52.9, F53.0, F53.1, F59, F60.0, F60.1, F60.2, F60.3, F60.4, F60.5, F60.6, F60.7, F60.81, F60.89, F60.9, F63.0, F63.1, F63.2, F63.3, F63.81, F63.89, F63.9, F64.0, F64.1, F64.2, F64.8, F64.9, F65.0, F65.1, F65.2, F65.3, F65.4, F65.50, F65.51, F65.52, F65.81, F65.89, F65.9, F66, F68.10, F68.11, F68.12, F68.13, F68.8, F68.A, F69, F80.0, F80.1, F80.2, F80.4, F80.81, F80.82, F80.89, F80.9, F81.0, F81.2, F81.81, F81.89, F81.9, F82, F84.0, F84.2, F84.3, F84.5, F84.8, F84.9, F88, F89, F90.0, F90.1, F90.2, F90.8, F90.9, F91.0, F91.1, F91.2, F91.3, F91.8, F91.9, F93.0, F93.8, F93.9, F94.0, F94.1, F94.2, F94.8, F94.9, F98.5, F99.5, F98.8, F99.9, F99.03, P31.1, F03.918, F03.92, F03.93, F03.94, F03.80, F03.811, F03.818, F03.82, F03.83, F03.84, F03.C0, F03.C11, F03.C18, F03.C2, F03.C3, F03.C4, F43.81, F43.89	To be used with appropriate visit codes for numerator compliance
Peer Support Services	HCPCS	G0177, H0024, H0025, H0038, H0039, H0040, H0046, H2014, H2023, S9445, T1012, T1016, G0140	With a mental health disorder diagnosis
Residential Behavioral Health Treatment	HCPCS	T2048, H0019, H0017, H0018	

## $FUI \ ({\sf continued})$



HEDIS VALUE SET NAME	CODE SET	CODE	COMMENTS
AOD Abuse and Dependence	ICD10CM	F10.10, F10.120, F10.121, F10.129, F10.130, F10.131, F10.132, F10.139, F10.14, F10.150, F10.151, F10.159, F10.180, F10.181, F10.182, F10.188, F10.19, F10.20, F10.220, F10.221, F10.229, F10.230, F10.231, F10.232, F10.239, F10.24, F10.250, F10.251, F10.259, F10.26, F10.27, F10.280, F10.281, F10.282, F10.288, F10.29, F11.10, F11.120, F11.121, F11.122, F11.129, F11.13, F11.14, F11.150, F11.151, F11.159, F11.181, F11.182, F11.188, F11.19, F11.20, F11.220, F11.221, F11.222, F11.229, F11.23, F11.24, F11.250, F11.251, F11.259, F11.281, F11.282, F11.288, F11.29, F12.10, F12.120, F12.121, F12.122, F12.129, F12.13, F12.150, F12.151, F12.159, F12.180, F12.188, F12.19, F12.20, F12.220, F12.221, F12.222, F12.223, F12.236, F12.250, F12.251, F12.259, F12.280, F12.288, F12.29, F13.130, F13.131, F13.132, F13.139, F13.14, F13.150, F13.151, F13.159, F13.180, F13.181, F13.182, F13.188, F13.19, F13.20, F13.220, F13.221, F13.229, F13.230, F13.231, F13.232, F13.233, F13.24, F13.250, F13.251, F13.259, F13.26, F13.27, F13.280, F13.281, F13.282, F13.288, F13.29, F14.10, F14.120, F14.121, F14.122, F14.129, F14.13, F14.14, F14.150, F14.151, F14.159, F14.180, F14.181, F14.182, F14.188, F14.19, F14.20, F14.221, F14.222, F14.229, F14.23, F14.24, F14.250, F14.251, F14.259, F14.280, F14.281, F14.282, F14.288, F14.29, F15.10, F15.120, F15.121, F15.122, F15.129, F15.13, F15.14, F15.150, F15.151, F15.159, F15.180, F15.181, F15.182, F15.188, F15.19, F15.20, F15.221, F15.222, F15.229, F15.23, F15.24, F15.250, F15.251, F15.259, F16.29, F16.120, F16.120, F16.221, F16.129, F16.140, F16.150, F16.151, F16.159, F16.180, F16.180, F16.180, F16.180, F16.181, F16.180, F16.283, F16.29, F18.27, F18.280, F18.281, F18.29, F18.14, F19.150, F19.151, F19.159, F19.16, F19.17, F19.180, F19.181, F19.132, F19.139, F19.139, F19.14, F19.150, F19.151, F19.159, F19.16, F19.77, F19.180, F19.181, F19.182, F19.2	To be used with appropriate visit codes for numerator compliance
Visit Setting Unspecified	CPT	90791, 90792, 90832, 90833, 90834, 90836, 90837, 90838, 90839, 90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221, 99222, 99223, 99231, 99232, 99233, 99238, 99239, 99252, 99253, 99254, 99255	With a principle diagnosis of substance use disorder AND one of the following:  Outpatient POS code  POS code 52  POS code 53  Telehealth POS code  Non-residential substance abuse treatment facility POS
BH Outpatient	CPT	98960, 98961, 98962, 99078, 99202, 99203, 99204, 99205, 99211, 99212, 99213, 99214, 99215, 99242, 99243, 99244, 99245, 99341, 99342, 99344, 99345, 99347, 99348, 99349, 99350, 99381, 99382, 99383, 99384, 99385, 99386, 99387, 99391, 99392, 99393, 99394, 99395, 99396, 99397, 99401, 99402, 99403, 99404, 99411, 99412, 99483, 99492, 99493, 99494, 99510	With a principal diagnosis of substance use disorder
BH Outpatient	HCPCS	G0155, G0176, G0177, G0409, G0463, G0512, H0002, H0004, H0031, H0034, H0036, H0037, H0039, H0040, H2000, H2010, H2011, H2013, H2014, H2015, H2016, H2017, H2018, H2019, H2020, T1015	With a principal diagnosis of substance use disorder
Partial Hospitalization or Intensive Outpatient	HCPCS	G0410, G0411, H0035, H2001, H2012, S0201, S9480, S9484, S9485	With a principal diagnosis of substance use disorder
Substance Use Disorder Services	CPT	99408, 99409	With a principal diagnosis of substance use disorder
Substance Use Disorder Services	HCPCS	G0396, G0397, G0443, H0001, H0005, H0007, H0015, H0016, H0022, H0047, H0050, H2035, H2036, T1006, T1012	With a principal diagnosis of substance use disorder

## $FUI \ \hbox{(continued)}$



HEDIS VALUE SET NAME	CODE SET	CODE	COMMENTS
OUD Monthly Office Based Treatment	HCPCS	G2086, G2087	With a principal diagnosis of substance use disorder
OUD Weekly Non Drug Service	HCPCS	G2071, G2074, G2075, G2076, G2077, G2080	With a principal diagnosis of substance use disorder
Online Assessments	CPT	98970, 98971, 98972, 99421, 99422, 99423, 99457, 99458, 98980, 98981	With a principal diagnosis of substance use disorder
Online Assessments	HCPCS	G0071, G2010, G2012, G2250, G2251, G2252	With a principal diagnosis of substance use disorder
Residential Behavioral Health Treatment	HCPCS	H0017, H0018, H0019, T2048	With a principal diagnosis of substance use disorder
Telephone Visits	CPT	98966, 98967, 98968, 99441, 99442, 99443	With a principal diagnosis of substance use disorder
AOD Medication Treatment	HCPCS	G2069, G2070, G2072, G2073, H0020, H0033, J0570, J0571, J0572, J0573, J0574, J0575, J0577, J0578 J2315, Q9991, Q9992, S0109	
OUD Weekly Drug Treatment Service	HCPCS	G2067, G2068, G2069, G2070, G2072, G2073	
Outpatient POS	POS	03, 05, 07, 09, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 22, 33, 49, 50, 71, 72	With a Visit Setting Unspecified code <u>AND</u> a principal diagnosis of substance abuse disorder
Non-residential Substance Abuse Treatment Facility POS	POS	57, 58	With a Visit Setting Unspecified code AND a principal diagnosis of substance abuse disorder
Direct Reference Code	POS	52, 53	With a Visit Setting Unspecified code <b>AND</b> a principal diagnosis of substance abuse disorder
Telehealth POS	POS	2, 10	With a Visit Setting Unspecified code AND a principal diagnosis of substance abuse disorder
Substance Abuse Counseling and Surveillance	ICD10CM	Z71.41, Z71.51	With a principal diagnosis of substance abuse disorder. Do not include laboratory claims (claims with POS code 81)





HEDIS VALUE SET NAME	CODE SET	CODE	COMMENTS
Mental Health Diagnosis	ICD10CM	F03.90, F03.91, F20.0, F20.1, F20.2, F20.3, F20.5, F20.81, F20.89, F20.9, F21, F22, F23, F24, F25.0, F25.1, F25.8, F25.9, F28, F29, F30.10, F30.11, F30.12, F30.13, F30.2, F30.3, F30.4, F30.8, F30.9, F31.0, F31.10, F31.11, F31.12, F31.13, F31.2, F31.30, F31.31, F31.32, F31.4, F31.5, F31.60, F31.61, F31.62, F31.63, F31.64, F31.70, F31.71, F31.72, F31.73, F31.74, F31.75, F31.76, F31.77, F31.78, F31.81, F31.89, F31.9, F32.0, F32.1, F32.2, F32.3, F32.4, F32.5, F32.81, F32.89, F32.9, F32.4, F33.0, F33.1, F33.2, F33.3, F33.40, F33.41, F33.42, F33.8, F33.9, F34.0, F34.1, F34.81, F34.89, F34.9, F39, F40.00, F40.01, F40.02, F40.10, F40.11, F40.210, F40.218, F40.220, F40.228, F40.230, F40.231, F40.232, F40.233, F40.240, F40.241, F40.242, F40.243, F40.249, F40.290, F40.291, F40.298, F40.8, F40.9, F41.0, F41.1, F41.3, F41.8, F41.9, F42.2, F42.3, F42.4, F42.8, F42.9, F43.0, F43.10, F43.11, F43.12, F43.20, F43.21, F43.22, F43.23, F43.24, F43.25, F43.29, F43.8, F43.9, F44.0, F44.1, F44.2, F44.4, F44.5, F44.6, F44.7, F44.81, F44.89, F44.9, F45.0, F45.11, F45.20, F45.21, F45.22, F50.2, F50.81, F50.82, F50.89, F50.9, F51.01, F51.03, F51.04, F51.04, F51.05, F51.09, F51.11, F51.12, F51.13, F51.19, F51.3, F51.4, F51.5, F51.8, F51.9, F52.0, F52.1, F52.22, F52.31, F52.32, F52.34, F52.54, F52.58, F52.8, F52.9, F53.0, F53.1, F59, F60.0, F60.1, F60.2, F60.3, F63.89, F63.9, F64.0, F64.1, F64.2, F64.8, F64.9, F65.0, F65.1, F65.2, F65.3, F65.4, F65.50, F65.51, F65.52, F65.81, F65.89, F65.9, F66, F68.10, F68.11, F68.12, F68.13, F68.8, F68.8, F69.9, F80.0, F80.1, F80.2, F80.4, F80.81, F80.82, F80.89, F80.9, F81.0, F81.2, F81.81, F81.89, F81.9, F82.79, F95.0, F95.11, F90.2, F90.8, F90.9, F91.0, F91.1, F91.2, F91.3, F91.8, F91.9, F93.0, F93.8, F93.9, F94.0, F94.1, F94.2, F94.8, F94.9, F95.0, F95.1, F95.2, F95.8, F95.9, F98.0, F98.1, F98.21, F98.29, F98.3, F98.8, F99.9, F93.0, F93.11, F03.918, F03.92, F03.93, F03.94, F03.00, F03.A11, F03.A18, F03.A2, F03.A3, F03.A4. F03.B0, F03.B11, F03.B18, F03.B2, F03.B3, F03.B4, F03.C0, F03.C11, F03.C18, F03.	To be used with appropriate visit codes for numerator compliance
Visit Setting Unspecified	СРТ	90791, 90792, 90832, 90833, 90834, 90836, 90837, 90838, 90839, 90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221, 99222, 99223, 99231, 99232, 99233, 99238, 99239, 99252, 99253, 99254, 99255	With one of the following:  Diagnosis of a metal health disorder  POS code 52  POS code 53  POS code 56  Telehealth POS code
BH Outpatient	CPT	98960, 98961, 98962, 99078, 99202, 99203, 99204, 99205, 99211, 99212, 99213, 99214, 99215, 99242, 99243, 99244, 99245, 99341, 99342, 99344, 99345, 99347, 99348, 99349, 99350, 99381, 99382, 99383, 99384, 99385, 99386, 99387, 99392, 99393, 99394, 99395, 99396, 99397, 99401, 99402, 99403, 99404, 99411, 99412, 99483, 99492, 99493, 99494, 99510	With any diagnosis of a mental health disorder OR with any diagnosis of intentional self-harm with any diagnosis of a mental health disorder
BH Outpatient	HCPCS	G0155, G0176, G0177, G0409, G0463, G0512, H0002, H0004, H0031, H0034, H0036, H0037, H0039, H0040, H2000, H2010, H2011, H2013, H2014, H2015, H2016, H2017, H2018, H2019, H2020, T1015	With any diagnosis of a mental health disorder OR with any diagnosis of intentional self-harm with any diagnosis of a mental health disorder
Partial Hospitalization or Intensive Outpatient	HCPCS	G0410, G0411, H0035, H2001, H2012, S0201, S9480, S9484, S9485	With a diagnosis of a mental health disorder

## $FUM \ \hbox{(continued)}$



HEDIS VALUE SET NAME	CODE SET	CODE	COMMENTS
Telephone Visits	CPT	98966, 98967, 98968, 99441, 99442, 99443	With a diagnosis of a mental health disorder
Online Assessments	CPT	98970, 98971, 98972, 99421, 99422, 99423, 99457, 99458, 98980, 98981	With a diagnosis of a mental health disorder
Online Assessments	HCPCS	G0071, G2010, G2012, G2250, G2251, G2252	With a diagnosis of a mental health disorder
Electroconvulsive Therapy	CPT	90870	With one of the following:  • Oupatient POS  • POS code 24  • POS code 52  • POS code 53
Electroconvulsive Therapy	ICD10PCS	GZB0ZZZ, GZB1ZZZ, GZB2ZZZ, GZB3ZZZ, GZB4ZZZ	With one of the following:  • Oupatient POS  • POS code 24  • POS code 52  • POS code 53
Outpatient POS	POS	03, 05, 07, 09, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 22, 33, 49, 50, 71, 72	With a Visit Setting Unspecified code <u>AND</u> diagnosis of a mental health disorder
Direct Reference Code	POS	52, 53	With a Visit Setting Unspecified code <b>OR</b> an Electroconvulsive therapy code
Direct Reference Code	POS	24	With a Visit Setting Unspecified code
Telehealth POS	POS	2, 10	With a Visit Setting Unspecified code <u>AND</u> a diagnosis of a mental health disorder
Direct Reference Code	POS	56	With a Visit Setting Unspecified code
Psychiatric Collaborative Care Management	СРТ	99492, 99493, 99494, G0512	
Peer Support Services	HCPCS	G0177, H0024, H0025, H0038, H0039, H0040, H0046, H2014, H2023, S9445, T1012, T1016, G0140	With a mental health disorder diagnosis
Residential Behavioral Health Treatment	HCPCS	T2048, H0019, H0017, H0018	





HEDIS VALUE SET NAME	CODE SET	CODE	COMMENTS
OUD Monthly Office Based Treatment	HCPCS	G2086, G2087	
Alcohol Abuse and Dependence	ICD10CM	F10.10, F10.120, F10.121, F10.129, F10.130, F10.131, F10.132, F10.139, F10.14, F10.150, F10.151, F10.159, F10.180, F10.181, F10.182, F10.188, F10.19, F10.20, F10.220, F10.221, F10.229, F10.230, F10.231, F10.232, F10.239, F10.24, F10.250, F10.251, F10.259, F10.26, F10.27, F10.280, F10.281, F10.282, F10.288, F10.29	To be used with appropriate visit codes for numerator compliance
Opioid Abuse and Dependence	ICD10CM	F11.10, F11.120, F11.121, F11.122, F11.129, F11.13, F11.14, F11.150, F11.151, F11.159, F11.181, F11.182, F11.188, F11.19, F11.20, F11.220, F11.221, F11.222, F11.229, F11.23, F11.24, F11.250, F11.251, F11.259, F11.281, F11.282, F11.288, F11.29	To be used with appropriate visit codes for numerator compliance
Other Drug Abuse and Dependence	ICD10CM	F12.10, F12.120, F12.121, F12.122, F12.129, F12.13, F12.150, F12.151, F12.159, F12.180, F12.188, F12.19, F12.20, F12.220, F12.221, F12.222, F12.229, F12.23, F12.250, F12.251, F12.259, F12.280, F12.288, F12.29, F13.10, F13.120, F13.121, F13.129, F13.130, F13.131, F13.132, F13.139, F13.14, F13.150, F13.151, F13.159, F13.180, F13.181, F13.182, F13.188, F13.29, F13.20, F13.220, F13.221, F13.229, F13.230, F13.231, F13.232, F13.239, F13.24, F13.250, F13.251, F13.259, F13.26, F13.27, F13.280, F13.281, F13.282, F13.288, F13.29, F14.10, F14.120, F14.121, F14.122, F14.129, F14.13, F14.14, F14.150, F14.151, F14.159, F14.180, F14.181, F14.182, F14.188, F14.19, F14.20, F14.220, F14.221, F14.222, F14.229, F14.23, F14.24, F14.250, F14.251, F14.259, F14.280, F14.281, F14.282, F14.288, F14.29, F15.10, F15.120, F15.121, F15.122, F15.129, F15.13, F15.14, F15.150, F15.151, F15.159, F15.180, F15.181, F15.182, F15.188, F15.19, F15.20, F15.220, F15.221, F15.222, F15.229, F15.23, F15.24, F15.250, F15.251, F15.259, F15.280, F15.281, F15.282, F15.288, F15.29, F16.10, F16.120, F16.121, F16.122, F16.129, F16.14, F16.150, F16.151, F16.159, F16.180, F16.183, F16.188, F16.19, F16.20, F16.220, F16.221, F16.229, F16.24, F16.250, F16.251, F16.259, F16.283, F16.288, F16.29, F18.10, F18.120, F18.121, F18.129, F18.14, F18.150, F18.151, F18.159, F18.17, F18.180, F18.188, F18.19, F18.20, F18.220, F18.221, F18.229, F18.24, F18.250, F18.251, F18.259, F18.27, F18.280, F18.288, F18.29, F19.10, F19.120, F19.121, F19.122, F19.129, F19.130, F19.131, F19.132, F19.139, F19.14, F19.150, F19.151, F19.159, F19.16, F19.17, F19.180, F19.181, F19.182, F19.188, F19.19, F19.20, F19.221, F19.222, F19.222, F19.229, F19.230, F19.231, F19.232, F19.239, F19.24, F19.250, F19.251, F19.259, F19.26, F19.27, F19.280, F19.281, F19.282, F19.288, F19.29	To be used with appropriate visit codes for numerator compliance
Visit Setting Unspecified	CPT	90791, 90792, 90832, 90833, 90834, 90836, 90837, 90838, 90839, 90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221, 99222, 99223, 99231, 99232, 99233, 99238, 99239, 99252, 99253, 99254, 99255	With a diagnosis of Alcohol Abuse / Dependence OR Opioid Abuse / Dependence OR Other Drug Abuse / Dependence AND one of the following:  Outpatient POS code POS code 52 POS code 53 Telehealth POS code Non-residential substance abuse treatment facility POS
BH Outpatient	CPT	98960, 98961, 98962, 99078, 99202, 99203, 99204, 99205, 99211, 99212, 99213, 99214, 99215, 99242, 99243, 99244, 99245, 99341, 99342, 99344, 99345, 99347, 99348, 99349, 99350, 99381, 99382, 99383, 99384, 99385, 99386, 99387, 99391, 99392, 99393, 99394, 99395, 99396, 99397, 99401, 99402, 99403, 99404, 99411, 99412, 99483, 99492, 99493, 99494, 99510	With a diagnosis of Alcohol Abuse / Dependence <u>OR</u> Opioid Abuse / Dependence <u>OR</u> Other Drug Abuse / Dependence





HEDIS VALUE SET NAME	CODE SET	CODE	COMMENTS
BH Outpatient	HCPCS	G0155, G0176, G0177, G0409, G0463, G0512, H0002, H0004, H0031, H0034, H0036, H0037, H0039, H0040, H2000, H2010, H2011, H2013, H2014, H2015, H2016, H2017, H2018, H2019, H2020, T1015	With a diagnosis of Alcohol Abuse / Dependence <u>OR</u> Opioid Abuse / Dependence <u>OR</u> Other Drug Abuse / Dependence
Partial Hospitalization or Intensive Outpatient	HCPCS	G0410, G0411, H0035, H2001, H2012, S0201, S9480, S9484, S9485	With a diagnosis of Alcohol Abuse / Dependence <u>OR</u> Opioid Abuse / Dependence <u>OR</u> Other Drug Abuse / Dependence
Telephone Visits	CPT	98966, 98967, 98968, 99441, 99442, 99443	With a diagnosis of Alcohol Abuse / Dependence <u>OR</u> Opioid Abuse / Dependence <u>OR</u> Other Drug Abuse / Dependence
Online Assessments	CPT	98970, 98971, 98972, 99421, 99422, 99423, 99457, 99458, 98980, 98981	With a diagnosis of Alcohol Abuse / Dependence <u>OR</u> Opioid Abuse / Dependence <u>OR</u> Other Drug Abuse / Dependence
Online Assessments	HCPCS	G0071, G2010, G2012, G2250, G2251, G2252	With a diagnosis of Alcohol Abuse / Dependence <u>OR</u> Opioid Abuse / Dependence <u>OR</u> Other Drug Abuse / Dependence
Substance Use Disorder Services	CPT	99408, 99409	With a diagnosis of Alcohol Abuse / Dependence <u>OR</u> Opioid Abuse / Dependence <u>OR</u> Other Drug Abuse / Dependence
Substance Use Disorder Services	HCPCS	G0396, G0397, G0443, H0001, H0005, H0007, H0015, H0016, H0022, H0047, H0050, H2035, H2036, T1006, T1012	With a diagnosis of Alcohol Abuse / Dependence <u>OR</u> Opioid Abuse / Dependence <u>OR</u> Other Drug Abuse / Dependence
OUD Weekly Non-Drug Service	HCPCS	G2071, G2074, G2075, G2076, G2077, G2080	With a diagnosis of Alcohol Abuse / Dependence <u>OR</u> Opioid Abuse / Dependence <u>OR</u> Other Drug Abuse / Dependence
OUD Monthly Office Based Treatment	HCPCS	G2086, G2087	Numerator Compliance for Engagement Only
Buprenorphine Implant	HCPCS	G2070, G2072, J0570	Numerator compliance for opioid use disorder cohort for Initiation rate only
Buprenorphine Injection	HCPCS	G2069, Q9991, Q9992	Numerator compliance for opioid use disorder cohort for Initiation rate only
Buprenorphine Naloxone	HCPCS	J0572, J0573, J0574, J0575	Numerator compliance for opioid use disorder cohort only
Buprenorphine Oral	HCPCS	H0033, J0571	Numerator compliance for opioid use disorder cohort only
Buprenorphine Oral Weekly	HCPCS	G2068, G2079	Numerator compliance for opioid use disorder cohort only
Direct Reference Code	POS	52, 53	With a Visit Setting Unspecified Code AND one of the following diagnoses: Alcohol Abuse / Dependence OR Opioid Abuse / Dependence OR Other Drug Abuse / Dependence
Methadone Oral	HCPCS	H0020, S0109, 310653000	Numerator compliance for opioid use disorder cohort only
Methadone Oral Weekly	HCPCS	G2067, G2078	Numerator compliance for opioid use disorder cohort only





HEDIS VALUE SET NAME	CODE SET	CODE	COMMENTS
Naltrexone Injection	HCPCS	J2315	Numerator compliance for Initiation rate only
Non-residential Substance Abuse Treatment Facility POS	POS	57, 58	With a Visit Setting Unspecified Code <u>AND</u> one of the following diagnoses: Alcohol Abuse / Dependence <u>OR</u> Opioid Abuse / Dependence <u>OR</u> Other Drug Abuse / Dependence
OUD Weekly Drug Treatment Service	HCPCS	G2067, G2068, G2069, G2070, G2072, G2073	
Outpatient	CPT	99202, 99203, 99204, 99205, 99211, 99212, 99213, 99214, 99215, 99241, 99242, 99243, 99244, 99245, 99341, 99342, 99343, 99344, 99345, 99347, 99348, 99349, 99350, 99381, 99382, 99383, 99384, 99385, 99386, 99387, 99391, 99392, 99393, 99394, 99395, 99396, 99397, 99401, 99402, 99403, 99404, 99411, 99412, 99429, 99455, 99456, 99483	With a Visit Setting Unspecified Code AND one of the following diagnoses: Alcohol Abuse / Dependence OR Opioid Abuse / Dependence OR Other Drug Abuse / Dependence
Outpatient	HCPCS	G0402, G0438, G0439, G0463, T1015, 77406008	With a Visit Setting Unspecified Code <u>AND</u> one of the following diagnoses: Alcohol Abuse / Dependence <u>OR</u> Opioid Abuse / Dependence <u>OR</u> Other Drug Abuse / Dependence
Substance Abuse Counseling and Surveillance	ICD10CM	Z71.41, Z71.51	With a diagnosis of Alcohol Abuse / Dependence OR Opioid Abuse / Dependence OR Other Drug Abuse / Dependence. Do not include laboratory claims (claims with POS code 81)
Telehealth	POS	02, 10	With a Visit Setting Unspecified Code AND one of the following diagnoses: Alcohol Abuse / Dependence OR Opioid Abuse / Dependence OR Other Drug Abuse / Dependence





HEDIS VALUE SET NAME	CODE SET	CODE	COMMENTS
Prenatal Bundled Services	CPT	59400, 59425, 59426, 59510, 59610, 59618	
Prenatal Bundled Services	HCPCS	H1005	
Stand Alone Prenatal Visits	СРТ	99500	
Stand Alone Prenatal Visits	CPT-CAT-II	0500F, 0501F, 0502F	Do not include codes with any of these CPT II Modifiers: 1P, 2P, 3P, 8P
Stand Alone Prenatal Visits	HCPCS	H1000, H1001, H1002, H1003, H1004	
Prenatal Visits	СРТ	99202, 99203, 99204, 99205, 99211, 99212, 99213, 99215, 99241, 99242, 99243, 99244, 99245, 99483, 98966, 98967, 98968, 98970, 98971, 98972, 98980, 98981, 99421, 99422, 99423, 99441, 99442, 99443, 99457, 99458	With a pregnancy-related diagnosis code
Prenatal Visits	HCPCS	G0463, T1015, G0071, G2010, G2012, G2250, G2251, G2252	With a pregnancy-related diagnosis code
Pregnancy Diagnosis	ICD10CM	09.00, 009.01, 009.02, 009.03, 009.10, 009.11, 009.12, 009.13, 009.211, 009.212, 009.213, 009.219, 009.291, 009.292, 009.293, 009.299, 009.30, 009.31, 009.32, 009.33, 009.40, 009.41, 009.42, 009.43, 009.511, 009.512, 009.513, 009.519, 009.521, 009.522, 009.523, 009.529, 009.611, 009.612, 009.613, 009.619, 009.622, 009.623, 009.629, 009.70, 009.71, 009.72, 009.73, 009.811, 009.812, 009.813, 009.819, 009.819, 009.821, 009.822, 009.829, 009.829, 009.891, 009.892, 009.893, 009.80, 009.91, 009.92, 009.93, 009.80, 009.41, 009.A2, 009.A3, 010.011, 010.012, 010.013, 010.019, 010.111, 010.112, 010.113, 010.119, 010.211, 010.212, 010.213, 010.219, 010.311, 010.312, 010.313, 010.319, 010.411, 010.412, 010.413, 010.419, 010.911, 010.912, 010.913, 010.919, 011.1, 011.2, 011.3, 011.9, 012.20, 012.20, 101.20, 101.20, 101.20, 101.20, 101.20, 101.20, 101.20, 012.10, 101.21, 012.21, 012.22, 012.23, 013.1, 013.2, 013.3, 013.9, 014.00, 014.02, 014.02, 014.02, 014.02, 014.02, 014.92, 014.92, 014.92, 014.93, 015.00, 015.02, 015.03, 015.1, 015.9, 016.1, 016.2, 016.3, 016.9, 020.0, 020.8, 020.9, 021.0, 021.1, 021.2, 021.8, 021.9, 022.00, 022.01, 022.30, 022.31, 022.32, 022.33, 022.40, 022.41, 022.42, 022.43, 022.50, 022.51, 022.52, 022.53, 022.831, 022.32, 022.33, 022.40, 022.41, 022.42, 022.43, 022.50, 022.51, 022.52, 022.53, 022.831, 023.39, 023.11, 023.12, 023.13, 023.20, 023.21, 023.22, 023.33, 023.30, 023.11, 023.12, 023.13, 023.20, 023.21, 023.22, 023.23, 023.30, 023.11, 023.12, 023.13, 023.593, 023.599, 023.599, 023.591, 023.592, 023.599, 023.599, 023.591, 023.592, 023.599, 023.599, 023.591, 023.512, 023.513, 023.512, 023.513, 023.512, 023.513, 023.512, 023.513, 023.512, 023.513, 023.512, 023.522, 023.523, 023.529, 023.591, 023.592, 023.599, 023.599, 023.511, 024.419, 024.411, 024.412, 024.413, 024.419, 024.411, 024.412, 024.413, 024.419, 024.411, 024.412, 024.413, 024.419, 024.411, 024.412, 024.413, 024.419, 024.411, 024.412, 024.413, 024.812, 024.813, 024.819, 024.911, 024.912, 024.919, 025.611, 026.612, 026.613, 026.614, 026	To be used with appropriate visit codes for numerator compliance





HEDIS VALUE SET NAME	CODE SET	CODE	COMMENTS
Pregnancy Diagnosis (continued)	ICD10CM	030.099, 030.101, 030.102, 030.103, 030.109, 030.111, 030.112, 030.113, 030.119, 030.119, 030.120, 030.120, 030.120, 030.123, 030.129, 030.131, 030.190, 030.103, 030.129, 030.213, 030.129, 030.130, 030.190, 030.103, 030.219, 030.213, 030.129, 030.131, 030.129, 030.131, 030.129, 030.130, 030.229, 030.230, 030.229, 030.231, 030.229, 030.231, 030.229, 030.230, 030.239, 030.241, 030.221, 030.222, 030.232, 030.239, 030.239, 030.241, 030.2421, 030.221, 030.222, 030.223, 030.229, 030.831, 030.839, 030.831, 030.839, 030.831, 030.839, 030.839, 030.839, 030.839, 030.839, 030.839, 030.839, 030.839, 030.839, 030.831, 030.832, 030.839, 03	With a prenatal visit OR Telephone Visit OR Online Assessment





HEDIS VALUE SET NAME	CODE SET	CODE	COMMENTS
Pregnancy Diagnosis (continued)	ICD10CM	036.22X0, 036.22X1, 036.22X2, 036.22X3, 036.22X4, 036.22X5, 036.22X9, 036.23X0, 036.23X0, 036.23X1, 036.23X2, 036.23X2, 036.23X3, 036.23X4, 036.23X3, 036.4XX1, 036.4XX1, 036.4XX1, 036.4XX1, 036.4XX1, 036.4XX1, 036.4XX1, 036.4XX1, 036.4XX1, 036.5X4X, 036.4XX1, 036.5X12, 036.5X13, 036.5X12, 036.6X12, 036.5X12, 036.5X	With a prenatal visit <u>OR</u> Telephone Visit <u>OR</u> Online Assessment





HEDIS VALUE SET NAME	CODE SET	CODE	COMMENTS
Pregnancy Diagnosis (continued)	ICD10CM	043.023, 043.029, 043.101, 043.102, 043.103, 043.109, 043.111, 043.112, 043.113, 043.119, 043.121, 043.122, 043.123, 043.129, 043.221, 043.222, 043.233, 043.239, 043.231, 043.222, 043.231, 043.222, 043.231, 043.222, 043.231, 043.232, 043.339, 043.839, 043.831, 043.831, 043.831, 043.831, 043.839, 043.839, 043.839, 043.831, 043.831, 043.831, 043.839, 043.851, 043.822, 044.23, 044.20, 044.41, 044.12, 044.13, 044.20, 044.51, 044.52, 044.53, 045.002, 045.003, 045.009, 045.001, 044.002, 044.50, 044.51, 044.52, 044.50, 044.51, 044.52, 044.50, 044.52, 045.022, 045.022, 045.023, 045.093, 045.091, 045.022, 045.033, 045.009, 045.011, 045.012, 045.013, 045.019, 045.022, 045.033, 046.001, 046.002, 046.003, 04	With a prenatal visit OR Telephone Visit OR Online Assessment
Cervical Cytology Lab Test	CPT	88141, 88142, 88143, 88147, 88148, 88150, 88152, 88153, 88164, 88165, 88166, 88167, 88174, 88175	
Cervical Cytology Lab Test	HCPCS	G0123, G0124, G0141, G0143, G0144, G0145, G0147, G0148, P3000, P3001, Q0091	
Postpartum Bundled Services	CPT	59400, 59410, 59510, 59515, 59610, 59614, 59618, 59622	

## $PPC \ (\hbox{continued})$



HEDIS VALUE SET NAME	CODE SET	CODE	COMMENTS
Postpartum Care	CPT	57170, 58300, 59430, 99501	
Postpartum Care	CPT-CAT-II	0503F	Do not include codes with any of these CPT II Modifiers: 1P, 2P, 3P, 8P
Postpartum Care	HCPCS	G0101	
Encounter for Postpartum Care	ICD10CM	Z01.411, Z01.419, Z01.42, 30.430, Z39.1, Z39.2	Do not include laboratory claims (claims with POS code 81)

