

Corporate Medical Policy

Genetic Testing for Epilepsy AHS – M2075

File Name: genetic_testing_for_epilepsy
Origination: 01/01/2019
Last Review: 5/2023

Description of Procedure or Service

Epilepsy is a group of disorders characterized by recurrent, unprovoked seizures due to abnormal, synchronized neuronal firing in the brain that can be distinguished by seizure type, age of onset, developmental status, co-morbid features and etiology (Berg et al., 2010; Myers & Mefford, 2015).

Related Policies

General Genetic Testing Germline Disorders
General Genetic Testing Somatic Disorders
Genetic Testing of Mitochondrial Disorders

Regulatory Status

Food and Drug Administration (FDA)

Many labs have developed specific tests that they must validate and perform in house. These laboratory-developed tests (LDTs) are regulated by the Centers for Medicare and Medicaid (CMS) as high-complexity tests under the Clinical Laboratory Improvement Amendments of 1988 (CLIA '88). LDTs are not approved or cleared by the U. S. Food and Drug Administration; however, FDA clearance or approval is not currently required for clinical use.

******Note: This Medical Policy is complex and technical. For questions concerning the technical language and/or specific clinical indications for its use, please consult your physician.***

Policy

BCBSNC will provide coverage for genetic testing for epilepsy when it is determined the medical criteria or reimbursement guidelines below are met.

Benefits Application

This medical policy relates only to the services or supplies described herein. Please refer to the Member's Benefit Booklet for availability of benefits. Member's benefits may vary according to benefit design; therefore member benefit language should be reviewed before applying the terms of this medical policy.

When Genetic Testing for Epilepsy is covered

Genetic Testing for Epilepsy AHS – M2075

Genetic testing for mutations associated with infantile- and early childhood-onset epilepsy syndromes (see Note 1) for individuals with infantile- and early-childhood-onset epilepsy syndromes (in which epilepsy is the core clinical symptom) is considered medically necessary only when a positive test result would lead to changes in at least **one** of the following:

- Medication management.
- Diagnostic testing (alternative potentially invasive tests are avoided).
- Reproductive decision making.

Targeted mutation testing is considered medically necessary for individuals with an epilepsy syndrome with a suspected genetic cause, to assess for the following clinical situations:

- *ALDH7A1* testing for pyridoxine-related epilepsy.
- *BTD* testing for biotinidase deficiency-related epilepsy.
- *FOLR1* testing for cerebral folate deficiency-related epilepsy.
- *GRIN2A* testing for GRIN2A-related epileptic encephalopathy.
- *KCNQ2* testing for KCNQ2-related epileptic encephalopathy.
- *KCNT1* testing for KCNT1-related migrating partial epilepsy of infancy.
- *PCDH19* testing for epilepsy female-restricted with mental retardation (EFMR).
- *PHGDH*, *PSAT1*, and *PSPH* testing for epilepsy due to serine biosynthesis defects.
- *SCN1A* testing for SCN1A-related seizure disorders.
- *SCN8A* testing for SCN8A-related epileptic encephalopathy.
- *SLC2A1* testing for glucose transporter type 1 deficiency syndrome.
- *SLC6A8*, *GATM*, and *GAMT* testing for epilepsy due to creatine deficiency syndromes.
- *TSC1* and *TSC2* testing for tuberous sclerosis complex-related epilepsy.

When Genetic Testing for Epilepsy is not covered

Genetic testing for epilepsy is considered investigational in all other clinical situations not described above and for any other mutations not listed above.

Note 1: For 5 or more gene tests being run on the same platform, please refer to Laboratory Procedures Reimbursement Policy AHS-R2162.

Table of Terminology

Term	Definition
AAN	American Academy of Neurology
<i>ALDH7A1</i>	<i>Aldehyde dehydrogenase 7 family member A1</i>
AMA	Antiseizure medication
<i>ANK2</i>	<i>Ankyrin 2</i>
ASD	Autism spectrum disorders
<i>BTD</i>	<i>Biotinidase</i>
<i>CACNA1A</i>	<i>Calcium voltage-gated channel subunit alpha 1A</i>
<i>CACNA1AE</i>	<i>Calcium voltage-gated channel subunit alpha 1AE</i>
<i>CACNA1D</i>	<i>Calcium voltage-gated channel subunit alpha 1D</i>
<i>CACNA1H</i>	<i>Calcium voltage-gated channel subunit alpha 1H</i>
<i>CACNA2D3</i>	<i>Calcium voltage-gated channel subunit alpha 1A</i>
CDKL5	Calcium voltage-gated channel subunit alpha2delta3
CGH	Comparative genomic hybridization

Genetic Testing for Epilepsy AHS – M2075

<i>CHD2</i>	<i>Chromodomain helicase DNA binding protein 2</i>
CLIA '88	Clinical Laboratory Improvement Amendments of 1988
CMA	Chromosomal microarray analysis
CMS	Centers for Medicare and Medicaid Services
CNS	Child Neurology Society
CNV	Copy number variant
DEE	Developmental and epileptic encephalopathy
<i>DLG4</i>	<i>Discs large homolog 4</i>
DS	Dravet syndrome
EE	Epileptic encephalopathy
EEG	Electroencephalography
EFMR	Epilepsy female-restricted with mental retardation
EFNS	European Federation of Neurological Societies
EP	Epileptic panels
<i>FGF12</i>	<i>Fibroblast growth factor 12</i>
<i>FOLR1</i>	<i>Folate receptor 1</i>
<i>GABRA1</i>	<i>Gamma-aminobutyric acid type A receptor alpha1</i>
<i>GABRG2</i>	<i>Gamma-aminobutyric acid type A receptor gamma2</i>
<i>GAMT</i>	<i>Guanidinoacetate n-methyltransferase</i>
<i>GATM</i>	<i>Glycine amidinotransferase</i>
GGE	Genetic generalized epilepsy
Glut1DS	Glucose transporter 1 deficiency syndrome
<i>GRIA2</i>	<i>Glutamate ionotropic receptor AMPA type subunit 2</i>
<i>GRIN2A</i>	<i>Glutamate ionotropic receptor NMDA type subunit 2a</i>
<i>GRIN2B</i>	<i>Glutamate ionotropic receptor NMDA type subunit 2b</i>
GS	Genome sequencing
GTAC	Genetic Testing Advisory Committee
<i>HNRNPU</i>	<i>Heterogeneous nuclear ribonucleoprotein U</i>
IC-CODE	International Classification of Cognitive Disorders in Epilepsy
ID	Intellectual disability
ILAE	International League Against Epilepsy
<i>KANSL1</i>	<i>KAT8 regulatory NSL complex subunit 1</i>
KB	Kilo-base pair
<i>KCNC1</i>	<i>Potassium voltage-gated channel subfamily C member 1</i>
<i>KCNQ2</i>	<i>Potassium voltage-gated channel subfamily Q member 2</i>
<i>KCNT1</i>	<i>Potassium sodium-activated channel subfamily T member 1</i>
KD	Ketogenic diet
LDTs	Laboratory-developed tests
<i>MECP2</i>	<i>Methyl-CpG binding protein 2</i>
MGP	Multigene panel
mTOR	Mammalian target of rapamycin

Genetic Testing for Epilepsy AHS – M2075

NDD	Neurodevelopmental disorders
NGS	Next-generation sequencing
NICE	National Institute for Health and Care Excellence
<i>PCDH19</i>	<i>Protocadherin 19</i>
<i>PHGDH</i>	<i>Phosphoglycerate dehydrogenase</i>
<i>PNPO</i>	<i>Pyridoxamine 5'-phosphate oxidase</i>
PRRT2	Proline-rich transmembrane protein 2
<i>PSAT1</i>	<i>Phosphoserine aminotransferase 1</i>
<i>PSPH</i>	<i>Phosphoserine phosphatase</i>
<i>PURA</i>	<i>Pur-alpha</i>
<i>SCN1A</i>	<i>Sodium voltage-gated channel alpha subunit 1</i>
<i>SCN2A</i>	<i>Sodium voltage-gated channel type 2 alpha subunit</i>
<i>SCN8A</i>	<i>Sodium voltage-gated channel type 8 alpha subunit</i>
SE	Status epilepticus
SIGN	Scottish Intercollegiate Guidelines Network
<i>SLC2A1</i>	<i>Solute carrier family 2 member 1</i>
<i>SLC6A1</i>	<i>Solute carrier family 6 member 1</i>
<i>SLC6A8</i>	<i>Solute carrier family 6 member 8</i>
SMEI	Severe myoclonic epilepsy of infancy
<i>STX1B</i>	<i>Syntaxin 1B</i>
STXBP1	Syntaxin-binding protein 1
TCF4	Transcription factor 4
TSC	Tuberous sclerosis complex
TSC1	Tuberous sclerosis complex 1
TSC2	Tuberous sclerosis complex 2
<i>UBE3A</i>	<i>Ubiquitin protein ligase E3A</i>
WES	Whole exome sequencing
WGS	Whole genome sequencing
<i>YWHAG</i>	<i>Tyrosine 3-monooxygenase/tryptophan 5-monooxygenase activation protein gamma</i>

Policy Guidelines

Epilepsy, defined as having two or more unprovoked seizures, is a common neurologic disorder, affecting an estimated 3 million adults and 470,000 children in the United States (CDC, 2020). The biological and genetic mechanisms that disturb the balance between excitatory and inhibitory neuronal circuits to result in epilepsy are extremely heterogeneous (Ottman et al., 2010; Williams & Battaglia, 2013). Approximately 20–30% of epilepsy diagnoses can be attributed to other primary conditions, such as stroke, tumor or head injury, but the remaining 70–80% of cases are believed to be due to one or more genetic factors (Hildebrand et al., 2013; Myers & Mefford, 2015).

The epilepsies can be classified by multiple approaches. Clinically, they can be broadly grouped into three classes: genetic generalized epilepsy (GGE), focal epilepsy, and epileptic encephalopathy (EE). GGE is characterized by generalized seizures that involve both sides of

Genetic Testing for Epilepsy AHS – M2075

the brain, start in childhood or adolescence, are usually associated with normal development and intellect, and include juvenile myoclonic epilepsy and childhood absence epilepsy, among others. Focal seizures originate in one hemisphere of the brain and include temporal lobe epilepsy and autosomal dominant nocturnal frontal lobe epilepsy. EE are severe, early onset conditions characterized by refractory seizures, developmental delay or regression associated with ongoing epileptic activity, and generally poor prognosis such as Dravet, Ohtahara, and West syndromes. Epilepsy is often a comorbid condition with intellectual disability (ID), autism, or schizophrenia, and may be a feature of many metabolic conditions and genetic syndromes (Myers & Mefford, 2015).

The International League Against Epilepsy (ILAE) classifies epilepsies at three levels: seizure types, epilepsy types, and epilepsy syndromes. The classification starts with seizure type as the least specific, epilepsy type as more specific, and epilepsy syndromes as the most specific classification. Seizure type is based on onset, of which there are three categories: focal, generalized, and unknown. The second category, epilepsy type, is divided into four categories; focal, generalized, combined generalized & focal, and unknown. This level of classification typically requires electroencephalography (EEG) data. The “combined generalized & focal” group is new to the 2017 edition of the ILAE classification, and it was created to include patients with both types of seizures, such as patients with Dravet Syndrome. Finally, the “Unknown” category refers to a case where there is not enough information to determine the epilepsy type. This can occur for numerous reasons, such as the unavailability of an EEG. The third level of classification is “epilepsy syndromes.” A syndrome refers to a cluster of features incorporating items such as seizure types, EEG results, and imaging results that tend to occur together. Other distinctive features, such as intellectual disability, may be a part of a syndrome (Scheffer et al., 2017).

Epilepsy is genetically heterogeneous, and extensive phenotypic heterogeneity has been observed even in many monogenic epilepsies. This heterogeneity makes testing each gene individually through Sanger sequencing impractical despite its status as the gold standard (Lemke et al., 2012). Mutations in ion channels, chromatin remodeling, transcriptional regulation, and regulation of the mammalian target of rapamycin (mTOR) protein have been implicated in the etiology of epilepsy (Moller et al., 2016).

Clinical Utility and Validity

The DNA sequencing methods for epilepsy diagnosis have high analytic validity; the methods are well-established in laboratories. Sequencing bi-directionally has been evaluated at over 98% sensitivity (Stenhouse et al., 2013). Next generation sequencing (NGS) allows for parallel sequencing of any number of genes, thereby allowing for far superior testing of heterogenous conditions such as epilepsy. Since the number of new epilepsy genes continues to grow and the phenotypes associated with mutations in each gene are so variable, NGS is a desirable option to diagnose epilepsy patients (Mefford, 2015).

New test options for epilepsy diagnosis have an increased yield of molecular diagnosis, particularly in patients with severe, early-onset epilepsies. For example, a study by Lemke et al. (2012) focused on targeted resequencing of 265 candidate genes, which identified mutations that were presumed to be pathogenic in 16 of 33 patients. Many of these patients had severe epilepsies associated with intellectual disability. This NGS method detected mutations that had been missed by the previous gold standard of Sanger sequencing (with at least one mutation missed due to artifacts from Sanger sequencing). Furthermore, the authors concluded that a patient with more than one possible gene responsible for their phenotype would benefit from this sequencing method compared to Sanger. The authors noted that NGS is only suited for conditions with at least one major gene effect and that whole exome sequencing (WES) would be a better option for more complex genetic conditions (Lemke et al., 2012).

Genetic Testing for Epilepsy AHS – M2075

In 2016, a gene panel targeting 46 epilepsy genes was used on 216 patients representing a wide spectrum of epilepsies with age of onset spanning from the neonatal period to adulthood. A presumed pathogenic variant was identified in 49 (23%) patients. The variants were found in 19 different genes including SCN1A, STXBP1, and CDKL5. Patients with neonatal-onset epilepsies had the highest rate of positive findings (57%). The overall yield for patients with EEs was 32%, compared to 17% among patients with generalized epilepsies and 16% in patients with focal or multifocal (Moller et al., 2016). Another study focusing on the diagnostic use of microarray was performed by Hrabik et al. (2015). Approximately 17.7% (26/147) of participants between the ages of birth to 23 years who were diagnosed with epilepsy and had a SNP microarray performed had an abnormal microarray as defined by laboratory guidelines (Hrabik et al., 2015). Gene panels and exome sequencing for clinical diagnostics provide better and less expensive options for testing and could be implemented early in the diagnostic process. Chromosome microarrays may be considered in severe cases and in GGE with co-morbid features (Myers & Mefford, 2015). Overall, sequencing of many genes at once may provide major benefits to those with epilepsy as epilepsy's genetic profile is widespread.

Copy number abnormalities also play an important role in patients with epilepsy. Of 973 patients who had chromosomal microarray (CMA) and ICD-9 codes for epilepsy or seizures, 805 patients satisfied criteria for epilepsy. 437 copy number variants (CNVs) in 323 patients (1-4 per patient), including 185 (42%) deletions and 252 (58%) duplications were observed. Forty (9%) were confirmed de novo, 186 (43%) were inherited, and parental data were unavailable for 211 (48%). Because the diagnostic yield of CMA for epilepsy patients is similar to the yield in autism spectrum disorders (ASD) and in prenatal diagnosis (which published guidelines recommending testing with CMA), the authors recommended implementation of CMA in the evaluation of unexplained epilepsy (Olson et al., 2014).

Berg et al. (2017) conducted a prospective cohort study of 775 children with newly diagnosed epilepsy with an onset at less than three years of age to determine the role of genetic testing in the initial evaluation of early life epilepsies. Ninety-five children had brain injury, and of the other 680, 327 underwent genetic testing, such as microarrays, WES, and gene panels. In total, 132 (40.4%) children were found to have pathogenic variants. A total of 446 children were deemed to have an unknown etiology without genetic testing, and 180 of these children were tested. Pathogenic variants were found in 48 or 26.7% of the children, however, epilepsy gene-sequencing panels and whole-exome sequencing (WES) had substantially greater diagnostic yields than CMA. Epilepsy panels detected 28 of 96 (29.2%), WES detected 5 of 18 or (27.8%), and CMA detected 8 for 101 or (7.9%). Without a clinically identified cause, testing yields were greater than 15% and as high as 47% depending on patient subgroups. The authors concluded that broad genetic sequencing methods have high diagnostic yields in diagnosing early-life epilepsies regardless of clinical features. Sequencing tests should be incorporated into the initial evaluation of newly presenting early-life epilepsies and not just reserved for those with severe presentations (Berg et al., 2017). Gene panels including the most commonly mutated genes have a 10%-50% diagnostic rate (depending on the panel used and the population) although it is possible that a greater number of genes in the panel does not necessarily equate to a higher diagnostic rate. A 67 gene panel and a 265 gene panel were found to have nearly equivalent diagnostic rates; however, the study using the 67 gene panel only included nineteen patients (Dunn et al., 2018; Mefford, 2015). Exome sequencing has been evaluated at a 25% diagnostic rate (without a prior diagnosis), and whole-genome sequencing (WGS) has been evaluated at as high as 60% with smaller studies and known phenotypes. The weaker sequencing depth contributes to the difficulty of locating copy number variants, which are a significant cause of epilepsy-related conditions. Nonetheless, the authors conclude that WES and WGS will become part of the regularly used tools for clinicians once their cost decreases (Dunn et al., 2018).

Stosser et al. (2017) conducted a retrospective analysis of “893 probands with epilepsy who had an epilepsy panel or WES performed and were positive for a pathogenic or likely

Genetic Testing for Epilepsy AHS – M2075

pathogenic variant in one of nine genes (CDKL5, GABRA1, GABRG2, GRIN2B, KCNQ2, MECP2, PCDH19, SCN1A, or SCN2A),” which found “mosaic pathogenic variants... at an overall frequency of 3.5% (95% CI, 2.4%-4.9%) in nine genes associated with epilepsy-related disorders.... Mosaicism was most common in the CDKL5, PCDH19, SCN2A, and SCN1A genes. Mosaicism was observed in GABRA1, GABRG2, and GRIN2B, which were not reported to have mosaicism prior to this study, and in KCNQ2 and MECP2. Parental mosaicism was observed for pathogenic variants in multiple genes including KCNQ2, MECP2, SCN1A, and SCN2A.” The authors concluded that patients with epilepsy who previously tested negative for pathogenic variants may benefit from an NGS test, which is superior at detecting mosaic variants. The authors also noted that targeted testing of parents of probands may use NGS to better assess risk and mosaicism (Stosser et al., 2017).

Some conditions have well-known mutations caused by a singular gene. For example, in Dravet syndrome, (DS, previously known as severe myoclonic epilepsy of infancy or SMEI) the occurrence of SCN1A mutations is 70%-80%. About 90% of these mutations are de novo. Truncating mutations have found to be more severe (earlier onset of symptoms, faster cognitive decline) compared to missense mutations. PCDH19-related epilepsy is a rare syndrome characterized by focal and/or generalized seizures, which are commonly fever-induced and in clusters. Previously referred to as epilepsy female-restricted with mental retardation (EFMR), PCDH19-related epilepsy occurs primarily in females and has an early onset. A mutation in PCDH19 can cause both DS and PCDH19-related epilepsy as PCDH19 is thought to account for 5% of patients with DS (Andrade & Nascimento, 2022). Another example is tuberous sclerosis complex (TSC); as many as 89% of cases have mutations in either the TSC1 or TSC2 gene (Randle, 2022). Other mutations include KCNQ2 with EE, SLC2A1 with glucose transporter deficiency syndrome, PRRT2 with general infantile convulsions, and ALDH7A1 and PNPO with severe early-onset epilepsy (Poduri et al., 2014). Missense variants in SLC32A1 can cause genetic epilepsy with febrile seizures plus (GEFS+) and idiopathic generalizes epilepsy (IGE) (Heron et al., 2021).

“Epilepsy-plus” patients tend to have highest yield for diagnostic testing; that is, patients with other symptoms such as autism in addition to their epilepsy. Still, even the diagnostic yield of these individuals only reached 50% with panel or exome sequencing (Poduri, 2017). The potential clinical utility of genetic testing for these syndromes is in avoiding further diagnostic testing, directing medication management, and assisting in reproductive decision making.

Establishing the genetic basis of epilepsy in a given patient will help in making treatment decisions, genetic counseling, and more (Poduri et al., 2014). For instance, mutations in ALDH7A1 or in the PNPO gene may lead to seizures not treated with typical antiepileptic drugs, but with pyridoxine (Falsaperla & Corsello, 2017). On the other hand, sodium channel agents may be avoided in patients with an SCN1A mutation. Genetic counseling revolving around topics such as recurrence or heritability risk for pregnancies or testing of relatives is also helpful, and proper care must be taken to ensure that the information is clearly explained to the patient and their family (Poduri et al., 2014).

Genetic testing for epilepsy could also be useful in diagnosing ASD. Peng et al. (2021) discussed, in their research on developing a multiplex gene and phenotype network to identify shared genes between epilepsy and autism, that mutations in genes for subunits of ion channels were related to epilepsy, and that general ion channel dysfunctions “are also linked to susceptibility to autism, as well as bipolar disorder, schizophrenia, and other neuropsychiatric disorders.” The researchers “prioritize ANK2, CACNA1AE, CACNA2D3, GRIA2, and DLG4... as candidate epilepsy genes because of their overlap with the epilepsy-focused module 2 of the WES [whole-exome sequencing] network;” these genes originated as having association only with autism prior to this study (Peng et al., 2021). A study with similar intention done on co-occurring epilepsy and ASD in Chinese children found SCN1A and MECP2 gene mutations to be the most common; SCN1A was more associated with

Genetic Testing for Epilepsy AHS – M2075

epilepsy, while MECP2 was more associated with Rett Syndrome. Mutations in SCN2A, CACNA1A, CACNA1H, CACNA1D, and KCNQ2 were also identified, although individual cases for each gene were small in the latter set of genes (Long et al., 2019). Both studies contribute to the understanding of the beneficial utility for genetic testing of epilepsy in the context of other comorbidities, such as ASD.

Sanchez Fernandez et al. (2019) evaluated the cost-effectiveness of genetic testing in patients with epilepsy of unknown etiology. 20 studies were included, with 8 evaluating chromosomal microarray (CMA), 9 evaluating epileptic panels (EP) with deletion/duplication testing, and 6 evaluating whole exome sequencing (WES). The authors found WES to have the highest diagnostic yield at 0.45, followed by EP at 0.23 and then CMA at 0.08. EP was found to be the most cost-effective, at \$15,848 per diagnosis. Although cost-effectiveness of strategies overlapped, the authors found CMA to be consistently less cost-effective than WES and EP (Sanchez Fernandez et al., 2019).

Borlot et al. (2019) analyzed the results of epilepsy gene panels from 64 patients. Up to 185 genes were tested. 14 probands were found to have “pathogenic or likely pathogenic” variants in the following genes: “SCN1A, GABRB3, UBE3A, KANSL1, SLC2A1, KCNQ2, SLC6A1, HNRNPU, STX1B, SCN2A, PURA, and CHD2”. The authors also identified 6 mutations arising de novo, with unknown inheritance for 8 mutations. Overall, the authors concluded that a commercial gene panel for epilepsy may be useful, as it detected etiology in 22% of patients with epilepsy and intellectual disability (Borlot et al., 2019).

Kim et al. (2020) analyzed the clinical utility of whole-exome sequencing (WES) for patients with infantile-onset epilepsy that had tested negative on gene panel tests for epilepsy. The study included 59 patients. Following WES, 55.4% of the participants received genetic confirmation of epilepsy, indicating that WES increased diagnostic yield by 8%. Three epilepsy-causing genes (that were not included on the original gene panel) were identified: YWHAG, KCNC1, and FGF12. The authors conclude that WES could be an important way to reanalyze novel epilepsy-linked genes without updated gene panels (Kim et al., 2021).

Willimsky et al. (2021) studied the use of next generation sequencing (NGS) by comparing the diagnostic yield of small and large gene panels. The authors completed a retrospective study of 190 patients under the age of 18 years who were diagnosed with epilepsy of unknown etiology. Small gene panels were defined as those under 25 kilo-base pair (kb), and large gene panels were defined as those over 25 kb. Diagnostic yield was defined as detection of pathogenic or likely pathogenic variants. The authors found that the diagnostic yield of large panels (29%) was significantly larger than small panels (13%). The authors then analyzed the differences in diagnostic yield in developmental and epileptic encephalography (DEE) and non-DEE. The authors found that the significant increase of diagnostic yield in large panels was only significant for non-DEE patients, and not for DEE patients. The authors conclude that large epilepsy gene panels have significantly higher diagnostic yield for non-DEE patients but note that small gene panels (a maximum of 10 genes) is sufficient for DEE patients (Willimsky et al., 2021).

Stefanski et al. (2021) completed a meta-analysis and systematic review on the success rate of genetic testing of neurodevelopmental disorders (NDD). The study measured the diagnostic yield, defined as the “percent of pathogenic variant carriers identified in a cohort.” The study included 103 clinical sequencing studies that used NGS in a total of 32,331 people with epilepsy, ASD, or ID. The diagnostic yield for epilepsy was 24%. The epilepsy subtypes with the highest diagnostic yield were epilepsy with ID (27.9%) and early onset seizures (36.9%). The authors then studied the diagnostic yield of each sequencing technology and found a diagnostic yield of 27.2% for exome sequencing and a diagnostic yield of 22.6% for target gene sequencing panels (Stefanski et al., 2021). Sheidley et al. (2021) completed a similar meta-analysis and systematic review on 5985 studies on genetic testing of patients with epilepsy. “The overall diagnostic yield across all test modalities was 17%, with the highest

Genetic Testing for Epilepsy AHS – M2075

yield for GS [genome sequencing] (48%), followed by ES [exome sequencing] (24%), MGP [multigene panel] (19%), and [genome-wide comparative genomic hybridization/chromosomal microarray] CGH/CMA (9%).” The authors also studied non-yield outcomes. 24 studies reported on changes in treatment based on genetic testing; “treatment changes were reported in 12-18% of patients with a genetic diagnosis, including avoiding, stopping, or initiating specific antiseizure medications (AMAs) or ketogenic diet (KD) and halting a plan for surgery in the presence of a specific genetic diagnosis” (Sheidley et al., 2021).

Beyond developing panels to identify genetic variants among epileptic patients, recent advancements in the study of epigenetics could potentially be of use with epilepsy symptoms but not a clear-cut genetic basis. In the patients with neurodevelopmental disorders an epilepsy phenotype, 23% of cases had rare differential methylations (Barbosa et al., 2018). “When the parents were able to be tested, ~40% of the methylation variants were de novo, suggesting that methylation abnormalities may be causative in 5-10% of their cohort. When identified, the underlying causes of the methylation changes were varied and included CNVs (copy number variants), sequence variants in regularly elements, or repeat expansions, each of which is easily missed by conventional (even next-generation) sequencing methods” (Hebbar & Mefford, 2020). Genetic and epigenetic testing could further help shape precision medicine in the best types of pharmaceutical drugs used to treat specific epilepsies, such as by understanding individual responses to antiepileptic drug treatment and alterations in drug pharmacokinetics and pharmacodynamics that could guide safer and more effective treatment strategies (Balestrini & Sisodiya, 2018).

McKnight et al. (2022) conducted a study on the diagnostic yield of genetic tests performed on adults with epilepsy. The study included 2,008 individuals and the authors analyzed their NGS targeted testing results of 89-189 genes and clinical information. Diagnostic yield was defined as a definitive molecular diagnosis. In total, 218 adults (10.9%) had clinically actionable findings. The authors compared the diagnostic yield with age of seizure onset and found that “the highest diagnostic yield was in adults with seizure onset during infancy (29.6%, 0–1 year), followed by in early childhood (13.6%, 2–4 years), late childhood (7.0%, 5–10 years), adolescence (2.4%, 11–17 years), and adulthood (3.7%, ≥18 years).” The authors also reported that “comorbid intellectual disability (ID) or developmental delay resulted in a high diagnostic yield (16.0%).” Overall, the authors conclude that genetic testing could impact clinical management and outcomes (McKnight et al., 2022).

Guidelines and Recommendations

International League Against Epilepsy (ILAE)

In 2015, the ILAE Commission of Pediatrics issued a task force report which included the following related to genetic testing in epilepsy.

- “Genetic screening should not be undertaken at a primary or secondary level of care.”
- “Standard care should permit genetic counseling by trained personnel to be undertaken at all levels of care (primary to quaternary).”
- “Genetic evaluation for Dravet syndrome and other infantile-onset epileptic encephalopathies should be available at tertiary and quaternary levels of care (optimal intervention would permit an extended genetic evaluation).”
- “Early diagnosis of some mitochondrial conditions may alter long-term outcome, but whether screening at quaternary level is beneficial is unknown.” (Wilmshurst et al., 2015)

In 2017, the ILAE released a position paper containing a stance on the genetic etiologies of epilepsies. The ILAE states that “the epilepsies in which a genetic etiology has been implicated are quite diverse and, in most cases, the underlying genes are not yet known”. The ILAE also notes that genetic is not equivalent to inherited; de novo mutations are being

Genetic Testing for Epilepsy AHS – M2075

identified with increasing frequency, thereby making an epileptic syndrome not inherited. Furthermore, the ILAE states that environmental contributions to an epileptic syndrome should not be minimized and that a genetic etiology should be of significant effect in causing epilepsy (Scheffer et al., 2017).

In the 2021 report, the ILAE and the International Neuropsychological Society address the lack of an international taxonomy for cognitive disorders in epilepsy and proposes the International Classification of Cognitive Disorders in Epilepsy (IC-CODE), stating that “a taxonomy built on the alignment of genetic, neuroimaging, and cognitive processes promises high precision for the diagnosis of cognitive disorders in epilepsy and their treatment” (Norman et al., 2021).

In 2022, ILAD released a position statement on epilepsy syndromes with onset in neonates and infants. The statement notes that “genetic testing [for Dravet syndrome, previously known as Severe Myoclonic Epilepsy of Infancy] is recommended at all ages, including in adults in whom the diagnosis is suspected but details of history in infancy may be difficult to access. A pathogenic variant in SCN1A is present in more than 80%–85% of cases.” The statement did not mention recommendations on genetic testing for any other form of epilepsy (Zuberi et al., 2022).

In 2022, the ILAD released a position statement on idiopathic generalized epilepsy syndromes. Genetic testing is mentioned throughout the statement in relation to epilepsy occurring in individuals with intellectual disabilities. In patients suspected of childhood absence epilepsy or generalized tonic–clonic seizures alone, “investigations, including genetic testing, to exclude other etiologies should be considered” in individuals with intellectual disabilities. All other mentions of genetic testing for the diagnosis of epilepsy note that genetic testing is not currently part of the routine diagnostic evaluation but may be used soon, as more genetic determinants are identified (Hirsch et al., 2022).

European Federation of Neurological Societies

In 2010, EFNS issued the following recommendations pertaining to epilepsy. The EFNS noted that molecular investigations may be useful for diagnosis but cannot be considered routine with the large number of mutations present and state that clinical or physiological methods are generally superior. However, they note severe myoclonic epilepsy of infancy (SMEI) has a mutation in the gene SCN1A in 80% of patients. Since 2010, this organization has been renamed to the European Academy of Neurology, but no new updates on epilepsy have been released (Burgunder et al., 2010).

American Academy of Neurology (AAN) and Child Neurology Society (CNS)

The AAN and CNS issued guidelines for clinicians on diagnostic assessment of the child with status epilepticus (SE) in 2006, reaffirmed 2019. The recommendations provided guidance for the assessment of laboratory studies, metabolic and genetic studies, electroencephalography, and neuroimaging in children with SE. The expert panel concluded that “there are insufficient data to support or refute whether genetic testing (chromosomal or molecular studies) should be done routinely in children with SE” (Riviello et al., 2006).

American Epilepsy Society Annual Meeting

In 2015, a talk presented at the American Epilepsy Society Annual Meeting focused on reviewing the genetics of epilepsy, which groups were likely to benefit from genetic testing, and how to approach genetic testing in the current evolution of medicine. With regards to who would benefit most, “it would make sense to focus testing at this time on those in whom there is a high likelihood of a diagnostic finding and on those whose refractory epilepsy may be influenced by a precise genetic diagnosis that can guide treatment.” This study concluded that

Genetic Testing for Epilepsy AHS – M2075

genetic testing is needed to determine if treatment regimens will be affected, especially in younger children and infants.

With regards to the proper way to approach genetic testing, “we should start with a clinical approach, defining a specific epilepsy syndrome when relevant, to guide the nature and sequence of genetic testing. Pre- and post- testing counseling, whether provided by a physician or genetic counselor, is critical to maintaining clear communication with patients about the implications and limitations of their genetic testing.” The types of testing include chromosomal microarray analysis (CMA), complete genome sequencing, and exome sequencing (Poduri, 2017).

International Glut1DS Study Group

The International Glut1DS (glucose transporter 1 deficiency syndrome) put forth recommendations on the diagnosis and treatment of Glut1DS in 2020, which involves “characteristic clinical features, definite hypoglycorrhachia, and pathogenic SLC2A1 variants.” Hypoglycorrhachia is defined as a low glucose level in cerebrospinal fluid in the setting normal glucose levels in blood, or normoglycemia. Their strength of support for a combination of the three features is shown below.

Symbol	diagnosis of Glut1DS	start KDT
✓✓✓	confirmed	yes
✓✓	probable	yes
✓	possible	consider
0	negative	not required

		Characteristic clinical features		
		+	-	
Hypoglycorrhachia	+	✓✓✓	✓✓	+
	+	✓✓	✓	-
	-	✓	✓	+
	-	0	0	-
				Pathogenic SLC2A1-variant

On genetic testing, the International Glut1DS group also stated that “the absence of SLC2A1 pathogenic variants does not always exclude Glut1DS...SLC2A1-negative patients can be diagnosed on the basis of hypoglycorrhachia and distinctive clinical features, especially when responsive to KDT [ketogenic diet therapies]” (Klepper et al., 2020).

Genetic Testing for Epilepsy AHS – M2075

North American Consensus Panel on Dravet Syndrome

This panel was convened to establish standards for evaluation of Dravet Syndrome. The panel recommended that genetic testing be performed in patients whose symptoms indicated Dravet syndrome; the panel noted that either specific SCN1A sequencing or a larger gene panel may be performed, and a chromosomal microarray was not necessary.

The panel also notes that genetic testing should be pursued in a child <12 months and with ≥ 2 febrile seizures >15 minutes as well as a child 12-35 months old with at least 1 febrile seizure over 15 minutes. However, genetic testing should not be performed for a child <12 months with only one focal or generalized febrile seizure.

Genetic counseling should be provided, particularly information about heritability and risk of epilepsy in siblings. Other information about SCN1A mutations may also be provided (Wirrell et al., 2017).

Scottish Intercollegiate Guidelines Network (SIGN)

SIGN lists features suggesting genetic generalized epilepsies, which are as follows:

- childhood or teenage onset
- triggered by sleep deprivation and alcohol
- early morning tonic-clonic seizures or myoclonic jerks
- short absence seizures
- photoparoxysmal response on electroencephalography (EEG)
- generalized 3 per second spike and wave or polyspike and wave on EEG.

Genetics services are recommended for patients with a “very strong” family history of epilepsy, or with “a clinical phenotype suggestive of a monogenic epilepsy syndrome” (SIGN, 2018).

National Institute for Health and Care Excellence (NICE)

The NICE clinical guideline on epilepsies, in children, young people, and adults was updated in 2022 to include recommendations on genetic testing.

- Discuss with a neurologist or geneticist any uncertainties about whether to offer genetic testing or which tests to offer to a person with epilepsy.
- Before carrying out genetic tests:
 - discuss the purpose of testing and the possible implications of the results with the person with epilepsy, and their family and carers if appropriate
 - obtain informed consent with appropriate genetic counselling in line with the NHS Genomic Medicine Service.
- Consider whole-genome sequencing for people with epilepsy of unknown cause who:
 - were aged under 2 years when epilepsy started or
 - have clinical features suggestive of a specific genetic epilepsy syndrome (for example, Dravet syndrome) or
 - have additional clinical features such as:
 - a learning disability
 - autism spectrum disorder
 - a structural abnormality (for example, dysmorphism or congenital malformation)
 - unexplained cognitive or memory decline.

Genetic Testing for Epilepsy AHS – M2075

- Consider whole-genome sequencing for people with epilepsy of unknown cause who were aged between 2 and 3 years when epilepsy started, if clinically agreed by a specialist multidisciplinary team (NICE, 2022).

United Kingdom Expert Group on Tuberous Sclerosis Complex

The expert group recommended, with consensus, to offer a genetic test at baseline to patients with definite or probable TSC. The group noted that a genetic test may clarify the diagnosis of TSC for patients that do not fulfill clinical criteria for the condition (Amin et al., 2018).

In 2019, the group released recommendations for provision of coordinated care. They state, “with wider availability of genetic testing, identification of pathogenic mutations in TSC1 or TSC2 is now sufficient to establish a diagnosis, regardless of the presence of clinical features, and is particularly useful in confirming a suspected diagnosis, as many clinical TSC manifestations are infrequent in young patients.” The group further recommends that:

- For patients with newly diagnoses or suspected TSC:
 - “Obtain three-generation family history to assess for additional family members at risk of TSC.
 - Offer genetic testing for family counseling or when TSC diagnosis is in question but cannot be clinically confirmed.”
- For patients already diagnosed with definite or possible TSC:
 - “Offer genetic testing for family counseling or when TSC diagnosis is in question but cannot be clinically confirmed” (Annear et al., 2019)

Genetic Testing Advisory Committee (GTAC) for Ontario, Canada

These guidelines recommend the following as indications for genetic testing:

- “When the clinical features (age of onset, seizure semiology and EEG features) are consistent with a distinct epilepsy syndrome as defined by the International League Against Epilepsy (ILAE), with the exception of syndromes outlined in the following section.”
- “When the prognosis based on clinical and EEG findings is poor or the likelihood of lethal outcome is high.”
- “When epileptic seizures are refractory to medical treatment as defined by the ILAE12 (with no apparent acquired cause).”
- “When epilepsy is associated with features suggestive of inborn errors of metabolism.”
- “When epilepsy is associated with distinctive patterns of malformations of cortical development identified on neuroimaging studies.”
- “When epilepsy is associated with clinical signs of neurodegeneration.”
- “When epilepsy is associated with paroxysmal neurological features such as paroxysmal dyskinesias, episodic ataxias and hemiplegic migraine.”
- “When epilepsy is associated with additional syndromic features such as developmental delay/intellectual disability, multiple congenital anomalies and dysmorphic features.”
- “When familial epilepsy is present, defined as at least two first- degree family members with related epilepsy syndromes, unless the epilepsy syndrome is benign.”

The guidelines also recommend against genetic testing in the following situations:

- “Recognizable seizure syndrome with benign course.”
- “Childhood epilepsy with centro-temporal spikes (previously termed benign rolandic epilepsy).”
- “Isolated mesial temporal lobe epilepsy with hippocampal sclerosis.”

Genetic Testing for Epilepsy AHS – M2075

- “Typical childhood Absence epilepsy (although if it is early-onset or medically refractory epilepsy, one should consider and test for GLUT1 deficiency).”
- “Juvenile myoclonic epilepsy, which is well controlled on medications and without intellectual disability or any signs of neurodegeneration.”
- “Acquired epilepsy.”

The guideline also provides a list of “potentially treatable” genetic/metabolic epilepsies, shown below

Table 3: Potentially treatable genetic/metabolic epilepsies

Effect on ion channels
Dravet syndrome (<i>SCN1A</i>)—avoidance of sodium-channel blockers
<i>SCN8A</i> -related epileptic encephalopathy—carbamazepine, phenytoin [sodium-channel blockers]
<i>KCNQ2</i> -related epileptic encephalopathy—retigabine [potassium-channel openers] or carbamazepine [sodium-channel blockers]
<i>KCNT1</i> -related migrating partial epilepsy of infancy—quinidine [gain-of-function mutations being treated by partial channel antagonist]
<i>GRIN2A</i> -related epileptic encephalopathy—memantine [NMDA (N-methyl-D-aspartate) receptor antagonists]
Alternative energy source
GLUT-1 deficiency syndrome (<i>SLC2A1</i> gene)—ketogenic diet
Modulate epileptogenesis
Tuberous sclerosis complex (<i>TSC1/2</i>)—mTOR inhibitors (Rapamycin/analogs)
Modulating biochemical pathways
Pyridoxine dependency (<i>ALDH7A1</i>)—B6 vitamin
Biotinidase deficiency (<i>BTD</i>)—biotin
Cerebral folate deficiency (<i>FOLR1</i>)—folic acid
Creatine deficiency syndromes (<i>SLC6A8</i> , <i>GATM</i> , <i>GAMT</i>)—creatine, other amino acid supplementation (glycine, arginine, ornithine) or restriction (arginine)
Serine biosynthesis defects (<i>PHGDH</i> , <i>PSAT1</i> , <i>PSPH</i>)—serine, glycine

Finally, the guideline also makes recommendations for panel testing. Some situations in which a gene panel may be considered are if:

- epilepsy phenotyping is broad and does not easily classify the patient into a distinct group.
- “if the clinical diagnosis is clear and genetic heterogeneity is high, but clinical diagnosis is not indicative of a treatable condition.”
- “if clinical diagnosis is unclear and genetic heterogeneity is unknown.”

The guidelines further list examples of these situations, such as seizures with fever as a major trigger and idiopathic generalized epilepsy refractory to treatment (Jain et al., 2019).

-

Finally, the guideline also makes recommendations for panel testing. Some situations in which a gene panel may be considered are if:

- epilepsy phenotyping is broad and does not easily classify the patient into a distinct

Genetic Testing for Epilepsy AHS – M2075

- group.
- “if the clinical diagnosis is clear and genetic heterogeneity is high, but clinical diagnosis is not indicative of a treatable condition.”
- “if clinical diagnosis is unclear and genetic heterogeneity is unknown.”

The guidelines further list examples of these situations, such as seizures with fever as a major trigger and idiopathic generalized epilepsy refractory to treatment (Jain et al., 2019)

Billing/Coding/Physician Documentation Information

This policy may apply to the following codes. Inclusion of a code in this section does not guarantee that it will be reimbursed. For further information on reimbursement guidelines, please see Administrative Policies on the Blue Cross Blue Shield of North Carolina web site at www.bcbsnc.com. They are listed in the Category Search on the Medical Policy search page.

Applicable service codes: 81404, 81405, 81406, 81407, 81419, 81479

BCBSNC may request medical records for determination of medical necessity. When medical records are requested, letters of support and/or explanation are often useful, but are not sufficient documentation unless all specific information needed to make a medical necessity determination is included.

Scientific Background and Reference Sources

Amin, S., Kingswood, J. C., Bolton, P. F., Elmslie, F., Gale, D. P., Harland, C., Johnson, S. R., Parker, A., Sampson, J. R., Smeaton, M., Wright, I., & O’Callaghan, F. J. (2018). The UK guidelines for management and surveillance of Tuberous Sclerosis Complex. *QJM: An International Journal of Medicine*, 112(3), 171-182. <https://doi.org/10.1093/qjmed/hcy215>

Andrade, D., & Nascimento, F. (2022, July 21). Dravet syndrome: Genetics, clinical features, and diagnosis. <https://www.uptodate.com/contents/dravet-syndrome-genetics-clinical-features-and-diagnosis>

Annear, N. M. P., Appleton, R. E., Bassi, Z., Bhatt, R., Bolton, P. F., Crawford, P., Crowe, A., Tossi, M., Elmslie, F., Finlay, E., Gale, D. P., Henderson, A., Jones, E. A., Johnson, S. R., Joss, S., Kerecuk, L., Lipkin, G., Morrison, P. J., O’Callaghan, F. J., . . . Kingswood, J. C. (2019). Tuberous Sclerosis Complex (TSC): Expert Recommendations for Provision of Coordinated Care. *Front Neurol*, 10, 1116. <https://doi.org/10.3389/fneur.2019.01116>

Balestrini, S., & Sisodiya, S. M. (2018). Pharmacogenomics in epilepsy. *Neurosci Lett*, 667, 27-39. <https://doi.org/10.1016/j.neulet.2017.01.014>

Barbosa, M., Joshi, R. S., Garg, P., Martin-Trujillo, A., Patel, N., Jadhav, B., Watson, C. T., Gibson, W., Chetnik, K., Tessereau, C., Mei, H., De Rubeis, S., Reichert, J., Lopes, F., Vissers, L., Kleefstra, T., Grice, D. E., Edelmann, L., Soares, G., . . . Sharp, A. J. (2018). Identification of rare de novo epigenetic variations in congenital disorders. *Nat Commun*, 9(1), 2064. <https://doi.org/10.1038/s41467-018-04540-x>

Berg, A. T., Berkovic, S. F., Brodie, M. J., Buchhalter, J., Cross, J. H., van Emde Boas, W., Engel, J., French, J., Glauser, T. A., Mathern, G. W., Moshe, S. L., Nordli, D., Plouin, P., & Scheffer, I. E. (2010). Revised terminology and concepts for organization of seizures and epilepsies: report of the ILAE Commission on Classification and Terminology, 2005-2009. *Epilepsia*, 51(4), 676-685. <https://doi.org/10.1111/j.1528-1167.2010.02522.x>

Berg, A. T., Coryell, J., Saneto, R. P., Grinspan, Z. M., Alexander, J. J., Kekis, M., Sullivan, J. E., Wirrell, E. C., Shellhaas, R. A., Mytinger, J. R., Gaillard, W. D., Kossoff, E. H.,

Genetic Testing for Epilepsy AHS – M2075

Valencia, I., Knupp, K. G., Wusthoff, C., Keator, C., Dobyns, W. B., Ryan, N., Loddenkemper, T., . . . Koh, S. (2017). Early-Life Epilepsies and the Emerging Role of Genetic Testing. *JAMA Pediatr*, 171(9), 863-871. <https://doi.org/10.1001/jamapediatrics.2017.1743>

Borlot, F., de Almeida, B. I., Combe, S. L., Andrade, D. M., Filloux, F. M., & Myers, K. A. (2019). Clinical utility of multigene panel testing in adults with epilepsy and intellectual disability. *Epilepsia*, 60(8), 1661-1669. <https://doi.org/10.1111/epi.16273>

Burgunder, J. M., Finsterer, J., Szolnoki, Z., Fontaine, B., Baets, J., Van Broeckhoven, C., Di Donato, S., De Jonghe, P., Lynch, T., Mariotti, C., Schöls, L., Spinazzola, A., Tabrizi, S. J., Tallaksen, C., Zeviani, M., Harbo, H. F., & Gasser, T. (2010). EFNS guidelines on the molecular diagnosis of channelopathies, epilepsies, migraine, stroke, and dementias. *European Journal of Neurology*, 17(5), 641-648. <https://doi.org/10.1111/j.1468-1331.2010.02985.x>

CDC. (2020). Epilepsy Data and Statistics. <https://www.cdc.gov/epilepsy/data/index.html>

Dunn, P., Albury, C. L., Maksemous, N., Benton, M. C., Sutherland, H. G., Smith, R. A., Haupt, L. M., & Griffiths, L. R. (2018). Next Generation Sequencing Methods for Diagnosis of Epilepsy Syndromes. *Front Genet*, 9, 20. <https://doi.org/10.3389/fgene.2018.00020>

Falsaperla, R., & Corsello, G. (2017). Pyridoxine dependent epilepsies: new therapeutical point of view. *Ital J Pediatr*, 43(1), 68. <https://doi.org/10.1186/s13052-017-0387-3>

Hebbar, M., & Mefford, H. C. (2020). Recent advances in epilepsy genomics and genetic testing. *F1000Res*, 9. <https://doi.org/10.12688/f1000research.21366.1>

Heron, S. E., Regan, B. M., Harris, R. V., Gardner, A. E., Coleman, M. J., Bennett, M. F., Grinton, B. E., Helbig, K. L., Sperling, M. R., Haut, S., Geller, E. B., Widdess-Walsh, P., Pelekanos, J. T., Bahlo, M., Petrovski, S., Heinzen, E. L., Hildebrand, M. S., Corbett, M. A., Scheffer, I. E., . . . Berkovic, S. F. (2021). Association of SLC32A1 Missense Variants With Genetic Epilepsy With Febrile Seizures Plus. *Neurology*, 96(18), e2251-e2260. <https://doi.org/10.1212/wnl.00000000000011855>

Hildebrand, M. S., Dahl, H. H., Damiano, J. A., Smith, R. J., Scheffer, I. E., & Berkovic, S. F. (2013). Recent advances in the molecular genetics of epilepsy. *J Med Genet*, 50(5), 271-279. <https://doi.org/10.1136/jmedgenet-2012-101448>

Hirsch, E., French, J., Scheffer, I. E., Bogacz, A., Alsaadi, T., Sperling, M. R., Abdulla, F., Zuberi, S. M., Trinkka, E., Specchio, N., Somerville, E., Samia, P., Riney, K., Nabbout, R., Jain, S., Wilmshurst, J. M., Auvin, S., Wiebe, S., Perucca, E., . . . Wirrell, E. C. (2022). ILAE definition of the Idiopathic Generalized Epilepsy Syndromes: Position statement by the ILAE Task Force on Nosology and Definitions. *Epilepsia*, 63(6), 1475-1499. <https://doi.org/https://doi.org/10.1111/epi.17236>

Hrabik, S. A., Standridge, S. M., Greiner, H. M., Neilson, D. E., Pilipenko, V. V., Zimmerman, S. L., Connor, J. A., & Spaeth, C. G. (2015). The Clinical Utility of a Single-Nucleotide Polymorphism Microarray in Patients With Epilepsy at a Tertiary Medical Center. *J Child Neurol*, 30(13), 1770-1777. <https://doi.org/10.1177/0883073815579972>

Jain, P., Andrade, D., Donner, E., Dymont, D., Prasad, A. N., Goobie, S., Boycott, K., Lines, M., & Snead, O. C. (2019). Development of Criteria for Epilepsy Genetic Testing in Ontario, Canada. *Can J Neurol Sci*, 46(1), 7-13. <https://doi.org/10.1017/cjn.2018.341>

Genetic Testing for Epilepsy AHS – M2075

Kim, S. Y., Jang, S. S., Kim, H., Hwang, H., Choi, J. E., Chae, J.-H., Kim, K. J., & Lim, B. C. (2021). Genetic diagnosis of infantile-onset epilepsy in the clinic: Application of whole-exome sequencing following epilepsy gene panel testing. *Clinical Genetics*, 99(3), 418-424. <https://doi.org/https://doi.org/10.1111/cge.13903>

Klepper, J., Akman, C., Armeno, M., Auvin, S., Cervenka, M., Cross, H. J., De Giorgis, V., Della Marina, A., Engelstad, K., Heussinger, N., Kossoff, E. H., Leen, W. G., Leiendecker, B., Monani, U. R., Oguni, H., Neal, E., Pascual, J. M., Pearson, T. S., Pons, R., . . . De Vivo, D. C. (2020). Glut1 Deficiency Syndrome (Glut1DS): State of the art in 2020 and recommendations of the international Glut1DS study group. *Epilepsia Open*, 5(3), 354-365. <https://doi.org/10.1002/epi4.12414>

Lemke, J. R., Riesch, E., Scheurenbrand, T., Schubach, M., Wilhelm, C., Steiner, I., Hansen, J., Courage, C., Gallati, S., Burki, S., Strozzi, S., Simonetti, B. G., Grunt, S., Steinlin, M., Alber, M., Wolff, M., Klopstock, T., Prott, E. C., Lorenz, R., . . . Biskup, S. (2012). Targeted next generation sequencing as a diagnostic tool in epileptic disorders. *Epilepsia*, 53(8), 1387-1398. <https://doi.org/10.1111/j.1528-1167.2012.03516.x>

Long, S., Zhou, H., Li, S., Wang, T., Ma, Y., Li, C., Zhou, Y., Zhou, S., Wu, B., & Wang, Y. (2019). The Clinical and Genetic Features of Co-occurring Epilepsy and Autism Spectrum Disorder in Chinese Children. *Front Neurol*, 10, 505. <https://doi.org/10.3389/fneur.2019.00505>

McKnight, D., Bristow, S. L., Truty, R. M., Morales, A., Stetler, M., Westbrook, M. J., Robinson, K., Riethmaier, D., Borlot, F., & Kellogg, M. (2022). Multigene panel testing in a large cohort of adults with epilepsy: diagnostic yield and clinically actionable genetic findings. *Neurology Genetics*, 8(1).

Mefford, H. C. (2015). Clinical Genetic Testing in Epilepsy. *Epilepsy Curr*, 15(4), 197-201. <https://doi.org/10.5698/1535-7511-15.4.197>

Moller, R. S., Larsen, L. H., Johannesen, K. M., Talvik, I., Talvik, T., Vaher, U., Miranda, M. J., Farooq, M., Nielsen, J. E., Svendsen, L. L., Kjelgaard, D. B., Linnet, K. M., Hao, Q., Uldall, P., Frangu, M., Tommerup, N., Baig, S. M., Abdullah, U., Born, A. P., . . . Dahl, H. A. (2016). Gene Panel Testing in Epileptic Encephalopathies and Familial Epilepsies. *Mol Syndromol*, 7(4), 210-219. <https://doi.org/10.1159/000448369>

Myers, C. T., & Mefford, H. C. (2015). Advancing epilepsy genetics in the genomic era. *Genome Med*, 7, 91. <https://doi.org/10.1186/s13073-015-0214-7>

NICE. (2022). Epilepsies in children, young people and adults. <https://www.nice.org.uk/guidance/ng217/chapter/1-Diagnosis-and-assessment-of-epilepsy#genetic-testing>

Norman, M., Wilson, S. J., Baxendale, S., Barr, W., Block, C., Busch, R. M., Fernandez, A., Hessen, E., Loring, D. W., McDonald, C. R., & Hermann, B. P. (2021). Addressing neuropsychological diagnostics in adults with epilepsy: Introducing the International Classification of Cognitive Disorders in Epilepsy: The IC CODE Initiative. *Epilepsia Open*, 6(2), 266-275. <https://doi.org/10.1002/epi4.12478>

Olson, H., Shen, Y., Avallone, J., Sheidley, B. R., Pinsky, R., Bergin, A. M., Berry, G. T., Duffy, F. H., Eksioglu, Y., Harris, D. J., Hisama, F. M., Ho, E., Irons, M., Jacobsen, C. M., James, P., Kothare, S., Khwaja, O., Lipton, J., Loddenkemper, T., . . . Poduri, A. (2014). Copy number variation plays an important role in clinical epilepsy. *Ann Neurol*, 75(6), 943-958. <https://doi.org/10.1002/ana.24178>

Genetic Testing for Epilepsy AHS – M2075

Ottman, R., Hirose, S., Jain, S., Lerche, H., Lopes-Cendes, I., Noebels, J. L., Serratosa, J., Zara, F., & Scheffer, I. E. (2010). Genetic testing in the epilepsies--report of the ILAE Genetics Commission. *Epilepsia*, 51(4), 655-670. <https://doi.org/10.1111/j.1528-1167.2009.02429.x>

Peng, J., Zhou, Y., & Wang, K. (2021). Multiplex gene and phenotype network to characterize shared genetic pathways of epilepsy and autism. *Sci Rep*, 11(1), 952. <https://doi.org/10.1038/s41598-020-78654-y>

Poduri, A. (2017). When Should Genetic Testing Be Performed in Epilepsy Patients? *Epilepsy Curr*, 17(1), 16-22. <https://doi.org/10.5698/1535-7511-17.1.16>

Poduri, A., Sheidley, B. R., Shostak, S., & Ottman, R. (2014). Genetic testing in the epilepsies—developments and dilemmas. *Nat Rev Neurol*, 10(5), 293-299. <https://doi.org/10.1038/nrneurol.2014.60>

Randle, S. (2022, June 5). Tuberous sclerosis complex: Genetics, clinical features, and diagnosis. <https://www.uptodate.com/contents/tuberous-sclerosis-complex-genetics-clinical-features-and-diagnosis>

Riviello, J. J., Ashwal, S., Hirtz, D., Glauser, T., Ballaban-Gil, K., Kelley, K., Morton, L. D., Phillips, S., Sloan, E., & Shinnar, S. (2006). Practice Parameter: Diagnostic assessment of the child with status epilepticus (an evidence-based review). *Neurology*, 67(9), 1542. <https://doi.org/10.1212/01.wnl.0000243197.05519.3d>

Sanchez Fernandez, I., Loddenkemper, T., Gainza-Lein, M., Sheidley, B. R., & Poduri, A. (2019). Diagnostic yield of genetic tests in epilepsy: A meta-analysis and cost-effectiveness study. *Neurology*. <https://doi.org/10.1212/wnl.0000000000006850>

Scheffer, I. E., Berkovic, S., Capovilla, G., Connolly, M. B., French, J., Guilhoto, L., Hirsch, E., Jain, S., Mathern, G. W., Moshé, S. L., Nordli, D. R., Perucca, E., Tomson, T., Wiebe, S., Zhang, Y.-H., & Zuberi, S. M. (2017). ILAE classification of the epilepsies: Position paper of the ILAE Commission for Classification and Terminology. *Epilepsia*, 58(4), 512-521. <https://doi.org/10.1111/epi.13709>

Sheidley, B. R., Malinowski, J., Bergner, A. L., Bier, L., Gloss, D. S., Mu, W., Mulhern, M. M., Partack, E. J., & Poduri, A. (2021). Genetic testing for the epilepsies: A systematic review. *Epilepsia*. <https://doi.org/10.1111/epi.17141>

SIGN. (2018). Diagnosis and management of epilepsy in adults. https://www.sign.ac.uk/media/1079/sign143_2018.pdf

Stefanski, A., Calle-López, Y., Leu, C., Pérez-Palma, E., Pestana-Knight, E., & Lal, D. (2021). Clinical sequencing yield in epilepsy, autism spectrum disorder, and intellectual disability: A systematic review and meta-analysis. *Epilepsia*, 62(1), 143-151. <https://doi.org/10.1111/epi.16755>

Stenhouse, S. A., Ellis, R., & Zuberi, S. (2013). SCN1A Genetic Test for Dravet Syndrome (Severe Myoclonic Epilepsy of Infancy and its Clinical Subtypes) for use in the Diagnosis, Prognosis, Treatment and Management of Dravet Syndrome. *PLoS Curr*, 5. <https://doi.org/10.1371/currents.eogt.c553b83d745dd79bfb61eaf35e522b0b>

Stosser, M. B., Lindy, A. S., Butler, E., Retterer, K., Piccirillo-Stosser, C. M., Richard, G., & McKnight, D. A. (2017). High frequency of mosaic pathogenic variants in genes causing epilepsy-related neurodevelopmental disorders. *Genet Med*. <https://doi.org/10.1038/gim.2017.114>

Genetic Testing for Epilepsy AHS – M2075

Williams, C. A., & Battaglia, A. (2013). Molecular biology of epilepsy genes. *Exp Neurol*, 244, 51-58. <https://doi.org/10.1016/j.expneurol.2011.12.001>

Willimsky, E. K., Munzig, A., Mayer, K., Biskup, S., Abicht, A., Hoertnagel, K., Voss, H. V., Klein, H. G., Rost, I., Larsen, L. H. G., Dahl, H. A., Hoelz, H., Stuelpnagel, C. V., & Borggraefe, I. (2021). Next Generation Sequencing in Pediatric Epilepsy Using Customized Panels: Size Matters. *Neuropediatrics*, 52(2), 92-97. <https://doi.org/10.1055/s-0040-1712488>

Wilmshurst, J. M., Gaillard, W. D., Vinayan, K. P., Tsuchida, T. N., Plouin, P., Van Bogaert, P., Carrizosa, J., Elia, M., Craiu, D., Jovic, N. J., Nordli, D., Hirtz, D., Wong, V., Glauser, T., Mizrahi, E. M., & Cross, J. H. (2015). Summary of recommendations for the management of infantile seizures: Task Force Report for the ILAE Commission of Pediatrics. *Epilepsia*, 56(8), 1185-1197. <https://doi.org/10.1111/epi.13057>

Wirrell, E. C., Laux, L., Donner, E., Jette, N., Knupp, K., Meskis, M. A., Miller, I., Sullivan, J., Welborn, M., & Berg, A. T. (2017). Optimizing the Diagnosis and Management of Dravet Syndrome: Recommendations From a North American Consensus Panel. *Pediatr Neurol*, 68, 18-34.e13. <https://doi.org/10.1016/j.pediatrneurol.2017.01.025>

Zuberi, S. M., Wirrell, E., Yozawitz, E., Wilmshurst, J. M., Specchio, N., Riney, K., Pressler, R., Auvin, S., Samia, P., Hirsch, E., Galicchio, S., Triki, C., Snead, O. C., Wiebe, S., Cross, J. H., Tinuper, P., Scheffer, I. E., Perucca, E., Moshé, S. L., & Nabbout, R. (2022). ILAE classification and definition of epilepsy syndromes with onset in neonates and infants: Position statement by the ILAE Task Force on Nosology and Definitions. *Epilepsia*, 63(6), 1349-1397. <https://doi.org/10.1111/epi.17239>

Policy Implementation/Update Information

- | | |
|----------|--|
| 1/1/19 | New policy developed. BCBSNC will provide coverage for genetic testing for epilepsy when it is determined to be medically necessary because the medical criteria and guidelines are met. Medical Director review 1/1/2019. Policy noticed 1/1/2019 for effective date 4/1/2019. (sk) |
| 5/14/19 | Reviewed by Avalon 1st Quarter 2019 CAB. Related policies added. Regulatory Status added. Not Covered statement reworded. Intent unchanged. Background section updated. Clinical Utility and Validity section added. Policy Guidelines updated. Medical Director review 4/2019. References added. (sk) |
| 10/29/19 | Wording in the Policy, When Covered, and/or Not Covered section(s) changed from Medical Necessity to Reimbursement language, where needed. |
| 11/26/19 | Specialty Matched Consultant Advisory Panel review 10/16/2019. Coding table removed from Billing/Coding section. (sk) |
| 5/12/20 | Reviewed by Avalon 1st Quarter 2020 CAB. Additional coverage criteria added. Not Covered statement reworded. Added code 81404. Medical Director review 4/2019. References added. (bb) |
| 11/10/20 | Specialty Matched Consultant Advisory Panel review 10/21/2020. (sk) |
| 5/4/21 | Reviewed by Avalon 1st Quarter 2021 CAB. Policy Guidelines updated. Added code 81419. Medical Director review 4/2021. References added. (sk) |

Genetic Testing for Epilepsy AHS – M2075

- 2/8/22 Specialty Matched Consultant Advisory Panel review 10/20/2021. (sk)
- 5/16/23 Reviewed by Avalon 1st Quarter 2023 CAB. Policy Guidelines updated. Medical Director review 4/2023. References added. (sk)

Medical policy is not an authorization, certification, explanation of benefits or a contract. Benefits and eligibility are determined before medical guidelines and payment guidelines are applied. Benefits are determined by the group contract and subscriber certificate that is in effect at the time services are rendered. This document is solely provided for informational purposes only and is based on research of current medical literature and review of common medical practices in the treatment and diagnosis of disease. Medical practices and knowledge are constantly changing and BCBSNC reserves the right to review and revise its medical policies periodically.