

Corporate Medical Policy

Diagnosis and Treatment of Sacroiliac Joint Pain

File Name: diagnosis_and_treatment_of_sacroiliac_joint_pain
Origination: 8/2010
Last Review: 4/2022

Description of Procedure or Service

Sacroiliac joint (SIJ) arthrography using fluoroscopic guidance with injection of an anesthetic has been explored as a diagnostic test for sacroiliac joint pain. Duplication of the patient's pain pattern with the injection of contrast medium suggests a sacroiliac etiology, as does relief of chronic back pain with injection of local anesthetic. Treatment of sacroiliac joint pain with corticosteroids, radiofrequency ablation (RFA), stabilization, or minimally invasive sacroiliac joint fusion has also been explored.

Similar to other structures in the spine, it is assumed that the sacroiliac joint may be a source of low back pain. In fact, prior to 1928, the sacroiliac joint was thought to be the most common cause of sciatica. In 1928, the role of the intervertebral disc was elucidated, and from that point forward the sacroiliac joint received less research attention.

Research into sacroiliac joint pain has been plagued by lack of a criterion standard to measure its prevalence and against which various clinical examinations can be validated. For example, sacroiliac joint pain is typically without any consistent, demonstrable radiographic or laboratory features and most commonly exists in the setting of morphologically normal joints. Clinical tests for sacroiliac joint pain may include various movement tests, palpation to detect tenderness, and pain descriptions by the patient. Further confounding the study of the sacroiliac joint is that multiple structures, such as posterior facet joints and lumbar discs, may refer pain to the area surrounding the sacroiliac joint.

Because of inconsistent information obtained from history and physical examination, some have proposed the use of image-guided anesthetic injection into the sacroiliac joint for the diagnosis of sacroiliac joint pain. Treatments being investigated for sacroiliac joint pain include prolotherapy, corticosteroid injection, radiofrequency ablation, stabilization, and arthrodesis. Some procedures have been referred to as SIJ fusion but may be more appropriately called fixation due to little to no bridging bone on radiographs. Devices for SIJ fixation/fusion that promote bone ingrowth to fixate the implants include a triangular implant (iFuse Implant System) and cylindrical threaded devices (Rialto, SImmetry, Silex, SambaScrew, SI-LOK). Some devices also have a slot in the middle where autologous or allogeneic bone can be inserted. This added bone is intended to promote the fusion of the SIJ.

This policy does not address the treatment of sacroiliac joint pain due to infection, trauma or neoplasm.

Regulatory Status

A number of radiofrequency generators and probes have been cleared for marketing by the U.S. Food and Drug Administration (FDA) through the 510(k) process. In 2005, the SInergy® (Halyard; formerly Kimberly-Clark), a water-cooled single-use probe, was cleared by the FDA, listing the Baylis Pain Management Probe as a predicate device. The intended use is in

Diagnosis and Treatment of Sacroiliac Joint Pain

conjunction with a radiofrequency generator to create radiofrequency lesions in nervous tissue.
FDA product code: GXD, GXI.

A number of percutaneous or minimally invasive fixation/fusion devices have been cleared for marketing by FDA through the 510(k) process. They include the iFuse® Implant System and iFuse-3D™ Implant (SI Bone), the Rialto™ SI Joint Fusion System (Medtronic), SIJ-Fuse (Spine Frontier), the SImmetry® Sacroiliac Joint Fusion System (Zyga Technologies), Silex™ Sacroiliac Joint Fusion System (XTANT Medical), SambaScrew® and FIREBIRD SI Fusion System (Orthofix), and the SI-LOK® Sacroiliac Joint Fixation System (Globus Medical) and the SIImpact Sacroiliac Joint Fixation System (Life Spine). FDA product code: OUR.

Related Policies

Facet Joint Denervation

Sacroiliac Joint Fusion

Vertebroplasty, Kyphoplasty, and Sacroplasty Percutaneous Prolotherapy

*****Note: This Medical Policy is complex and technical. For questions concerning the technical language and/or specific clinical indications for its use, please consult your physician.**

Policy

BCBSNC will provide coverage for diagnosis and treatment of sacroiliac joint pain when it is determined to be medically necessary because the medical criteria and guidelines noted below are met.

Arthrography and radiofrequency denervation of the sacroiliac joint are considered investigational for all applications. BCBSNC does not provide coverage for investigational services or procedures.

Benefits Application

This medical policy relates only to the services or supplies described herein. Please refer to the Member's Benefit Booklet for availability of benefits. Member's benefits may vary according to benefit design; therefore member benefit language should be reviewed before applying the terms of this medical policy.

When diagnosis and treatment of sacroiliac joint pain is covered

Injection of anesthetic for diagnosing sacroiliac joint pain may be considered medically necessary when the following criteria have been met:

- Pain has failed to respond to 3 months of conservative management, which may consist of therapies such as nonsteroidal anti-inflammatory medications, acetaminophen, manipulation, physical therapy, and a home exercise program; AND
- Dual (controlled) diagnostic blocks with 2 anesthetic agents with differing duration of action are used; AND
- The injections are performed under imaging guidance

Injection of corticosteroid may be considered medically necessary for the treatment of sacroiliac joint pain when the following criteria have been met:

- Pain has failed to respond to 3 months of conservative management, which may consist of therapies such as nonsteroidal anti-inflammatory medications, acetaminophen, manipulation, physical therapy, and a home exercise program; AND
- The injection is performed under imaging guidance; AND
- No more than 3 injections are given in one year

Diagnosis and Treatment of Sacroiliac Joint Pain

When diagnosis and treatment of sacroiliac joint pain is not covered

Arthrography of the sacroiliac joint is considered investigational.

Radiofrequency denervation of the sacroiliac joint is considered investigational.

Policy Guidelines

Summary of Evidence

Diagnostic

For individuals who have suspected SIJ pain who receive a diagnostic sacroiliac block, the evidence includes systematic reviews. Relevant outcomes are test validity, symptoms, functional outcomes, quality of life, medication use, and treatment-related morbidity. Current evidence is conflicting on the diagnostic utility of SIJ blocks. Based on the established use of injections to diagnose pain in other joints, the evidence is sufficient to determine the effects of the technology on health outcomes for controlled diagnostic (two blocks with anesthetics of different duration) injections.

Therapeutic

For individuals who have SIJ pain who receive therapeutic corticosteroid injections, the evidence includes small RCTs and case series. Relevant outcomes are symptoms, functional outcomes, quality of life, medication use, and treatment-related morbidity. In general, the literature on injection therapy of joints in the back is of poor quality. Based on the established use of injections to treat pain in other joints, the evidence is sufficient to determine the effects of the technology on health outcomes for therapeutic (corticosteroid) injections.

For individuals who have SIJ pain who receive RFA, the evidence includes 5 RCTs using different radiofrequency applications and case series. Relevant outcomes are symptoms, functional outcomes, quality of life, medication use, and treatment-related morbidity. Meta-analysis of available sham-controlled RCTs suggests that there may be a small effect of RFA on SIJ pain at short-term (1-3 months) follow-up. However, the randomized trials of RFA have methodologic limitations, and there is limited data on the duration of the treatment effect. The single RCT with 6 and 12 month follow-up showed no significant benefit of RFA compared to an exercise control group at these time points. In addition, heterogeneity of RFA treatment techniques precludes generalizing results across different studies. For RFA with a cooled probe, the 2 small RCTs reported short-term benefits, but these are insufficient to determine the overall effect on health outcomes. The RCT on palisade RFA of the SIJ did not include a sham control. Another sham-controlled randomized trial showed no benefit of RFA. Further high-quality controlled trials are needed to compare this procedure in defined populations with sham control and alternative treatments. The evidence is insufficient to determine the effects of the technology on health outcomes.

The American Society of Interventional Pain Physicians (2013) guidelines have been updated. The updated guidelines recommend the use of controlled SIJ blocks with placebo or controlled comparative local anesthetic block when indications are satisfied with suspicion of SIJ pain. A positive response to a joint block is considered to be at least a 75% improvement in pain or in the ability to perform previously painful movements.

Billing/Coding/Physician Documentation Information

This policy may apply to the following codes. Inclusion of a code in this section does not guarantee that it will be reimbursed. For further information on reimbursement guidelines, please see Administrative Policies on the Blue Cross Blue Shield of North Carolina web site at www.bcsnc.com. They are listed in the Category Search on the Medical Policy search page.

Diagnosis and Treatment of Sacroiliac Joint Pain

Applicable codes: 27096, 27279, G0259, G0260, 64625

BCBSNC may request medical records for determination of medical necessity. When medical records are requested, letters of support and/or explanation are often useful, but are not sufficient documentation unless all specific information needed to make a medical necessity determination is included.

Scientific Background and Reference Sources

For guideline titled “Sacroiliac Joint Arthroscopy and Injection

BCBSA Medical Policy Reference Manual [Electronic Version]. 6.01.23, 2/1/10

Boswell MV, Trescot AM, Sukdeb D, et al. Interventional techniques: evidence-based practice guidelines in the management of chronic spinal pain. *Pain Physician* 2007; 10:7-111. Retrieved on July 20, 2010 from <http://www.ncbi.nlm.nih.gov/pubmed/17256025>

Manchikanti L, Boswell MV, Singh V et al. Comprehensive evidence-based guidelines for interventional techniques in the management of chronic spinal pain. *Pain Physician* 2009; 12(4):699-802. Retrieved on July 20, 2010 from <http://www.ncbi.nlm.nih.gov/pubmed/19644537>

Manchikanti L, Datta S, Derby R, et al. A critical review of the American Pain Society clinical practice guidelines for interventional techniques: part 1. Diagnostic interventions. *Pain Physician*. 2010 May-Jun; 13(3):E141-74. Retrieved on July 20, 2010 from http://www.painphysicianjournal.com/linkout_vw.php?issn=1533-3159&vol=13&page=E141

Rupert MP, Lee M, Manchikanti L et al. Evaluation of sacroiliac joint interventions: a systematic appraisal of the literature. *Pain Physician* 2009; 12(2):399-418. Retrieved on July 20, 2010 from <http://www.ncbi.nlm.nih.gov/pubmed/19305487>

Senior Medical Director review 8/2010

Manchikanti L, Datta S, Gupta S et al. A critical review of the American Pain Society Clinical practice guidelines for interventional techniques: part 2. Therapeutic interventions. *Pain Physician*. 2010; 13(4):E215-64.

BCBSA Medical Policy Reference Manual [Electronic Version]. 6.01.23, 2/10/11

Specialty Matched Consultant Advisory Panel review 7/2011

For guideline titled “Diagnosis and Treatment of Sacroiliac Joint Pain”

BCBSA Medical Policy Reference Manual [Electronic Version]. 6.01.23, 2/9/12

Specialty Matched Consultant Advisory Panel review 7/2012

BCBSA Medical Policy Reference Manual [Electronic Version]. 6.01.23, 2/14/13

Specialty Matched Consultant Advisory Panel – 5/2013

BCBSA Medical Policy Reference Manual [Electronic Version]. 6.01.23, 5/22/14

Specialty Matched Consultant Advisory Panel – 5/2014

For policy titled “Diagnosis and Treatment of Sacroiliac Joint Pain”

Diagnosis and Treatment of Sacroiliac Joint Pain

BCBSA Medical Policy Reference Manual [Electronic Version]. 6.01.23, 4/23/15

Specialty Matched Consultant Advisory Panel – 5/2015

BCBSA Medical Policy Reference Manual [Electronic Version]. 6.01.23, 11/12/15

Specialty Matched Consultant Advisory Panel – 5/2016

BCBSA Medical Policy Reference Manual [Electronic Version]. 6.01.23, 8/11/16

BCBSA Medical Policy Reference Manual [Electronic Version]. 6.01.23, 10/13/16

BCBSA Medical Policy Reference Manual [Electronic Version]. 6.01.23, 12/14/2017

BCBSA Medical Policy Reference Manual [Electronic Version]. 6.01.23, 11/8/2018

Specialty Matched Consultant Advisory Panel – 04/2020

BCBSA Medical Policy Reference Manual [Electronic Version]. 6.01.23, 11/14/19

Manchikanti L, Abdi S, Atluri S, et al. An update of comprehensive evidence-based guidelines for interventional techniques in chronic spinal pain. Part II: guidance and recommendations. *Pain Physician*. Apr 2013;16(2 Suppl): S49-283. PMID 23615883

BCBSA Medical Policy Reference Manual [Electronic Version]. 6.01.23, 11/12/2020

Specialty Matched Consultant Advisory Panel – 04/2021

Medical Director review 4/2021

Specialty Matched Consultant Advisory Panel – 04/2022

Medical Director review 4/2022

Policy Implementation/Update Information

For guideline titled “Sacroiliac Joint Arthroscopy and Injection

8/31/10 New Evidence Based Guideline implemented. Sacroiliac joint arthrography and/or injection are not recommended as treatment for sacroiliac pain. (mco)

4/26/11 References updated. No changes to guideline statements. (mco)

8/16/11 Specialty Matched Consultant Advisory Panel review 7/2011. No changes to guideline statements. (mco)

12/30/11 Deleted code 73542 from “Billing/Coding” section. (mco)

For guideline titled “Diagnosis and Treatment of Sacroiliac Joint Pain”

5/1/12 Guideline titled changed from “Sacroiliac Joint Arthroscopy and Injection” to “Diagnosis and Treatment of Sacroiliac Joint Pain.” Description section updated. “Not Recommended” section updated. The following statement added to the Evidence Based

Diagnosis and Treatment of Sacroiliac Joint Pain

- Guidelines: “Radiofrequency ablation of the sacroiliac joint is not recommended as a treatment for sacroiliac pain.” References updated. Medical Director review 4/2012. (mco)
- 5/15/12 Information regarding radiofrequency ablation of the sacroiliac joint deleted. Description section updated to include reference for BCBSNC policy titled, “Facet Joint Denervation.” Medical Director review 5/2012. (mco)
- 8/7/12 Specialty Matched Consultant Advisory Panel review 7/2012. No changes to guideline statements. (mco)
- 4/16/13 References updated. Added “Sacral Joint Fusion” as a related policy. (mco)
- 7/16/13 Specialty Matched Consultant Advisory Panel review 5/15/2013. No change to guideline. (btw)
- 1/28/14 Added HCPCS codes G0259 and G0260 to Billing/Coding section. (btw)
- 11/25/14 Reference added. Specialty Matched Consultant Advisory Panel review 5/27/2014. Guideline changed to may be appropriate for controlled diagnostic injections and for therapeutic injections with corticosteroid. Medical Director review. (sk)

For policy titled “Diagnosis and Treatment of Sacroiliac Joint Pain”

- 7/28/15 Evidence based guideline converted to corporate medical policy. Medical director review. Reference added. Specialty Matched Consultant Advisory Panel review 5/27/2015. Notification given 7/28/15 for policy effective date 10/1/15. (sk)
- 2/29/16 Reference added. Related policy added. Policy Guidelines updated. (sk)
- 7/1/16 Policy Guidelines updated. Specialty Matched Consultant Advisory Panel review 5/25/2016. (sk)
- 11/22/16 References added. Policy Guidelines updated. (sk)
- 6/30/17 Specialty Matched Consultant Advisory Panel review 4/26/2017. No change to policy statement. (an)
- 6/8/18 Codes 27096 and 27279 added to Billing/Coding section. Reference added. Specialty Matched Consultant Advisory Panel review 5/23/2018. No change to policy statement. (an)
- 4/30/19 Updated Description and Policy Guidelines sections. Deleted code 20552 from Billing/Coding section. Reference added. Specialty Matched Consultant Advisory Panel review 4/17/2019. (an)
- 4/28/20 References added. Policy guidelines updated. No change to policy statement. Specialty Matched Consultant Advisory Panel review 4/15/2020. (eel)
- 5/4/21 Updated Description and Policy Guidelines section. Code 64625 added to Billing/Coding section. Reference added. Specialty Matched Consultant Advisory Panel review 4/2021. Medical Director review 4/2021. (bb)
- 11/16/21 Correction made to Billing/Coding section. Added code 64625. (tt)
- 5/3/22 References added. Related policies updated. No change to policy statement. Specialty Matched Consultant Advisory Panel review 4/2022. (tt)

Diagnosis and Treatment of Sacroiliac Joint Pain

Medical policy is not an authorization, certification, explanation of benefits or a contract. Benefits and eligibility are determined before medical guidelines and payment guidelines are applied. Benefits are determined by the group contract and subscriber certificate that is in effect at the time services are rendered. This document is solely provided for informational purposes only and is based on research of current medical literature and review of common medical practices in the treatment and diagnosis of disease. Medical practices and knowledge are constantly changing and BCBSNC reserves the right to review and revise its medical policies periodically.