

## Corporate Medical Policy

### Diagnosis of Vaginitis including Multi-target PCR Testing AHS – M2057

<b>File Name:</b>	diagnosis_of_vaginitis_including_multi_target_pcr_testing
<b>Origination:</b>	04/2019
<b>Last CAP Review:</b>	03/2021
<b>Next CAP Review:</b>	03/2022
<b>Last Review:</b>	04/2021

#### Description of Procedure or Service

---

##### Definition

Vaginitis is defined as inflammation of the vagina with symptoms of discharge, itching, and discomfort often due to a disruption of the vaginal microflora. The most common infections are bacterial vaginosis, *Candida* vulvovaginitis, and trichomoniasis (Sobel, 1999). Other causes include vaginal atrophy in postmenopausal women, cervicitis, foreign body, irritants, and allergens (Sobel, 2020a).

Bacterial vaginosis (BV) is characterized by a shift in microbial species from the normally dominant hydrogen-peroxide producing *Lactobacillus* species to *Gardnerella vaginalis* and anaerobic commensals (Eschenbach et al., 1989; Hill, 1993; Lamont et al., 2011; Ling et al., 2010; Sobel, 2020b).

Vulvovaginal candidiasis (VVC) is characterized by *Candida* species. It is the second most common cause of vaginitis symptoms (after BV) and accounts for approximately one-third of vaginitis cases (Sobel, 2018c; Workowski & Bolan, 2015).

Trichomoniasis is caused by the flagellated protozoan *Trichomonas vaginalis*, which principally infects the squamous epithelium in the urogenital tract: vagina, urethra, and paraurethral glands (Kissinger, 2015; Sobel, 2020).

##### Related Policies:

Cervical Cancer Screening  
Diagnostic Testing of Sexually Transmitted Infections  
Identification of Microorganisms Using Nucleic Acid Probes  
Pathogen Panel Testing

**\*\*\*Note: This Medical Policy is complex and technical. For questions concerning the technical language and/or specific clinical indications for its use, please consult your physician.**

#### Policy

---

**BCBSNC will provide coverage for diagnosis of vaginitis including multi-target PCR testing when it is determined the medical criteria or reimbursement guidelines below are met.**

#### Benefits Application

---

# Diagnosis of Vaginitis including Multi-target PCR Testing AHS – M2057

This medical policy relates only to the services or supplies described herein. Please refer to the Member's Benefit Booklet for availability of benefits. Member's benefits may vary according to benefit design; therefore member benefit language should be reviewed before applying the terms of this medical policy.

## **When diagnosis of vaginitis including multi-target PCR testing is covered**

---

Reimbursement is allowed for testing of pH, testing for the presence of amines, saline wet mount, hydrogen peroxide (KOH) wet mount and microscopic examination of vaginal fluids in patients with symptoms of vaginitis.

Reimbursement is allowed for direct Probe DNA-based identification of *Gardnerella*, *Trichomonas*, and *Candida* in patients with symptoms of vaginitis.

Reimbursement is allowed for vaginal culture or *Candida* species for the diagnosis of vulvovaginal candidiasis in patients with clinical signs and symptoms of vaginitis and negative findings on wet-mount preparations and a normal pH test.

Reimbursement is allowed for measurement of sialidase activity in vaginal fluid for the diagnosis of bacterial vaginosis in women with symptoms of vaginitis.

Reimbursement is allowed for Nucleic Acid Amplification Test (NAAT) or Polymerase Chain Reaction (PCR)-based identification of *Trichomonas vaginalis* in patients with symptoms of vaginitis.

Reimbursement is allowed for screening for *Trichomonas* for women with risk factors including: new or multiple partners; history of sexually transmitted diseases (STDs); exchange of sex for payment; or injection drug use.

Reimbursement is allowed for rapid identification of *Trichomonas* by enzyme immunoassay in patients with symptoms of vaginitis.

## **When diagnosis of vaginitis including multi-target PCR testing is not covered**

---

Reimbursement is not allowed for Polymerase Chain Reaction (PCR) based identification of *Candida* for any indication.

Reimbursement is not allowed for screening for Trichomoniasis and bacterial vaginosis in asymptomatic patients, including asymptomatic pregnant patients at average or high risk for premature labor.

Reimbursement is not allowed for PCR testing and Multitarget polymerase chain reaction (PCR) testing for diagnosis of bacterial vaginosis.

Reimbursement is not allowed for molecular-based panel testing, including, but not limited to testing such as SmartJane™, to test for microorganisms involved in vaginal flora imbalance and/or infertility.

## **Policy Guidelines**

---

### **Background**

Vaginitis is characterized by several symptoms including odor, itching, abnormal vaginal discharge, burning and irritation; this inflammatory ailment is considered the most common gynecologic diagnosis in primary care as most women experience vaginitis at least once in their

# Diagnosis of Vaginitis including Multi-target PCR Testing AHS – M2057

lives (Paladine & Desai, 2018). A diagnosis of vaginitis can be given based on a combination of symptoms, physical examination, and office or laboratory-based testing methods.

The squamous epithelium of the vagina in premenopausal women is rich in glycogen, a substrate for lactobacilli, which create an acidic vaginal environment (pH 4.0 to 4.5). This acidity helps maintain the normal vaginal flora and inhibits growth of pathogenic organisms. Disruption of the normal ecosystem by menstrual cycle, sexual activity, contraceptive, pregnancy, foreign bodies, estrogen level, sexually transmitted diseases, and use of hygienic products or antibiotics can lead to development of vaginitis. Bacterial vaginosis (BV), vulvovaginal candidiasis (VVC), and trichomoniasis are the three most common infections responsible for vaginitis. Other causes include: vaginal atrophy in postmenopausal women, cervicitis, foreign body, irritants and allergens (Sobel, 2020a).

Bacterial vaginosis is caused by an imbalance of naturally occurring vaginal bacteria, characterized by both a change in the most common type of bacteria present, along with an increase in the total number of bacteria present. Normal vaginal microbiota is dominated by the species *Lactobacilli*, which are known to produce hydrogen peroxide and lactic acid, which help to keep the acidic vaginal environment below pH 4.5 (Jones, 2019; Kairys & Garg, 2020). Though the origin of vaginal bacterial infections is still unclear, it is believed that most of such infections are the result of another bacteria, *Gardnerella vaginalis*, creating a biofilm which allows opportunistic bacteria to grow within the vagina, causing a decrease in the *Lactobacilli* and subsequent disruption of the pH of the system. An entire host of etiologic organisms have been identified as possible instigators and exacerbators, including *Atopobium vaginae*, *Megasphaera* phylotype 1 and 2, *Leptotrichia aminionii*, *Mobiluncus spp*, *Prevotella spp*, *Mycoplasma hominis*, *Bacteroides spp*, *Sneathia*, and BV-associated bacteria (BVAB)1, 2, and 3, though as aforementioned the causative mechanism and the interaction between these species are still uncertain (Jones, 2019).

Laboratory documentation of the etiology of vaginitis is important before initiating therapy, given the nonspecific nature and considerable overlap of the symptoms (Anderson, Klink, & Cochrane, 2004; Ellis, Lerch, & Whitcomb, 2001; Landers, Wiesenfeld, Heine, Krohn, & Hillier, 2004). Diagnostic testing enables targeted treatment, increases therapeutic compliance, and increases the likelihood of partner notification (Sobel, 2020a; Workowski & Bolan, 2015).

Measurement of vaginal pH is the primary initial finding that drives the diagnostic. The pH of the normal vaginal secretions in premenopausal women with relatively high estrogen levels is 4.0 to 4.5. The pH of normal vaginal secretions in premenarchal and postmenopausal women in whom estrogen levels are low is  $\geq 4.7$ . An elevated pH in a premenopausal woman suggests infections, such as BV (pH > 4.5) or trichomoniasis (pH 5 to 6) and helps to exclude *Candida* vulvovaginitis (pH 4 to 4.5). Vaginal pH may also be altered by lubricating gels, semen, douches, intravaginal medications and in pregnant women, leakage of amniotic fluid (Anderson et al., 2004; Sobel, 2020a).

## Analytical Validity

Microscopic examination of normal vaginal discharge reveals a predominance of squamous epithelial cells, rare polymorphonuclear leukocytes (PMNs), and *Lactobacillus* species. The primary goal of the examination is to look for candidal buds or hyphae, motile trichomonads, epithelial cells studded with adherent coccobacilli (clue cells), and increased numbers of PMNs (Sobel, 2020a). The microscopic evaluation of BV is usually based on Amsel criteria (Amsel et al., 1983). Amsel criteria state that the presence of at least three out of the following four criteria are indicative of a BV diagnosis: increased homogeneous thin vaginal discharge, pH secretion > 4.5, amine odor when potassium hydroxide 10% solution is added to a vaginal secretion sample, and the presence of clue cells in wet preparations (Amsel et al., 1983). If clinical criteria are used to

## Diagnosis of Vaginitis including Multi-target PCR Testing AHS – M2057

define infection, then reported sensitivity may range from 62 to 100 percent (Spiegel, 1991). Using Gram's stain as the standard for diagnosing BV, the sensitivity of Amsel criteria for diagnosis of BV is over 90 percent and specificity is 77 percent (Landers et al., 2004). The Nugent score is also available as a Gram staining scoring system to diagnose BV based on vaginal swab samples (Amegashie et al., 2017). Because BV represents complex changes in the vaginal flora, vaginal culture has **no** role in diagnosis. If microscopy is not available, commercial diagnostic testing methods (eg, rapid antigen and nucleic acid amplification tests) are used for confirming the clinical suspicion of BV. Polymerase chain reaction (PCR)-based assays to quantify BV-associated bacteria (Cartwright et al., 2012; Menard, Fenollar, Henry, Bretelle, & Raoult, 2008) have good sensitivity and specificity compared with standard clinical tests (Dumonceaux et al., 2009; Menard et al., 2010). However, they are expensive and of limited utility (Sobel, 2020b).

Trichomoniasis can be diagnosed by the presence of motile trichomonads on wet mount, but it is identified in only 60 to 70 percent of culture-confirmed cases. Culture on Diamond's medium was considered the gold standard method for diagnosing a *T. vaginalis* infection (Workowski & Bolan, 2015); however, nucleic acid amplification tests (Baron et al., 2013) have become the accepted gold standard for the diagnosis of *T. vaginalis*. One study found the sensitivities for *T. vaginalis* using wet mount, culture, rapid antigen testing, and transcription-mediated amplification testing were 65, 96, 90, and 98 percent, respectively (Huppert et al., 2007). Coexistence of *T. vaginalis* and BV pathogens is common, with coinfection rates of 60 to 80 percent (Sobel, 2019; Sobel, Subramanian, Foxman, Fairfax, & Gygax, 2013).

Microscopy is negative in up to 50 percent of patients with culture-confirmed VVC (Sobel, 1985). Since there are no reliable point of care tests for *Candida* available in the United States (Abbott, 1995; Chatwani et al., 2007; Dan, Leshem, & Yeshaya, 2010; Hopwood, Evans, & Carney, 1985; Marot-Leblond et al., 2009; Matsui et al., 2009), culture must be obtained. PCR methods have high sensitivity and specificity and a shorter turn-around time than culture (Diba, Namaki, Ayatollahi, & Hanifian, 2012; Mahmoudi Rad, Zafarghandi, Amel Zabihi, Tavallae, & Mirdamadi, 2012; Tabrizi, Pirota, Rudland, & Garland, 2006; Weissenbacher et al., 2009), but they are costly and offer no proven benefit over culture in symptomatic women (Sobel, 2020c).

Lynch et al. (2019) collected vaginal swabs from 93 women in a cross-sectional study; results from microscopy were compared to two molecular approaches (a qPCR assay with a BV interpretive algorithm and a microbiome profiling test of the 16S rRNA gene produced by Illumina) (Lynch et al., 2019). Results show that “Microscopy plus BV Nugent score had 76% overall agreement with the qPCR plus BV interpretive algorithm method”; further, “Microscopic identification of *Candida* versus that by qPCR had 94% agreement (9 positive, 78 negative) (Lynch et al., 2019).” The qPCR assays gave additional information regarding the types of bacteria present, and the 16S microbiome analysis identified differentiating patterns between BV, aerobic vaginitis (AV), and *Lactobacillus* type infections.

Cartwright, Pherson, Harris, Clancey, and Nye (2018) have published data regarding the clinical validity of a PCR-based assay for the detection of BV. This multicenter study included 1579 patients and compared PCR results to samples realized by both the Nugent gram stain and a clinical evaluation using Amsel criteria. Next-generation sequencing was used to confirm differing results. After the resolution of discordant test results using next-generation sequencing, the BV-PCR assay reported a sensitivity of 98.7%, a specificity of 95.9%, a positive predictive value of 92.9% and a negative predictive value of 96.9% (Cartwright et al., 2018). These results show that this PCR-based assay can diagnose BV in symptomatic women efficiently.

### *Clinical Validity and Utility*

As previously stated, microscopy, rather than bacterial culture, is the standard of care for diagnosing BV, and commercially available tests are available in the absence of microscopy but

## Diagnosis of Vaginitis including Multi-target PCR Testing AHS – M2057

are not widely used. A study of 176 women using the Affirm VP III test (a DNA hybridization probe test that identifies high concentrations of *G. vaginalis*) reported comparable results to wet mount examination with no false positives and only three false negatives for *T. vaginalis*, and three false positives and four false negatives for *G. vaginalis* (Briselden & Hillier, 1994). This test “takes less than one hour to perform and is the best option when findings on physical examination suggest BV... but microscopy cannot be performed to look for clue cells (Sobel, 2020b).”

The OSOM BVBlue chromogenic diagnostic point-of-care test is a CLIA-waived test with a reported 10 minute read time. One study of 173 pregnant women reported a sensitivity and specificity of 94% and 96% respectively, as compared to Gram stain score (Sumeksri, Kopraser, & Panichkul, 2005). These results were comparable to the previously reported values of 91.7% sensitivity and 97.8% specificity in an earlier, smaller study of non-menstruating women (n=57) (Myziuk, Romanowski, & Johnson, 2003). A larger study (n=288 women) reported a sensitivity of 88% and specificity of 91% as compared to the Amstel criteria. The authors of this report concluded that women who “are not in settings where the conventional diagnostic methods are either practical or possible... would greatly benefit from access to rapid and reliable point-of-care tests to improve the diagnosis and management of BV (Bradshaw et al., 2005).”

The FDA approved the use of the BD MAX Vaginal Panel as “an automated qualitative *in vitro* diagnostic test for the direct detection of DNA targets from bacteria associated with BV (qualitative results reported based on detection and quantitation of targeted organism markers), *Candida* species associated with vulvovaginal candidiasis, and *Trichomonas vaginalis* from vaginal swabs in patients who are symptomatic for vaginitis/vaginosis. The test utilizes real-time PCR for the amplification of specific DNA targets and utilizes fluorogenic target-specific hybridization probes to detect and differentiate DNA (FDA, 2016).” A 2017 cross-sectional, multi-site study into the clinical validation of this system (n=1740 symptomatic women) reported a sensitivity and specificity of 90.9% and 94.1%, respectively for the *Candida* group and 90.5% sensitivity and 85.8% specificity for BV. For *C. glabrata* specifically, the assay had only 75.9% sensitivity but 99.7% specificity. For trichomoniasis, the sensitivity and specificity were 93.1% and 99.3%, respectively (Gaydos et al., 2017). These researchers also compared the results of this test to clinician assessment. Again, to qualify for the study, the women must have at least one symptom of BV. Using Amsel’s criteria, the investigational test sensitivity was 92.7% as compared to the 75.6% sensitivity of the clinician assessment. The authors conclude, “The investigational test showed significantly higher sensitivity for detecting vaginitis, involving more than one cause, than did clinician diagnosis. Taken together, these results suggest that a molecular investigational test can facilitate accurate detection of vaginitis (Schwebke et al., 2018).” It should be noted, however, that these studies only included symptomatic women, and, therefore, the possible clinical nonspecificity (i.e. instances where an asymptomatic woman would test positive) is not addressed. Sherrard (2019) compared BV, candidiasis, and trichomoniasis diagnostic results from the BD MAX Vaginal Panel to a current test used in a UK specialist sexual health service center. The authors reported that the BD MAX Vaginal Panel had a sensitivity of 86.4% and specificity of 86.0% for *Candida* species, and a sensitivity of 94.4% and specificity of 79% for BV; the specificity for BV was lower in this study than what has been previously reported (Sherrard, 2019).

SureSwab®(Quest Diagnostics, Inc.) is a multi-target PCR test using RT-PCR to screen for a number of microorganisms involved in vaginal flora imbalances, including *B. vaginalis*, *T. vaginalis*, *C. albicans*, *C. glabrata*, *C. tropicalis*, and *C. parapsilosis*, from a vaginal swab. The swab can be collected either by a physician or the patient (Quest, 2018a). Similarly, Quest Diagnostics also offers the SureSwab® Vaginosis/Vaginitis Plus test, which tests for the presence of *Chlamydia trachomatis* and *Neisseria gonorrhoeae* in addition to the microorganisms of the SureSwab® test (Quest, 2018b). The test is based on the unique set of primers synthesized by a CDC research team to identify *Candida* that purports to diagnose vulvovaginal candidiasis while ruling out other genital infections (CDC, 2016). The CDC research group, led by Dr. C.J.

## Diagnosis of Vaginitis including Multi-target PCR Testing AHS – M2057

Morrison, developed the DNA probes to identify medically important *Candida* species by the internal transcribed spacer 2 region of ribosomal DNA. The specific hybridization was measured by a sample-to-background ratio of 58.7, 53.2, 46.9, 59.9, and 54.7 for *C. albicans*, *C. tropicalis*, *C. glabrata*, *C. parapsilosis*, and *C. krusei*, respectively. The negative control sample-to-background ratio was 0.9 (Das, Brown, Kellar, Holloway, & Morrison, 2006).

The OSOM *Trichomonas vaginalis* (TV) Rapid Test by Sekisui Diagnostics is “an antigen-detection test using immunochromatographic capillary flow dipstick technology that can be performed at the point of care (CDC, 2015b).” The diagnostic accuracy of the OSOM TV Rapid assay was tested against the common laboratory-based Anyplex II STI-7 Detection in a South African cross-sectional study; all irregular results were further tested with the Fast Track Diagnostics (FTD) STD9 assay (Garrett et al., 2019). Vaginal swabs from 247 women were tested for this study. “The sensitivity and specificity of OSOM TV were 75.0% (45.0-100) and 100% (100-100)”, respectively, showing a very high specificity and lower sensitivity (Garrett et al., 2019).

The AMPLISwab™ by MedLabs is a comprehensive test created to assess the different organisms responsible for a variety of female genital tract infections, including causative pathogens for cervicitis, nongonococcal urethritis, pelvic inflammatory disease and infertility, sexually transmitted infections, and vaginitis (e.g., bacterial vaginosis, candidiasis and trichomoniasis). The test requires one swab to test for 23 total organisms, broken down into four categories (7 yeast, 12 bacteria and 1 reference bacteria, 1 parasite, and 2 types of herpes viruses), employing testing methodologies such as automated DNA/RNA extraction, transcription mediated amplification (TMA), and real-time polymerase chain reaction (RT-PCR) for the quantification of select organisms implicated in bacterial vaginosis (MedLabs, 2015).

The multiplex PCR assay SmartJane™ measures a specimen’s vaginal flora (such as *Lactobacillus iners* or *Treponema pallidum*). The test proposes that the results can provide a health snapshot of the environment tested based on the levels of microorganisms detected. The procedure for the test requires the user to self-sample by collecting a vaginal swab and sending the sample back to Ubiome where it is analyzed. The labs use Precision Sequencing technology to extract DNA from the microorganisms in the sample and Illumina Next-Generation to sequence the targeted genes. Then, phylogenetic algorithms are used to analyze and organize the DNA from those microorganisms. Finally, a clinical report detailing the levels of the targeted microorganisms is sent to the user and medical provider (Ubiome, 2018). The report contains measurements of its targeted microorganisms, informing the patient whether those measurements are within the normal reference ranges for certain conditions, and whether certain high danger pathogens are present. The manufacturers state that on average SmartJane™ has a sensitivity and specificity for the species of microorganism of 99.4% and 100.0%, respectively. SmartJane™ tests for 19 different HPV strains and common pathological agents involved in sexually transmitted infections in addition to more than 20 different microorganisms involved in BV, including *G. vaginalis* (Ubiome, 2017).

Even though studies have shown that PCR methods have a higher specificity and sensitivity than culture and shorter turn-around time in identifying *Candida* (Diba et al., 2012; Mahmoudi Rad et al., 2012; Tabrizi et al., 2006; Weissenbacher et al., 2009), their use may be adding to clinical nonspecificity. Tabrizi et al. (2006) reported that PCR “detected four additional *Candida albicans*, three *Candida parapsilosis* and one *Candida tropicalis* when compared with culture. All but one case additionally detected by PCR were found in patients with no VVC symptoms (Tabrizi et al., 2006).” These data support the earlier findings by Giraldo et al. (2000) where, unlike culture testing, “*Candida* was identified by PCR in a similar proportion of patients with previous recurrent vulvovaginal candidiasis (30%) and in controls (28.8%).” Taken together, these studies indicate that, even though PCR is more sensitive than culture, it may be identifying cases of *Candida* in asymptomatic women that are clinically irrelevant.

# Diagnosis of Vaginitis including Multi-target PCR Testing AHS – M2057

## Applicable State and Federal Regulations

A search of the term “vaginitis” on the FDA Device database on 01/29/2021 yielded 145 records. Additionally, many labs have developed specific tests that they must validate and perform in house. These laboratory-developed tests (LDTs) are regulated by the Centers for Medicare and Medicaid (CMS) as high-complexity tests under the Clinical Laboratory Improvement Amendments of 1988 (CLIA '88). As an LDT, the U. S. Food and Drug Administration has not approved or cleared this test; however, FDA clearance or approval is not currently required for clinical use.

## Guidelines and Recommendations

### Centers for Disease Control and Prevention (CDC) (CDC, 2015a, 2015b, 2015c)

The CDC published recommendations for the evaluation of diseases characterized by vaginal discharge in the 2015 Sexually Transmitted Diseases Treatment Guidelines (CDC, 2015a). These guidelines state, “Various diagnostic methods are available to identify the etiology of an abnormal vaginal discharge...In the clinician’s office, the cause of vaginal symptoms might be determined by pH, a potassium hydroxide (KOH) test, and microscopic examination of fresh samples of the discharge” and “In settings where pH paper, KOH, and microscopy are not available, alternative commercially available point-of-care tests or clinical laboratory testing can be used to diagnose vaginitis (CDC, 2015a).”

For the evaluation of BV, the CDC recommends that “BV can be diagnosed by the use of clinical criteria (i.e., Amsel’s Diagnostic Criteria) (Amsel et al., 1983) or Gram stain”; further, “Other tests, including Affirm VP III (Becton Dickinson, Sparks, MD), a DNA hybridization probe test for high concentrations of *G. vaginalis*, and the OSOM BV Blue test (Sekisui Diagnostics, Framingham, MA), which detects vaginal fluid sialidase activity, have acceptable performance characteristics compared with Gram stain. Although a prolineaminopeptidase card test is available for the detection of elevated pH and trimethylamine, it has low sensitivity and specificity and therefore is not recommended. PCR has been used in research settings for the detection of a variety of organisms associated with BV, but evaluation of its clinical utility is still underway. Detection of specific organisms might be predictive of BV by PCR (Cartwright et al., 2012; Fredricks, Fiedler, Thomas, Oakley, & Marrazzo, 2007). Additional validation is needed before these tests can be recommended to diagnose BV. Culture of *G. vaginalis* is not recommended as a diagnostic tool because it is not specific. Cervical Pap tests have no clinical utility for the diagnosis of BV because of their low sensitivity and specificity (CDC, 2015a).” The guidelines also state that “evidence is insufficient to recommend routine screening for BV in asymptomatic pregnant women at high or low risk for preterm delivery for the prevention of preterm birth (CDC, 2015a)”, which is in compliance with the 2008 USPSTF recommendations (USPSTF, 2008).

For the evaluation of vulvovaginal candidiasis, the CDC recommends: “Examination of a wet mount with KOH preparation should be performed for all women with symptoms or signs of VVC, and women with a positive result should be treated. For those with negative wet mounts but existing signs or symptoms, vaginal cultures for *Candida* should be considered...PCR testing for yeast is not FDA-cleared; culture for yeast remains the gold standard for diagnosis (CDC, 2015c).” DNA hybridization probe tests are not addressed.

For the evaluation of Trichomoniasis, the CDC recommends: “Diagnostic testing for *T. vaginalis* should be performed in women seeking care for vaginal discharge... The use of highly sensitive and specific tests is recommended for detecting *T. vaginalis*. Among women, NAAT is highly sensitive, often detecting three to five times more *T. vaginalis* infections than wet-mount microscopy, a method with poor sensitivity (51%–65%) (CDC, 2015b; Hollman, Coupey, Fox, & Herold, 2010; Roth et al., 2011).” Regarding point of care testing, it is stated that “Other FDA-cleared tests to detect *T. vaginalis* in vaginal secretions include the OSOM *Trichomonas* Rapid

## Diagnosis of Vaginitis including Multi-target PCR Testing AHS – M2057

Test (Sekisui Diagnostics, Framingham, MA), an antigen-detection test using immunochromatographic capillary flow dipstick technology that can be performed at the point of care, and the Affirm VP III (Becton Dickinson, Sparks, MD), a DNA hybridization probe test that evaluates for *T. vaginalis*, *G. vaginalis*, and *Candida albicans*. The results of the OSOM *Trichomonas* Rapid Test are available in approximately 10 minutes, with sensitivity 82%–95% and specificity 97%–100% (Campbell, Woods, Lloyd, Elsayed, & Church, 2008; Huppert et al., 2007). Self-testing might become an option, as a study of 209 young women aged 14–22 years found that >99% could correctly perform and interpret her own self-test using the OSOM assay, with a high correlation with clinician interpretation (96% agreement,  $\kappa = 0.87$ ) (Huppert et al., 2010). The results of the Affirm VP III are available within 45 minutes. Sensitivity and specificity are 63% and 99.9%, respectively, compared with culture and TMA; sensitivity might be higher among women who are symptomatic (Andrea & Chapin, 2011; Brown, Fuller, Jasper, Davis, & Wright, 2004; CDC, 2015b).”

### **American Academy of Family Physicians (AAFP) (Hainer & Gibson, 2011; Paladine & Desai, 2018)**

The AAFP published an article (Hainer & Gibson, 2011) on the diagnosis of vaginitis which states that: “Physicians traditionally diagnose vaginitis using the combination of symptoms, physical examination, pH of vaginal fluid, microscopy, and the whiff test. When combined, these tests have a sensitivity and specificity of 81 and 70 percent, respectively, for BV; 84 and 85 percent for vulvovaginal candidiasis; and 85 and 100 percent for trichomoniasis when compared with the DNA probe standard...A cost-effectiveness analysis of diagnostic strategies for vaginitis undiagnosed by pelvic examination, wet-mount preparation, and related office tests showed that the least expensive strategy was to perform yeast culture, gonorrhea and chlamydia probes at the initial visit, and Gram stain and *Trichomonas* culture only when the vaginal pH exceeded 4.9. Other strategies cost more and increased duration of symptoms by up to 1.3 days (Hainer & Gibson, 2011).”

In 2018, the AAFP has published the following guidelines:

- “Symptoms alone cannot differentiate between the causes of vaginitis. Office-based or laboratory testing should be used with the history and physical examination findings to make the diagnosis. (C evidence rating)
- Do not obtain culture for the diagnosis of bacterial vaginosis because it represents a polymicrobial infection. (C evidence rating)
- Nucleic acid amplification testing is recommended for the diagnosis of trichomoniasis in symptomatic or high-risk women. (C evidence rating) (Paladine & Desai, 2018).”

### **U.S. Preventive Services Task Force Recommendations (USPSTF) (Owens et al., 2020; USPSTF, 2008)**

In 2020, the USPSTF published recommendations discouraging the use of screening for BV in pregnancy: “The USPSTF recommends against screening for bacterial vaginosis in pregnant persons not at increased risk for preterm delivery”. On a similar note, the USPSTF maintains its 2008 recommendation stating “that the current evidence is insufficient to assess the balance of benefits and harms of screening for bacterial vaginosis in pregnant persons at increased risk for preterm delivery” (Owens et al., 2020).

### **American College of Obstetrics and Gynecology (ACOG) (ACOG, 2006, 2017, 2018, 2020)**

ACOG published recommendations (ACOG, 2006) for the evaluation of vaginitis in 2006, and reaffirmed in 2017 (ACOG, 2017, 2018), which state: “Evaluation of women with vaginitis should include a focused history about the entire spectrum of vaginal symptoms, including change in discharge, vaginal malodor, itching, irritation, burning, swelling, dyspareunia, and dysuria.” Further, “During speculum examination, samples should be obtained for vaginal pH, amine (“whiff”) test, and saline (wet mount) and 10% potassium hydroxide (KOH) microscopy. The pH



## Diagnosis of Vaginitis including Multi-target PCR Testing AHS – M2057

and amine testing can be performed either through direct measurement or by colorimetric testing.” With a Level B recommendation, ACOG states, “Microscopy is the first line for diagnosing vulvovaginal candidiasis and trichomoniasis. In selected patients, culture for yeast and *T. vaginalis* should be obtained in addition to standard office-based testing.” Additionally, “A vaginal Gram stain for Nugent scoring of the bacterial flora may help to identify patients with BV. Other currently available ancillary tests for diagnosing vaginal infections include rapid tests for enzyme activity from BV-associated organisms, *Trichomonas vaginalis* antigen, and point-of care testing for DNA of *G. vaginalis*, *T. vaginalis*, and *Candida* species; however, the role of these tests in the proper management of patients with vaginitis is unclear. Depending on risk factors, DNA amplification tests can be obtained for *Neisseria gonorrhoeae* and *Chlamydia trachomatis* (ACOG, 2006).”

The ACOG published in 2020 Practice Bulletin Number 215 on vaginitis in nonpregnant patients. In these guidelines, the ACOG made these recommendations for diagnostic testing based on good and consistent scientific evidence (Level B):

- “The use of Amsel clinical criteria or Gram stain with Nugent scoring is recommended for the diagnosis of bacterial vaginosis.”
- “Nucleic acid amplification testing is recommended for the diagnosis of trichomoniasis.”
- “In a symptomatic patient, diagnosis of vulvovaginal candidiasis requires one of the following two findings: 1) visualization of spores, pseudohyphae, or hyphae on wet-mount microscopy or 2) vaginal fungal culture or commercial diagnostic test results positive for *Candida* species.”

The ACOG also published recommendations based on limited or inconsistent scientific evidence (Level B), along with a series of recommendations based on consensus and expert opinion (Level C). Those relating to diagnostic testing are reported below:

- “Patients should be retested within 3 months after treatment for *T vaginalis* because of the high rates of infection recurrence.” (Level B)
- “Pap tests are not reliable for the diagnosis of vaginitis. Diagnostic confirmation is recommended for incidental findings of vulvovaginal candidiasis, bacterial vaginosis, or trichomoniasis on a Pap test.” (Level B)
- “A complete medical history, physical examination of the vulva and vagina, and clinical testing of vaginal discharge (ie. pH testing, a potassium hydroxide [KOH] “whiff test”, and microscopy) are recommended for the initial evaluation of patients with vaginitis symptoms.” (Level C)

### **Infectious Diseases Society of America (IDSA) Clinical Practice Guidelines (Miller et al., 2018; Pappas et al., 2016)**

IDSA has published an updated clinical guideline (Pappas et al., 2016) for the management of candidiasis in which recommendations include diagnosing vulvovaginal candidiasis before proceeding with empiric antifungal therapy. The usual diagnosis is clinical based on signs and symptoms of vaginitis such as pruritus, irritation, vaginal soreness, vulvar edema, erythema and many others. Clinical signs and symptoms are nonspecific and could be attributed to causes other than vulvovaginal candidiasis. Therefore, authors recommend confirming clinical diagnosis by a wet -mount preparation with saline and 10% KOH to demonstrate the presence of yeast and a normal pH. In cases where signs and symptoms are suggestive of vulvovaginal candidiasis, but microscopic findings and pH are negative, culture testing confirms the diagnosis according to published guidelines. The IDSA also discusses the possible use of PCR in diagnosing invasive candidiasis, even though the guidelines later state that “Cultures of blood or other samples collected under sterile conditions have long been considered diagnostic gold standards for invasive candidiasis...The role of PCR in testing samples other than blood is not established (Pappas et al., 2016).”

# Diagnosis of Vaginitis including Multi-target PCR Testing AHS – M2057

In the 2018 IDSA *A guide to Utilization of the Microbiology Laboratory for Diagnosis of Infectious Diseases*, the IDSA states, “For vaginosis (altered vaginal flora) a Gram stain and recently available microbiome-based assays are more specific than culture and probe testing for *Gardnerella vaginalis* alone... A number of point-of-care tests can be performed from a vaginal discharge specimen while the patient is in the healthcare setting. Although point-of-care tests are popular, the sensitivity and specificity for making a specific diagnosis vary widely and these assays, while rapid, are often diagnostically poor (Miller et al., 2018).” The IDSA notes that the FDA has approved the use of the Max Vaginal Panel by Becton Dickinson in symptomatic females. “Preliminary data show greater specificity of this approach compared to methods that identify only *G. vaginalis*, as well as consistency in both reproducible as well as standardized results (Miller et al., 2018).”

## **Society of Obstetricians and Gynecologists of Canada (SOGC) (van Schalkwyk & Yudin, 2015; Yudin & Money, 2017)**

The SOGC published guidelines for the screening and management of BV in pregnancy. These guidelines state that the following:

- “In symptomatic pregnant women, testing for and treatment of bacterial vaginosis is recommended for symptom resolution. Diagnostic criteria are the same for pregnant and non-pregnant women (I-A).
- Asymptomatic women and women without identified risk factors for preterm birth should not undergo routine screening for or treatment of bacterial vaginosis (I-B).
- Women at increased risk for preterm birth may benefit from routine screening for and treatment of bacterial vaginosis (I-B).
- Testing should be repeated one month after treatment to ensure that cure was achieved (III-L) (Yudin & Money, 2017).”

The SOGC also published guidelines regarding the screening and management of trichomoniasis, VVC, and BV. These guidelines state that “Bacterial vaginosis should be diagnosed using either clinical (Amsel’s) or laboratory (Gram stain with objective scoring system) criteria (II-2A) (van Schalkwyk & Yudin, 2015).”

## **Billing/Coding/Physician Documentation Information**

---

This policy may apply to the following codes. Inclusion of a code in this section does not guarantee that it will be reimbursed. For further information on reimbursement guidelines, please see Administrative Policies on the Blue Cross Blue Shield of North Carolina web site at [www.bcbsnc.com](http://www.bcbsnc.com). They are listed in the Category Search on the Medical Policy search page.

*Applicable service codes: 81513, 81514, 82120, 83986, 87070, 87149, 87150, 87210, 87480, 87481, 87482, 87510, 87511, 87512, 87660, 87661, 87797, 87798, 87799, 87800, 87808, 87905, Q0111*

BCBSNC may request medical records for determination of medical necessity. When medical records are requested, letters of support and/or explanation are often useful, but are not sufficient documentation unless all specific information needed to make a medical necessity determination is included.

## **Scientific Background and Reference Sources**

---

Abbott, J. (1995). Clinical and microscopic diagnosis of vaginal yeast infection: a prospective analysis. *Ann Emerg Med*, 25(5), 587-591. Retrieved from <http://dx.doi.org/>

# Diagnosis of Vaginitis including Multi-target PCR Testing AHS – M2057

ACOG. (2006). ACOG Practice Bulletin. Clinical management guidelines for obstetrician-gynecologists, Number 72, May 2006: Vaginitis. *Obstet Gynecol*, 107(5), 1195-1206. Retrieved from <http://dx.doi.org/>

ACOG. (2017). College Publications. *Obstet Gynecol*, 129(6), 1147-1148. doi:10.1097/aog.0000000000002107

ACOG. (2018). Practice Bulletins. Retrieved from <https://www.acog.org/Clinical-Guidance-and-Publications/Practice-Bulletins-List>

ACOG. (2020). Vaginitis in Nonpregnant Patients: ACOG Practice Bulletin, Number 215. *Obstetrics & Gynecology*, 135(1), e1-e17. doi:10.1097/aog.0000000000003604

Amegashie, C. P., Gilbert, N. M., Peipert, J. F., Allsworth, J. E., Lewis, W. G., & Lewis, A. L. (2017). Relationship between nugent score and vaginal epithelial exfoliation. *PLoS One*, 12(5), e0177797. doi:10.1371/journal.pone.0177797

Amsel, R., Totten, P. A., Spiegel, C. A., Chen, K. C., Eschenbach, D., & Holmes, K. K. (1983). Nonspecific vaginitis. Diagnostic criteria and microbial and epidemiologic associations. *Am J Med*, 74(1), 14-22. Retrieved from <http://dx.doi.org/>

Anderson, M. R., Klink, K., & Cochrane, A. (2004). Evaluation of vaginal complaints. *Jama*, 291(11), 1368-1379. doi:10.1001/jama.291.11.1368

Andrea, S. B., & Chapin, K. C. (2011). Comparison of Aptima Trichomonas vaginalis transcription-mediated amplification assay and BD affirm VPIII for detection of T. vaginalis in symptomatic women: performance parameters and epidemiological implications. *J Clin Microbiol*, 49(3), 866-869. doi:10.1128/jcm.02367-10

Baron, E. J., Miller, J. M., Weinstein, M. P., Richter, S. S., Gilligan, P. H., Thomson, R. B., Jr., . . . Pritt, B. S. (2013). A guide to utilization of the microbiology laboratory for diagnosis of infectious diseases: 2013 recommendations by the Infectious Diseases Society of America (IDSA) and the American Society for Microbiology (ASM)(a). *Clin Infect Dis*, 57(4), e22-e121. doi:10.1093/cid/cit278

Bradshaw, C. S., Morton, A. N., Garland, S. M., Horvath, L. B., Kuzevska, I., & Fairley, C. K. (2005). Evaluation of a point-of-care test, BVBlue, and clinical and laboratory criteria for diagnosis of bacterial vaginosis. *J Clin Microbiol*, 43(3), 1304-1308. doi:10.1128/jcm.43.3.1304-1308.2005

Briselden, A. M., & Hillier, S. L. (1994). Evaluation of affirm VP Microbial Identification Test for Gardnerella vaginalis and Trichomonas vaginalis. *J Clin Microbiol*, 32(1), 148-152. Retrieved from <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC262986/>

Brown, H. L., Fuller, D. D., Jasper, L. T., Davis, T. E., & Wright, J. D. (2004). Clinical evaluation of affirm VPIII in the detection and identification of Trichomonas vaginalis, Gardnerella vaginalis, and Candida species in vaginitis/vaginosis. *Infect Dis Obstet Gynecol*, 12(1), 17-21. doi:10.1080/1064744042000210375

Campbell, L., Woods, V., Lloyd, T., Elsayed, S., & Church, D. L. (2008). Evaluation of the OSOM Trichomonas rapid test versus wet preparation examination for detection of Trichomonas vaginalis vaginitis in specimens from women with a low prevalence of infection. *J Clin Microbiol*, 46(10), 3467-3469. doi:10.1128/jcm.00671-08

Cartwright, C. P., Lembke, B. D., Ramachandran, K., Body, B. A., Nye, M. B., Rivers, C. A., & Schwabke, J. R. (2012). Development and validation of a semiquantitative, multitarget PCR assay

## Diagnosis of Vaginitis including Multi-target PCR Testing AHS – M2057

for diagnosis of bacterial vaginosis. *J Clin Microbiol*, 50(7), 2321-2329. doi:10.1128/jcm.00506-12

Cartwright, C. P., Pherson, A. J., Harris, A. B., Clancey, M. S., & Nye, M. B. (2018). Multicenter study establishing the clinical validity of a nucleic-acid amplification-based assay for the diagnosis of bacterial vaginosis. *Diagn Microbiol Infect Dis*, 92(3), 173-178. doi:10.1016/j.diagmicrobio.2018.05.022

CDC. (2015a, 06/04/2015). Bacterial Vaginosis - 2015 STD Treatment Guidelines. Retrieved from <https://www.cdc.gov/std/tg2015/bv.htm>

CDC. (2015b, 08/12/2016). Trichomoniasis - 2015 STD Treatment Guidelines. Retrieved from <https://www.cdc.gov/std/tg2015/trichomoniasis.htm>

CDC. (2015c, 06/04/2015). Vulvovaginal Candidiasis - 2015 STD Treatment Guidelines. Retrieved from <https://www.cdc.gov/std/tg2015/candidiasis.htm>

CDC. (2016, 10/11/2017). A Diagnostic Test to Detect Different Vaginal Yeast Infection Types. *Office of the Associate Director for Science (OADS) Success Stories*. Retrieved from <https://www.cdc.gov/od/science/technology/techtransfer/successstories/candida.htm>

Chatwani, A. J., Mehta, R., Hassan, S., Rahimi, S., Jeronis, S., & Dandolu, V. (2007). Rapid testing for vaginal yeast detection: a prospective study. *Am J Obstet Gynecol*, 196(4), 309.e301-304. doi:10.1016/j.ajog.2006.11.025

Dan, M., Leshem, Y., & Yeshaya, A. (2010). Performance of a rapid yeast test in detecting *Candida* spp. in the vagina. *Diagn Microbiol Infect Dis*, 67(1), 52-55. doi:10.1016/j.diagmicrobio.2009.12.010

Das, S., Brown, T. M., Kellar, K. L., Holloway, B. P., & Morrison, C. J. (2006). DNA probes for the rapid identification of medically important *Candida* species using a multianalyte profiling system. *FEMS Immunol Med Microbiol*, 46(2), 244-250. doi:10.1111/j.1574-695X.2005.00031.x

Diba, K., Namaki, A., Ayatollahi, H., & Hanifian, H. (2012). Rapid identification of drug resistant *Candida* species causing recurrent vulvovaginal candidiasis. *Med Mycol J*, 53(3), 193-198. Retrieved from <http://dx.doi.org/>

Dumonceaux, T. J., Schellenberg, J., Goleski, V., Hill, J. E., Jaoko, W., Kimani, J., . . . Severini, A. (2009). Multiplex detection of bacteria associated with normal microbiota and with bacterial vaginosis in vaginal swabs by use of oligonucleotide-coupled fluorescent microspheres. *J Clin Microbiol*, 47(12), 4067-4077. doi:10.1128/jcm.00112-09

Ellis, I., Lerch, M. M., & Whitcomb, D. C. (2001). Genetic testing for hereditary pancreatitis: guidelines for indications, counselling, consent and privacy issues. *Pancreatology*, 1(5), 405-415. Retrieved from <http://dx.doi.org/>

Eschenbach, D. A., Davick, P. R., Williams, B. L., Klebanoff, S. J., Young-Smith, K., Critchlow, C. M., & Holmes, K. K. (1989). Prevalence of hydrogen peroxide-producing *Lactobacillus* species in normal women and women with bacterial vaginosis. *J Clin Microbiol*, 27(2), 251-256. Retrieved from <http://dx.doi.org/>

FDA. (2016). EVALUATION OF AUTOMATIC CLASS III DESIGNATION FOR BD MAX Vaginal Panel. Retrieved from [https://www.accessdata.fda.gov/cdrh\\_docs/reviews/DEN160001.pdf](https://www.accessdata.fda.gov/cdrh_docs/reviews/DEN160001.pdf)

# Diagnosis of Vaginitis including Multi-target PCR Testing AHS – M2057

- Fredricks, D. N., Fiedler, T. L., Thomas, K. K., Oakley, B. B., & Marrazzo, J. M. (2007). Targeted PCR for detection of vaginal bacteria associated with bacterial vaginosis. *J Clin Microbiol*, *45*(10), 3270-3276. doi:10.1128/jcm.01272-07
- Garrett, N., Mitchev, N., Osman, F., Naidoo, J., Dorward, J., Singh, R., . . . Mindel, A. (2019). Diagnostic accuracy of the Xpert CT/NG and OSOM Trichomonas Rapid assays for point-of-care STI testing among young women in South Africa: a cross-sectional study. *BMJ Open*, *9*(2), e026888. doi:10.1136/bmjopen-2018-026888
- Gaydos, C. A., Beqaj, S., Schwebke, J. R., Lebed, J., Smith, B., Davis, T. E., . . . Cooper, C. K. (2017). Clinical Validation of a Test for the Diagnosis of Vaginitis. *Obstet Gynecol*, *130*(1), 181-189. doi:10.1097/aog.0000000000002090
- Giraldo, P., von Nowaskonski, A., Gomes, F. A., Linhares, I., Neves, N. A., & Witkin, S. S. (2000). Vaginal colonization by *Candida* in asymptomatic women with and without a history of recurrent vulvovaginal candidiasis. *Obstet Gynecol*, *95*(3), 413-416. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/?term=10711554%5Buid%5D>
- Hainer, B. L., & Gibson, M. V. (2011). Vaginitis: Diagnosis and Treatment. *American Family Physician*, *83*(7), 807-815. Retrieved from /afp/2011/0401/p807.pdf
- Hill, G. B. (1993). The microbiology of bacterial vaginosis. *Am J Obstet Gynecol*, *169*(2 Pt 2), 450-454. Retrieved from <http://dx.doi.org/>
- Hollman, D., Coupey, S. M., Fox, A. S., & Herold, B. C. (2010). Screening for *Trichomonas vaginalis* in high-risk adolescent females with a new transcription-mediated nucleic acid amplification test (NAAT): associations with ethnicity, symptoms, and prior and current STIs. *J Pediatr Adolesc Gynecol*, *23*(5), 312-316. doi:10.1016/j.jpjag.2010.03.004
- Hopwood, V., Evans, E. G., & Carney, J. A. (1985). Rapid diagnosis of vaginal candidosis by latex particle agglutination. *J Clin Pathol*, *38*(4), 455-458. Retrieved from <http://dx.doi.org/>
- Huppert, J. S., Hesse, E., Kim, G., Kim, M., Agreda, P., Quinn, N., & Gaydos, C. (2010). Adolescent women can perform a point-of-care test for trichomoniasis as accurately as clinicians. *Sex Transm Infect*, *86*(7), 514-519. doi:10.1136/sti.2009.042168
- Huppert, J. S., Mortensen, J. E., Reed, J. L., Kahn, J. A., Rich, K. D., Miller, W. C., & Hobbs, M. M. (2007). Rapid antigen testing compares favorably with transcription-mediated amplification assay for the detection of *Trichomonas vaginalis* in young women. *Clin Infect Dis*, *45*(2), 194-198. doi:10.1086/518851
- Jones, A. (2019). Bacterial Vaginosis: A Review of Treatment, Recurrence, and Disparities. *The Journal for Nurse Practitioners*, *15*(6), 420-423. doi:<https://doi.org/10.1016/j.nurpra.2019.03.010>
- Kairys, N., & Garg, M. (2020). *Bacterial Vaginosis*. Retrieved from <https://www.ncbi.nlm.nih.gov/books/NBK459216/>
- Kissinger, P. (2015). Epidemiology and treatment of trichomoniasis. *Curr Infect Dis Rep*, *17*(6), 484. doi:10.1007/s11908-015-0484-7
- Lamont, R. F., Sobel, J. D., Akins, R. A., Hassan, S. S., Chaiworapongsa, T., Kusanovic, J. P., & Romero, R. (2011). The vaginal microbiome: new information about genital tract flora using molecular based techniques. *Bjog*, *118*(5), 533-549. doi:10.1111/j.1471-0528.2010.02840.x

## Diagnosis of Vaginitis including Multi-target PCR Testing AHS – M2057

- Landers, D. V., Wiesenfeld, H. C., Heine, R. P., Krohn, M. A., & Hillier, S. L. (2004). Predictive value of the clinical diagnosis of lower genital tract infection in women. *Am J Obstet Gynecol*, *190*(4), 1004-1010. doi:10.1016/j.ajog.2004.02.015
- Ling, Z., Kong, J., Liu, F., Zhu, H., Chen, X., Wang, Y., . . . Xiang, C. (2010). Molecular analysis of the diversity of vaginal microbiota associated with bacterial vaginosis. *BMC Genomics*, *11*, 488. doi:10.1186/1471-2164-11-488
- Lynch, T., Peirano, G., Lloyd, T., Read, R., Carter, J., Chu, A., . . . Church, D. (2019). Molecular Diagnosis of Vaginitis: Comparing Quantitative PCR and Microbiome Profiling Approaches to Current Microscopy Scoring. *J Clin Microbiol*, *57*(9). doi:10.1128/jcm.00300-19
- Mahmoudi Rad, M., Zafarghandi, A., Amel Zabihi, M., Tavallaee, M., & Mirdamadi, Y. (2012). Identification of Candida species associated with vulvovaginal candidiasis by multiplex PCR. *Infect Dis Obstet Gynecol*, *2012*, 872169. doi:10.1155/2012/872169
- Marot-Leblond, A., Nail-Billaud, S., Pilon, F., Beucher, B., Poulain, D., & Robert, R. (2009). Efficient diagnosis of vulvovaginal candidiasis by use of a new rapid immunochromatography test. *J Clin Microbiol*, *47*(12), 3821-3825. doi:10.1128/jcm.01168-09
- Matsui, H., Hanaki, H., Takahashi, K., Yokoyama, A., Nakae, T., Sunakawa, K., & Omura, S. (2009). Rapid detection of vaginal Candida species by newly developed immunochromatography. *Clin Vaccine Immunol*, *16*(9), 1366-1368. doi:10.1128/cvi.00204-09
- MedLabs. (2015). AMPLISwab™ Women's Health. Retrieved from <http://www.medlabdx.com/AmpliSwab.html>
- Menard, J. P., Fenollar, F., Henry, M., Bretelle, F., & Raoult, D. (2008). Molecular quantification of Gardnerella vaginalis and Atopobium vaginae loads to predict bacterial vaginosis. *Clin Infect Dis*, *47*(1), 33-43. doi:10.1086/588661
- Menard, J. P., Mazouni, C., Fenollar, F., Raoult, D., Boubli, L., & Bretelle, F. (2010). Diagnostic accuracy of quantitative real-time PCR assay versus clinical and Gram stain identification of bacterial vaginosis. *Eur J Clin Microbiol Infect Dis*, *29*(12), 1547-1552. doi:10.1007/s10096-010-1039-3
- Miller, J. M., Binnicker, M. J., Campbell, S., Carroll, K. C., Chapin, K. C., Gilligan, P. H., . . . Yao, J. D. (2018). A Guide to Utilization of the Microbiology Laboratory for Diagnosis of Infectious Diseases: 2018 Update by the Infectious Diseases Society of America and the American Society for Microbiology. *Clinical Infectious Diseases*, ciy381-ciy381. doi:10.1093/cid/ciy381
- Myziuk, L., Romanowski, B., & Johnson, S. C. (2003). BVBlue test for diagnosis of bacterial vaginosis. *J Clin Microbiol*, *41*(5), 1925-1928. Retrieved from <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC154737/>
- Owens, D. K., Davidson, K. W., Krist, A. H., Barry, M. J., Cabana, M., Caughey, A. B., . . . Wong, J. B. (2020). Screening for Bacterial Vaginosis in Pregnant Persons to Prevent Preterm Delivery: US Preventive Services Task Force Recommendation Statement. *Jama*, *323*(13), 1286-1292. doi:10.1001/jama.2020.2684
- Paladine, H. L., & Desai, U. A. (2018). Vaginitis: Diagnosis and Treatment. *Am Fam Physician*, *97*(5), 321-329. Retrieved from <https://www.aafp.org/afp/2018/0301/p321.html>
- Pappas, P. G., Kauffman, C. A., Andes, D. R., Clancy, C. J., Marr, K. A., Ostrosky-Zeichner, L., . . . Sobel, J. D. (2016). Clinical Practice Guideline for the Management of Candidiasis: 2016 Update



# Diagnosis of Vaginitis including Multi-target PCR Testing AHS – M2057

by the Infectious Diseases Society of America. *Clin Infect Dis*, 62(4), e1-50.  
doi:10.1093/cid/civ933

Quest. (2019a). SureSwab®, Bacterial Vaginosis/Vaginitis. *Test Directory*. Retrieved from <https://testdirectory.questdiagnostics.com/test/test-detail/15509/sureswab-bacterial-vaginosisvaginitis?p=r&q=Sureswab&cc=MASTER>

Quest. (2019b). SureSwab®, Bacterial Vaginosis/Vaginitis Plus. *Test Directory*. Retrieved from <https://testdirectory.questdiagnostics.com/test/test-detail/17333/sureswab-vaginosisvaginitis-plus?p=r&q=Sureswab&cc=MASTER>

Roth, A. M., Williams, J. A., Ly, R., Curd, K., Brooks, D., Arno, J., & Van Der Pol, B. (2011). Changing sexually transmitted infection screening protocol will result in improved case finding for trichomonas vaginalis among high-risk female populations. *Sex Transm Dis*, 38(5), 398-400.  
doi:10.1097/OLQ.0b013e318203e3ce

Schwebke, J. R., Gaydos, C. A., Nyirjesy, P., Paradis, S., Kodosi, S., & Cooper, C. K. (2018). Diagnostic Performance of a Molecular Test versus Clinician Assessment of Vaginitis. *J Clin Microbiol*, 56(6). doi:10.1128/jcm.00252-18

Sherrard, J. (2019). Evaluation of the BD MAX Vaginal Panel for the detection of vaginal infections in a sexual health service in the UK. *Int J STD AIDS*, 30(4), 411-414.  
doi:10.1177/0956462418815284

Sobel, J. D. (1985). Epidemiology and pathogenesis of recurrent vulvovaginal candidiasis. *Am J Obstet Gynecol*, 152(7 Pt 2), 924-935. Retrieved from <http://dx.doi.org/>

Sobel, J. D. (1999). Vulvovaginitis in healthy women. *Compr Ther*, 25(6-7), 335-346. Retrieved from <http://dx.doi.org/>

Sobel, J. D. (2020a, 4/8/2020). Approach to women with symptoms of vaginitis - UpToDate. *UpToDate*. Retrieved from [https://www.uptodate.com/contents/approach-to-women-with-symptoms-of-vaginitis?source=search\\_result&search=bacterial%20vaginosis&selectedTitle=2~97](https://www.uptodate.com/contents/approach-to-women-with-symptoms-of-vaginitis?source=search_result&search=bacterial%20vaginosis&selectedTitle=2~97)

Sobel, J. D. (2020b, 9/4/2020). Bacterial vaginosis: Clinical manifestations and diagnoses. *UpToDate*. Retrieved from <https://www.uptodate.com/contents/bacterial-vaginosis-clinical-manifestations-and-diagnosis>

Sobel, J. D. (2020c, 3/24/2020). Candida vulvovaginitis: Clinical manifestations and diagnosis. *UpToDate*. Retrieved from [https://www.uptodate.com/contents/candida-vulvovaginitis?source=see\\_link#H6](https://www.uptodate.com/contents/candida-vulvovaginitis?source=see_link#H6)

Sobel, J. D., & Mitchell, C. (2020, 12/1/5/2020). Trichomoniasis. *UpToDate*. Retrieved from [https://www.uptodate.com/contents/trichomoniasis?source=see\\_link&sectionName=Rapid%20anti-gen%20and%20DNA%20hybridization%20probes&anchor=H10#H10](https://www.uptodate.com/contents/trichomoniasis?source=see_link&sectionName=Rapid%20anti-gen%20and%20DNA%20hybridization%20probes&anchor=H10#H10)

Sobel, J. D., Subramanian, C., Foxman, B., Fairfax, M., & Gygax, S. E. (2013). Mixed vaginitis—more than coinfection and with therapeutic implications. *Curr Infect Dis Rep*, 15(2), 104-108.  
doi:10.1007/s11908-013-0325-5

Spiegel, C. A. (1991). Bacterial vaginosis. *Clin Microbiol Rev*, 4(4), 485-502. Retrieved from <http://dx.doi.org/>

# Diagnosis of Vaginitis including Multi-target PCR Testing AHS – M2057

Sumeksri, P., Kopraser, C., & Panichkul, S. (2005). BVBLUE test for diagnosis of bacterial vaginosis in pregnant women attending antenatal care at Phramongkutklo Hospital. *J Med Assoc Thai*, 88 Suppl 3, S7-13.

Tabrizi, S. N., Pirotta, M. V., Rudland, E., & Garland, S. M. (2006). Detection of *Candida* species by PCR in self-collected vaginal swabs of women after taking antibiotics. In *Mycoses* (Vol. 49, pp. 523-524). Germany.

Ubiome. (2017). SmartJane Physician's Guidelines Clinical Approach and Recommendations. Retrieved from <http://s3-us-west-1.amazonaws.com/ubiome-assets/wp-content/uploads/2017/12/06202128/SmartJane-Physician-Guidelines.pdf>

Ubiome. (2018). SmartGut. Retrieved from <https://ubiome.com/providers/smartgut/>

USPSTF. (2008). Screening for bacterial vaginosis in pregnancy to prevent preterm delivery: U.S. Preventive Services Task Force recommendation statement. *Ann Intern Med*, 148(3), 214-219. Retrieved from <http://dx.doi.org/>

van Schalkwyk, J., & Yudin, M. H. (2015). Vulvovaginitis: screening for and management of trichomoniasis, vulvovaginal candidiasis, and bacterial vaginosis. *J Obstet Gynaecol Can*, 37(3), 266-274. doi:10.1016/s1701-2163(15)30316-9

Weissenbacher, T., Witkin, S. S., Ledger, W. J., Tolbert, V., Gingelmaier, A., Scholz, C., . . . Mylonas, I. (2009). Relationship between clinical diagnosis of recurrent vulvovaginal candidiasis and detection of *Candida* species by culture and polymerase chain reaction. *Arch Gynecol Obstet*, 279(2), 125-129. doi:10.1007/s00404-008-0681-9

Workowski, K. A., & Bolan, G. A. (2015). Sexually transmitted diseases treatment guidelines, 2015. *MMWR Recomm Rep*, 64(Rr-03), 1-137. Retrieved from <http://dx.doi.org/>

Yudin, M. H., & Money, D. M. (2017). No. 211-Screening and Management of Bacterial Vaginosis in Pregnancy. *J Obstet Gynaecol Can*, 39(8), e184-e191. doi:10.1016/j.jogc.2017.04.018

Specialty Matched Consultant Advisory Panel 4/2020

Medical Director review 4/2021



# Diagnosis of Vaginitis including Multi-target PCR Testing AHS – M2057

## Policy Implementation/Update Information

---

- 1/1/19 New policy developed. BCBSNC will provide coverage for diagnosis of vaginitis including multi-target PCR testing when it is determined to be medically necessary because the medical criteria and guidelines are met. Medical Director review 1/1/2019. Policy noticed 1/1/2019 for effective date 4/1/2019. (an)
- 5/14/19 Reviewed by Avalon 1<sup>st</sup> Quarter 2019 CAB. “Using molecular-based panel testing, including, but not limited to testing such as SmartJane™, to test for microorganisms involved in vaginal flora imbalance and/or infertility is considered investigational” was added to list of NonCovered indications. Policy Guidelines updated. Codes 87905 and 0068U added to Billing/Coding section. References added. Medical Director review 4/2019. **Policy noticed 5/14/2019 for effective date 7/16/2019.** (an)
- 10/1/19 Policy statement revised to read: BCBSNC will provide coverage for diagnosis of vaginitis including multi-target PCR testing when it is determined the medical criteria or reimbursement guidelines noted below are met. Wording revised in When Covered section. “Medically Necessary” changed to “Reimbursement is allowed...” Wording revised in the Not Covered section. “Not Medically Necessary” and “investigational” changed to read “Reimbursement is not allowed...” Deleted coding grid. (an)
- 5/12/20 Reviewed by Avalon 1<sup>st</sup> Quarter 2020 CAB. Medical Director review 4/2020. Code Q0111 added to Billing/Coding section. Description, Policy Guidelines, and Reference section updated. “Reimbursement is allowed for rapid identification of Trichomonas by enzyme immunoassay in patients with symptoms of vaginitis.” Added to When covered section. “Specialty Matched Consultant Advisory Panel review 4/30/2020. (eel)
- 11/10/20 Reviewed by Avalon 3<sup>rd</sup> Quarter 2020 CAB. Medical Director review 10/2020. Code 0068U removed from Billing/Coding section. (bb)
- 3/31/21 Specialty Matched Consultant Advisory Panel 3/9/21. No change to policy. (bb)
- 5/4/21 Reviewed by Avalon 1<sup>st</sup> Quarter 2021 CAB. Description, Policy Guidelines, and References updated. Codes 81513 and 81514 added to Billing/Coding section. Medical Director review 4/2021. (bb)

---

Medical policy is not an authorization, certification, explanation of benefits or a contract. Benefits and eligibility are determined before medical guidelines and payment guidelines are applied. Benefits are determined by the group contract and subscriber certificate that is in effect at the time services are rendered. This document is solely provided for informational purposes only and is based on research of current medical literature and review of common medical practices in the treatment and diagnosis of disease. Medical practices and knowledge are constantly changing and BCBSNC reserves the right to review and revise its medical policies periodically.