

## Corporate Medical Policy

### Gene Expression Profiling and Protein Biomarkers for Prostate Cancer AHS - M2166

**File Name:** gene\_expression\_profiling\_and\_protein\_biomarkers\_for\_prostate\_cancer  
**Origination:** 9/2019  
**Last Review:** 5/2022

#### **Description of Procedure or Service**

---

Prostate cancer is characterized by malignancy which originates in the small walnut-shaped gland in men that produces the seminal fluid. Heterogeneous in both molecular alterations and progression, clinical course ranges from a microscopic tumor that never becomes clinically significant to aggressive disease that can cause metastases, morbidity, and death (Benedettini et al., 2008; Taplin, & Smith, 2022).

Gene expression assays quantify specific mRNAs being transcribed to assess the genes that are active in a particular cell or tissue. Analyses of gene expression can be clinically useful for disease classification, diagnosis, prognosis, and tailoring treatment to underlying genetic determinants of pharmacologic response (Steiling & Christenson, 2021). Protein expression-based assays measure the expression of the translation end-product(s) to assess cell-cycle progression. Similar to gene expression assays, protein biomarker-based assays can be clinically useful for disease classification and possible surveillance (Blume-Jensen et al., 2015; Ross et al., 2021).

#### **Related Policies:**

Prostate Biopsies AHS-G2007  
Prostate Specific Antigen (PSA) Testing AHS-G2008  
Serum Tumor Markers for Malignancies AHS-G2124

***\*\*\*Note: This Medical Policy is complex and technical. For questions concerning the technical language and/or specific clinical indications for its use, please consult your physician.***

#### **Policy**

---

**Gene expression profiling and protein biomarkers for prostate cancer are considered investigational for all applications. BCBSNC does not provide coverage for investigational services or procedures.**

#### **Benefits Application**

---

This medical policy relates only to the services or supplies described herein. Please refer to the Member's Benefit Booklet for availability of benefits. Member's benefits may vary according to benefit design; therefore member benefit language should be reviewed before applying the terms of this medical policy.

#### **When Gene Expression Profiling and Protein Biomarkers for Prostate Cancer is covered**

---

1. The one-time use of Prolaris®, Oncotype DX®, OR Decipher® tumor-based assays to guide management of prostate cancer is considered medically necessary in individuals with low-risk or favorable intermediate-risk disease, as defined by the NCCN (see Note 1), only if **ALL** of the following criteria have been met:

# Gene Expression Profiling and Protein Biomarkers for Prostate Cancer

## AHS - M2166

- a. Needle biopsy with localized adenocarcinoma of prostate with no clinical evidence of metastasis or lymph node involvement; **AND**
  - b. No presence of significant co-morbidities, including advanced age, to suggest individual has an estimated life expectancy of less than 10 years
2. The one-time use of Prolaris® or Decipher® tumor-based assays to guide management of prostate cancer is considered medically necessary in individuals with unfavorable intermediate-risk and high-risk disease, as defined by the NCCN (see Note 1), only if **ALL** of the following criteria have been met:
- a. Needle biopsy with localized adenocarcinoma of prostate with no clinical evidence of metastasis or lymph node involvement; **AND**
  - b. No presence of significant co-morbidities, including advanced age, to suggest individual has an estimated life expectancy of less than 10 years
3. The one-time use of the EPI biomarker test prior to initial biopsy is considered medically necessary in individuals with a prostate only if **ALL** of the following criteria have been met:
- a. Moderately elevated PSA levels
    - i. For individuals ages 50 – 75 years, PSA levels between 3 – 10 ng/mL
    - ii. For individuals over the age of 75, PSA levels between 4 – 10 ng/mL
  - b. No other relative indication for prostate biopsy including **ANY** of the following:
    - i. DRE suspicious for cancer
    - ii. Persistently elevated PSA
    - iii. Positive multiparametric MRI, if performed
    - iv. Ethnicity at higher risk for prostate cancer (See Note 2)
    - v. First-degree relative with prostate cancer
    - vi. Known to have a high-penetrance prostate cancer risk gene(s) per NCCN guidelines (See Note
  - c. No other relative contraindication for prostate biopsy including **ANY** of the following:
    - i. <10-year life expectancy
    - ii. Benign disease not ruled out

### **When Gene Expression Profiling and Protein Biomarkers for Prostate Cancer is not covered**

---

1. The following tests to assess and/or monitor prostate cancer are considered not medically necessary:
  - a. Ki-67 immunohistochemistry
  - b. *PTEN* loss
2. The following tests are considered not medically necessary, including, but not limited to:
  - All other urine testing for gene expression profile and/or protein biomarkers to assess prostate cancer, including Select MDx.
  - Other screening tests for prostate cancer, including, but not limited to, alpha-methylacyl coenzyme A racemase (AMACR), early prostate cancer antigen, endoglin, E twenty-six (ETS) gene fusions, human kallikrein 2, analysis of prostatic fluid electrolyte composition, interleukin-6, transforming growth factor-beta 1, TMPRSS2:ERG gene fusion, MyProstateScore, gene hypermethylation, *PCA3/CLK3* (prostate cancer antigen 3

# Gene Expression Profiling and Protein Biomarkers for Prostate Cancer

## AHS - M2166

[non-protein coding]/kallikrein-related peptidase 3 [prostate-specific antigen] ratio, 4Kscore, Prostate Health Index (PHI), PCA3 score or ConfirmMDx.

- Other tests using cellular and biologic features of a tumor, including use in predicting risk of recurrence in patients with prostate cancer.

Reimbursement is not allowed for other screening tests for prostate cancer that are not listed in the when covered section, including, but not limited to, alpha-methylacyl coenzyme A racemase (AMACR).

NOTE 1: NCCN Prostate Cancer Initial Risk Stratification and Staging Workup for Clinically Localized Disease (NCCN, 2022).

Risk Group	Clinical/Pathological Features		
Very Low	Has all of the following: <ul style="list-style-type: none"> <li>· cT1c; AND</li> <li>· Grade Group 1</li> <li>· PSA &lt;10 ng/mL</li> <li>· Fewer than 3 prostate biopsy fragments/cores positive, ≤50% cancer in each fragment/core</li> <li>· PSA density &lt;0.15 ng/mL/g</li> </ul>		
Low	Has all of the following but does not qualify for very low risk: <ul style="list-style-type: none"> <li>· cT1-cT2a</li> <li>· Grade Group 1</li> <li>· PSA &lt;10 ng/mL</li> </ul>		
Intermediate	Has all of the following: <ul style="list-style-type: none"> <li>· No high-risk group features</li> <li>· No very-high-risk group features</li> <li>· Has one or more intermediate risk factors               <ul style="list-style-type: none"> <li>» cT2b-cT2c</li> <li>» Grade Group 2 or 3</li> <li>» PSA 10-20 ng/mL</li> </ul> </li> </ul>	Favorable Intermediate	Has all of the following: <ul style="list-style-type: none"> <li>· 1 IRF</li> <li>· Grade Group 1 or 2</li> <li>· &lt;50% biopsy cores positive</li> </ul>
		Unfavorable Intermediate	Has one or more of the following: <ul style="list-style-type: none"> <li>· 2 or 3 IRFs</li> <li>· Grade Group 3</li> <li>· ≥50% biopsy cores positive</li> </ul>
High	Has no very-high-risk features and has at least one high-risk feature: <ul style="list-style-type: none"> <li>· cT3a OR</li> <li>· Grade Group 4 or Grade Group 5 OR</li> <li>· PSA &gt;20 ng/mL</li> </ul>		
Very High	Has at least one of the following: <ul style="list-style-type: none"> <li>· T3b-T4</li> <li>· Primary Gleason pattern 5</li> <li>· 2 or 3 high-risk features</li> <li>· &gt;4 cores with Grade Group 4 or 5</li> </ul>		

# Gene Expression Profiling and Protein Biomarkers for Prostate Cancer

## AHS - M2166

NOTE 2: According to the NCCN Prostate Cancer Early Detection guidelines, “African-American men, men with a family history of prostate cancer, and those with a known genetic predisposition represent high-risk groups (NCCN, 2021).”

NOTE 3: According to the NCCN Prostate Cancer Early Detection guidelines, the main high-penetrance cancer risk genes include *BRCA1*, *BRCA2*, *ATM*, *MLH1*, *MSH2*, *MSH6*, *HOXB13*, *CHEK2*, *NBN*, *PALB2*, *RAD51D*, and *TP53* (NCCN, 2021).

### Policy Guidelines

---

Prostate cancer (PCa) is the most common cancer in American men and the second leading cause of death in men over 65 (Balducci et al., 1997; Tabayoyong & Abouassaly, 2015). In 2021, the American Cancer Society estimates that approximately 248,530 new prostate cancer diagnoses and approximately 34,130 prostate cancer deaths will occur; although, the 5-year survival rate between 2007-2013 was 99%. About 1 man in 8 will be diagnosed with prostate cancer during his lifetime in the United States (ACS, 2021; Siegel et al., 2018).

Many cases of prostate cancer do not become clinically evident, as indicated in autopsy studies, where prostate cancer is detected in approximately 30 percent of men aged 55 or older and approximately 60 percent of men by age 80 (Bell, Del Mar, Wright, Dickinson, & Glasziou, 2015). These data suggest that prostate cancer often grows so slowly that most men die of other causes before the disease becomes clinically advanced (Hoffman, 2021).

Prostate cancer survival is related to many factors, especially the extent of tumor at the time of diagnosis. The five-year relative survival among men with cancer localized to the prostate or with regional spread is 100%, compared with 31% among those diagnosed with distant metastases (Hoffman, 2021). Gene expression profiling has been proposed as a method of risk stratification for prostate cancer. Several tests evaluating the expression levels of various genes have been produced to be used in conjunction with other tools such as Gleason score and PSA assessment.

Tissue-based gene expression classifiers (GEC) are now widely used to assist in prostate cancer prognosis. These tests are RNA-based prognostic biomarkers that analyze a distinct multigene panel to predict cancer progression, from the chance of having the disease to the probability of death at ten years due to prostate cancer. Genomic tests can predict prostate cancer aggressiveness, detect potentially dangerous prostate cancer-related genomic activity, and utilize biopsy samples to deliver prognostic information via immunofluorescence imaging. Additionally, researchers have identified the potential of microRNAs as human prostate cancer biomarkers (Song et al., 2018). While several types of biomarker tests exist, the NCCN specifically recommends Prolaris, Oncotype DX Prostate, Decipher, and ProMark as tumor-based molecular assays to consider during initial risk stratification (NCCN, 2022). Ki-67 and PTEN are also listed in NCCN guidelines, but are not recommended (NCCN, 2022).

#### *Proprietary Testing, Clinical Utility and Analytical Validity*

Hu et al. (2018) evaluated the utility of three genomic expression classifiers (GEC), including Decipher, Oncotype, and Prolaris. 747 patients underwent GEC testing. The authors found that “Among patients with clinical favorable risk of cancer, the rate of active surveillance (AS) differed significantly among patients with a GEC result above the threshold (46.2%), those with a GEC result below the threshold (75.9%), and those who did not undergo GEC (57.9%)”. The authors further estimated that “for every nine men with favorable risk of cancer who undergo GEC testing, one additional patient may have their disease initially managed with AS (Hu et al., 2018).”

#### *Prolaris*

The test “Prolaris” (created by Myriad Genetics) has been used to inform decision making on active surveillance (AS) and whether to proceed to a treatment option, such as radiation or surgery. Prolaris is an assessment of the average expression of 31 cell-cycle progression (CCP) genes compared to 15 reference genes. This score is combined with the patient’s age, PSA, percent positive cores, clinical stage, Gleason score, and AUA risk category and is intended to provide a 10-year prostate cancer-specific mortality risk. Scores range from 0 to 10, with each unit increase representing a doubling of disease-risk progression. Prolaris may also be used to assess risk post-

# Gene Expression Profiling and Protein Biomarkers for Prostate Cancer

## AHS - M2166

prostatectomy, and the same scale of 0-10 is used. Each unit increase represents a doubling of risk of biochemical recurrence (BCR) (Alford et al., 2017).

CCP expression has found to correlate with mortality rate of prostate cancer. Cuzick et al. (2015) found that not only was there a relationship between CCP expression and mortality rate, the increased expression of CCP was predictive of BCR after 10 years. Even after adjusting for factors such as PSA and Gleason score, the CCP was both “highly significant” and “independent” of prostate cancer mortality rate. The authors noted that the CCP score could be created from minimal tumor mass (as little as 0.5 mm), with a 90% success rate with >0.5 mm visible tumor, as well as Prolaris’ objective criteria compared to the Gleason score (Cuzick et al., 2015).

Prolaris may be used to lower unnecessary treatment by providing a molecular indication of the disease’s progression. Radical treatments, such as prostatectomies, are often unnecessary, and there is utility in a biomarker metric than can reliably inform providers of a course of treatment or condition. An AS status is preferable to treatment. Hu et al. (2018) used data provided by the CCP score (along with two other biomarker tests) to perform risk stratification and assess whether further treatment was needed or if the condition could be managed by active surveillance. Lin et al. (2018) clearly separated high- and low-risk patients using the CCP score. The study combined the CCP score as well as a clinical assessment from CAPRA into a cell-cycle risk (CCR) score. This CCR score was used to select patients for an AS status. The threshold created from both the molecular measures and the clinical measures has the advantage of including higher-risk patients whose clinical features may be lower-risk. Furthermore, the patients that fell below the threshold were found to have a mortality risk of 2.5%, and the probability of survival of patients with scores under the threshold was 100% (Hu et al., 2018; Lin et al., 2018). Finally, Prolaris has been used by providers to inform clinician decision making. A survey by Carneiro et al found that the course of treatment for prostate cancer patients was influenced by Prolaris’ results. About 65% of cases were reported to have shifted in the intended treatment based on the test results, and about 40% were reported to have opted for the AS choice (a “decrease” in treatment) (Carneiro et al., 2018).

Tward et al. (2020) studied the ability of CCR to predict prostate cancer metastasis using Prolaris. According to a CCR threshold of 2.112, 29.5% patients were hypothesized to be high risk metastasis (CCR>2.112) and 70.5% were unfavorable intermediate risk patients (CCR < 2.112). Patients were followed five years later to determine if CCR accurately predicted metastasis in men undergoing multimodality therapy (androgen deprivation with surgery) or radiation therapy. According to the results, the CCR score does provide a clinically meaningful different risk of metastasis for patients receiving multimodality therapy or radiation therapy. Multimodality therapy reduced patients' risk of metastasis and treatment benefit can be evaluated as a function of the CCR score. For those with CCR scores below the threshold of 2.112 (27% of high-risk group and 73% of the unfavorable intermediate group), radiation therapy was considered after assessing the difference in the risk of metastasis (Tward et al., 2020).

### *Oncotype DX*

Oncotype DX is similar to Prolaris in that it assesses levels of gene expression, could be used for lower-risk patients, and could inform clinicians about the possible course of treatment. The primary difference is that Oncotype DX only tests 12 genes, with 5 reference genes (compared to 31 and 15, respectively, for Prolaris). These expression levels are combined into an algorithm to produce a GPS score of 0-100. This GPS score correlated with prediction of cancer aggression (outcomes such as death or recurrence) (Cullen et al., 2015).

Cullen et al. (2015) found that the GPS score correlated well with BCR. The researchers noted that OncoType DX is a good predictor of both early and late BCR and is validated for adverse pathology whereas Prolaris is validated for 10-year mortality or BCR after radical prostatectomy (Alford et al., 2017; Cullen et al., 2015; Davis, 2014; NCCN, 2022). Oncotype DX was recently validated in a group of men separated by race, showing that this tool is an independent predictor of adverse pathology with similar predictive accuracy in both African American (n=96) and European American (n=76) men (Murphy et al., 2020).

### *OncoType DX AR-V7 Nucleus*

# Gene Expression Profiling and Protein Biomarkers for Prostate Cancer

## AHS - M2166

This test evaluates the AR-V7 protein in the nucleus of circulating tumor cells and is intended to identify metastatic castration-resistant prostate cancer patients who will not respond to androgen-receptor targeted therapies (OncoType, 2019, 2021).

Scher et al. (2016) examined 161 patients with progressive metastatic castration-resistant prostate cancer (mCRPC) to assess its association with AR-V7. Out of 191 samples (128 pre-ARS inhibitor and 63 pretaxane), the investigators found AR-V7-positive circulating tumor cells in 34 samples, and those samples were found to have worse clinical outcomes and overall survival than those without AR-V7. Scher et al. (2016) concluded that “the results validate CTC nuclear expression of AR-V7 protein in men with mCRPC as a treatment-specific biomarker that is associated with superior survival on taxane therapy over ARS-directed therapy in a clinical practice setting (Scher et al., 2016).”

Further, Chen et al. (2018) studied the overexpression of the nuclear AR-V7 protein in prostate cancer cases. A total of 401 men participated in this study. Participants were split into two cohorts: cohort I included those who were high-risk (n=238), and cohort II included those who were not considered high-risk (n=238). Analyses showed that high nuclear AR-V7 protein expression was detected in approximately 30-40% of participants, and a “High baseline expression of nuclear AR-V7 protein was associated with an unfavorable BCR-free survival in the high-risk patient cohort I but not in the unselected consecutive cohort II. Remarkably, AR-V7 was an independent negative prognostic factor in high-risk prostate cancer patients of cohort I who were selected to receive adjuvant treatment (Chen et al., 2018).”

Graf et al. (2020) studied the clinical utility of AR-V7 as a biomarker for patients with progressing metastatic castration-resistant prostate cancer (mCRPC). The results were used by physicians to make a second line of therapy choice of either an androgen receptor signaling inhibitor (ARSI) or taxane chemotherapy. 255 samples of circulating tumor cells (CTCs) were tested for AR-V7. Patients with detectable AR-V7 in the CTCs had superior survival with taxane treatment over ARSIs and patients who were AR-V7- negative had superior survival on ARSIs over taxanes. These results showed that men who tested AR-V7- positive were more likely to survive longer on taxane chemotherapy. Overall, the authors suggest that the use of AR-V7 CTC test “to inform treatment choice can improve patient outcomes relative to decisions based solely on standard-of-care measures (Graf et al., 2020).”

### *Decipher*

Decipher is a genomic prognostic test that is used to predict cancer outcomes in patients that have undergone a radical prostatectomy (RP). It relies on the expression levels of 22 RNA markers in the RP specimen and is primarily used to predict likeliness of metastases or mortality. The algorithm score ranges from 0 to 1, where a higher score corresponds with higher chance of metastasis. This algorithm was shown to have outperformed the traditional assessment of clinical and pathological features in predicting metastasis (0.75 accuracy compared to 0.69) as well as 17 other genetic tests (0.54 to 0.68 accuracy) (Alford et al., 2017; Dalela, et al., 2016).

Van den Broeck et al. (2019) aimed to validate the Decipher test in the prediction of distant metastatic recurrence in men with high-risk nonmetastatic prostate cancer 10 years after the surgery was completed. A total of 298 men participated in this study. Results showed that “the median Decipher scores were higher in the population that developed metastases” suggesting that this study “validates Decipher as a predictor for metastatic recurrence even in patients with high-risk, nonmetastatic PC [prostate cancer] within 10-yr follow-up (Van den Broeck et al., 2019).” Specifically, the data showed that each 10% increase in Decipher score resulted in an increased risk of distant metastatic prostate cancer recurrence.

In a prospective trial by Marascio et al. (2020), the clinical utility of the Decipher tumor test on postoperative management of prostate cancer post prostatectomy was discussed. 3,455 males were enrolled in the study and the change in treatment decision-making was recorded. In the cohort, 61% of the patients had high-risk tumors with a two-year prostate cancer reoccurrence. As a result of genome classifier testing, providers’ recommendations changed for 39% of the patients, translating to a number needed to test of three to change one treatment decision. This study demonstrated that genome classifier testing favorably impacts treatment decision making post radical

# Gene Expression Profiling and Protein Biomarkers for Prostate Cancer

## AHS - M2166

prostatectomy, promoting more post-operative radiotherapy. This translated to improved patient reported quality of life (Marascio et al., 2020).

### *ExoDX Prostate (IntelliScore)*

ExoDX is a urinary test that detects the expression level of three genetic biomarkers (ERG, PCA3, and SPDEF) (AMA, 2019; ExoSome, 2019, 2021). This test integrates the expression levels of these three biomarkers and assigns an individualized risk score to predict the risk of high-grade prostate cancer (Gleason score  $\geq 7$ ). This test is intended for men 50 or over with a PSA level of 2-10 ng/mL presenting for an initial biopsy (prior to a DRE) (ExoSome, 2019, 2021).

McKiernan et al. (2016) used ExoDX to discriminate between benign prostate cancer (Gleason score 6 and under) and high-risk cancer (Gleason score  $\geq 7$ ). The prognostic score was derived from a sample of 499 patients with PSA levels of 2-20 ng/mL; it was then validated in a sample of 1064 patients and evaluated in a population of 255. The test was compared to the standard of care practices (SOC), and the area under the curve (AUC) of the test was 0.77 compared to the SOC's 0.66. An independent validation found the AUC of the test to be 0.73 compared to the SOC's 0.63. The authors calculated that 138 of 519 biopsies (27%) would have been avoided and that the test only missed 5% of patients with high-risk disease (McKiernan et al., 2016). Within a second phase of the long-term study, McKiernan and colleagues report that using the EPI validated cut-point of 15.6 results in avoiding 26% of unnecessary prostate biopsies and a 20% decrease in all biopsies. If the EPI cut-point is raised to 20, then 31% of total biopsies would be avoided, including 40% of unnecessary biopsies (McKiernan et al., 2018).

A study published in 2018 did a cost-effectiveness analysis and comparison of not only ExoDx (EPI), but also Prostate Health Index (PHI), 4Kscore, and SelectMDx to current standard care of care. Using 2017 US dollars for their calculations, the cost and quality adjusted life-years (QALY) for the current standard of care—transrectal ultrasound guided biopsy (TRUS biopsy)—was \$3,863 and 18.0865, respectively. The authors of the study note that EPI, PHI, and SelectMDx cost less than performing TRUS biopsy. They note, “The EPI provided the highest QALY with an incremental cost-effectiveness ratio of \$58,404 per QALY. The use of biomarkers could reduce the number of unnecessary biopsies by 24% to 34% compared to the current standard of care... Using SelectMDx or the EPI following elevated prostate specific antigen but before proceeding to biopsy is a cost-effective strategy in this setting (Sathianathen Niranjana et al., 2018).”

A randomized, blinded, two-armed clinical utility study was published in 2020 using ExoDx (EPI) in individuals presenting for initial biopsy with PSA values in the intermediate range (2 – 10 ng/mL). This large study (n = 1,094) included 72 urologists from 24 different practices. All patients had an EPI test performed, but the patients were divided into two different groups (control and experimental) where only the experimental group received results prior to their biopsy decision. Of the individuals within the experimental group who received negative EPI scores, 74% deferred biopsy. For individuals within the experimental group who received positive EPI scores, 87% were recommended by their urologists to undergo the biopsy, and ultimately 72% did. As compared to the control arm of the study, there is a 30% increase in the detection of high-grade prostate cancer [HGPC], and the authors “estimate that 49% fewer HGPC were missed due to deferrals compared to standard of care (SOC). Overall, 68% of urologists reported that the EPI test influenced their biopsy decision (Tutrone et al., 2020).”

### *ConfirmMDX*

ConfirmMDX uses methylation-specific polymerase chain reaction (PCR) to identify whether a patient with a previously negative prostate biopsy should undergo a repeat biopsy. This test identifies methylation of three genes (*GSTPI*, *APC*, and *RASSF1*) (MDx, 2018a; MDx\_Health, 2020a). This test has been evaluated by Van Neste, Partin, et al. (2016) and was found to have a negative predictive value (NPV) of 96% for high-grade prostate cancer. A total of 7899 prostate core biopsies from 803 patients were assessed, and the NPV of finding low levels of DNA methylation was 89.2% for all cancers. The positive predictive value (PPV) of the genetic assay was found to be 28.2% (for detection of any cancer on a repeat biopsy), and this was calculated to be “significantly higher” than the PPV of standard of care practices. The final algorithm was optimized to a maximum of 0.742 AUC (Van

# Gene Expression Profiling and Protein Biomarkers for Prostate Cancer

## AHS - M2166

Neste, Partin, et al., 2016). Wojno et al. (2014) evaluated the utility of this test and found that out of 138 patients that the test had been performed on, only 6 with a negative result had undergone a repeat biopsy.

### *SelectMDX*

SelectMDX evaluates two mRNA cancer-related biomarkers (HOXC6 and DLX1 with *KLK3* as a reference gene) to assist a clinician in deciding to continue routine screening or to order a prostate biopsy. This test is considered a “non-invasive urine test” and reports a binary result of “increased risk” or “very low risk” (MDx, 2018b; MDx\_Health, 2020b). Van Neste, Hendriks, et al. (2016) evaluated this test at a 0.90 AUC in a validation cohort. The authors concluded that the mRNA signature was one of the most significant components of the validation results (Van Neste, Hendriks, et al., 2016). Shore (2018) assessed the effect of SelectMDX results on clinical decision making and found that out of 253 patients that SelectMDX evaluated as “negative,” only 12% underwent a biopsy (Shore, 2018).

### *Proveri*

Proveri evaluates 114 diagnostic biomarkers and 15 prognostic biomarkers to assess prostate cancer. The diagnostic biomarkers were validated with 364 samples (243 tumor samples, 121 control) and assessed at a sensitivity of 88% and specificity of 98% (97% accuracy). The prognostic biomarkers were validated with 49 samples (40 relapse patients, 9 with indolent disease) and were calculated to have a sensitivity of 88%, a specificity of 80% and were 87% accurate (Proveri, 2013). This test has since been discontinued.

### *Progensis PCA3*

Progensis *PCA3* is an FDA-approved assay that examines the concentration of the prostate cancer gene 3 (*PCA3*) and compares it to the amount of prostate-specific antigen RNA. This test is intended for assistance in decision making for a repeat biopsy in men 50 years or older, and a *PCA3* score under 25 was associated with a decreased likelihood of a positive biopsy. However, the manufacturer states this test should not be used for men with atypical small acinar proliferation on their most recent biopsy (Hologic, 2017). A total of 466 samples were provided, and 102 of these samples were evaluated to require a repeat biopsy. This assay was evaluated at a 77.5% sensitivity, a 57.1% specificity, a 33.6% positive predictive value, and a 90.0% negative predictive value (Gittelman et al., 2013).

Rodríguez and García-Perdomo (2020) performed a systematic review and meta-analysis of the diagnostic accuracy of *PCA3* prior to a patient’s first prostate biopsy. They found that with a cutoff of 35, the sensitivity of the diagnostic tests was 0.69 (95% confidence interval 0.61-0.75), specificity was 0.65 (95% confidence interval 0.553-0.733), the diagnostic odds ratio was 4.244 (95% confidence interval 3.487-5.166), and the AUC was 0.734 (95% confidence interval 0.674-0.805). This study suggests that there may be a greater clinical utility with 35 as the cutoff as opposed to the 25 approved by the FDA, and ultimately urinary *PCA3* can “be used as a guide for directing the performance of the first prostate biopsy and decreasing unnecessary biopsies” (Matuszczak et al., 2021; Rodríguez & García-Perdomo, 2020).

### *ProMark*

Another test that may have utility is ProMark. It measures the levels of eight proteins through quantitative immunofluorescence of a biopsy specimen. It is used to predict cancer aggression in patients with a Gleason score of 3+3 or 3+4. The proteins chosen have roles in cell proliferation, signaling, or stress response, and the score is reported from 1-100. This score represents individualized risk. Blume-Jensen et al. (2015) narrowed down the 8 primary protein biomarkers used (down from the 12 proposed by an earlier study) as well as assessed its ability to predict clinical endpoints of favorable and nonfavorable disease. They recommended a cutoff of 0.33 (on a scale of 0-1) for “nonfavorable” pathology (83.6% of patients with favorable disease fell below this cutoff). Conversely, a cutoff of 0.8 was recommended for favorable pathology as 76.9% of patients with nonfavorable pathology were

# Gene Expression Profiling and Protein Biomarkers for Prostate Cancer

## AHS - M2166

above this cutoff. The authors concluded that this assay provided useful information, especially when differentiating between Gleason scores (Alford et al., 2017; Blume-Jensen et al., 2015).

### *4kScore*

4kScore, from OPKO, is intended to assess the risk for “aggressive” prostate cancer. The test incorporates total PSA, free PSA, “intact” PSA, and “hk2” [human kallikrein 2] (NCCN, 2022; OPKO, 2021). These biomarkers, along with other patient clinical information (such as age and prior biopsy status) are evaluated by the OPKO 4kScore Algorithm. Finally, the test provides a risk assessment for aggressive cancer (% risk of Gleason 7 or higher, if a biopsy were to be performed).

Zappala et al. (2017) performed a meta-analysis of 4kScore validation studies. A total of 12 studies encompassing 11134 patients were included, and the pooled area under curve (AUC) for the test to “discriminate for high-grade PCa [prostate cancer] was found to be 0.81 (Zappala, Scardino, Okrongly, Linder, & Dong, 2017).

Wysock et al., (2020) compared the performance of 4K score to SelectMDx in detecting prostate cancer in 114 patients who received both tests. These tests were analyzed to provide guidance on whether to perform biopsy. Based on the results, the two scores lead to different biopsy recommendations. 50 of the 144 patients underwent biopsy based on the test results. 22 of the 50 patients (44%) were found to have clinically significant prostate cancer. In addition, the specificity of 4K score was significantly greater compared to SelectMDx while sensitivity was similar. The area under the curve for 4K score was 0.830 and SelectMDx was 0.672. The authors state that "the 4Kscore when combined with magnetic resonance imaging was superior to the SelectMDx" in detecting prostate cancer (Wysock et al., 2020).

### *Prostate Health Index*

Prostate Health Index (PHI) measures total PSA, fPSA (free non-protein bound PSA), and p2PSA (an isoform of fPSA). Levels of these three proteins are combined and calculated, implying that men with a higher total PSA and p2PSA and a lower fPSA have a higher risk of presenting with prostate cancer (Couñago et al., 2020). PHI is clinically used to reduce the number of unnecessary biopsies in men with border-line PSA levels, predict biochemical recurrence after radical prostatectomy, and enhance the predictive value of multi-parametric MRI. PHI is not recommended in primary screening for prostate cancer (Duffy, 2020).

Jia et al. (2020) compared the diagnostic value of PCA3 and PHI for detection of prostate cancer at initial biopsy in a meta-analysis of 10,376 patients from 20 studies. The pooled sensitivity for PCA3 and PHI was 0.55 and 0.88, respectively. The pooled specificity for PCA3 and PHI was 0.74 and 0.36. The area under the curve, measuring overall quality of the diagnostic test, was 0.72 for PCA3 and 0.76 for PHI. The combination use of PCA3 and PHI resulted in a higher area under the curve of 0.79. Overall, this study suggests that both PCA3 and PHI show acceptable results and a "combination of these two diagnostic tests may be more helpful than the use of either test alone in prostate cancer management (Jia et al., 2020)."

White et al. (2018) evaluated the clinical utility of the PHI on “biopsy decision management” among patients with “non-suspicious DRE findings and tPSA in the 4-10 ng/mL range” in an observational study at several large urology group practices. They found that there was a “significant reduction in biopsy procedures performed” in men receiving a PHI test when comparing to the control group (36.4% biopsy vs 60.3% biopsy), and that the “PHI score impacted physician’s patient management plan in 73% of cases, including biopsy deferrals when the PHI score was low, and decisions to perform biopsies when the PHI score indicated an intermediate or high probability of prostate cancer,” defined as a score greater than or equal to 36. This altogether conveyed the importance of the PHI score in clinical decision making in terms of how to proceed with individual patient circumstances (Matuszczak et al., 2021; White et al., 2018).

### *Ki-67 and PTEN*

# Gene Expression Profiling and Protein Biomarkers for Prostate Cancer

## AHS - M2166

Finally, the NCCN specifically recommends *against* two particular tests in assessment of prostate cancer; Ki-67 staining and PTEN loss (NCCN, 2022).

Ki-67 is a nuclear protein involved in cell cycle proliferation and is intended to provide prognostic information on metastasis and prostate cancer-specific mortality (NCCN, 2022; Ross et al., 2021). Ki-67 staining has shown some promising results. However, the primary limitation with these studies is that most active surveillance populations will have a Gleason Score of 6 or less, which is considered “low-risk”. This population will most likely have low Ki-67 levels, clouding its utility in populations trying to decide between immediate and deferred treatment (Ross et al., 2021).

PTEN loss is a relatively early event in the course of prostate cancer. PTEN is a tumor suppressor gene on chromosome 10q and is involved in cell cycle regulation. PTEN is intended to provide prognostic information on prostate cancer-specific mortality, biochemical recurrence, and cancer progression (NCCN, 2022; Ross et al., 2021). Data on prognostic value of PTEN loss post-treatment have been conflicting. It is possible that active treatments contribute to the disruption of the PTEN pathway or the high correlation between PTEN loss and clinicopathologic factors. Lotan et al found that when clinicopathologic factors, such as Gleason Score and surgical margin status, were included in their multivariable analysis, PTEN’s association with metastasis and prostate cancer-specific mortality decreased significantly (Lotan et al., 2011).

### **National Cancer Coalition Network (NCCN)**

Patients with low or favorable intermediate-risk disease and life expectancy  $\geq 10$  y may consider the use of the following tumor-based molecular assays: Decipher, Oncotype DX Prostate, and Prolaris. Patients with unfavorable intermediate- and high-risk disease and life expectancy  $\geq 10$  y may consider the use of Decipher and Prolaris tumor-based molecular assays. Retrospective studies have shown that molecular assays performed on prostate biopsy or RP specimens provide prognostic information independent of NCCN or CAPRA risk groups. These include, but are not limited to, likelihood of death with conservative management, likelihood of biochemical progression after RP or EBRT [external beam radiation therapy], and likelihood of developing metastasis after RP or salvage radiotherapy (NCCN, 2022).” Furthermore, they note that clinicians may consider testing patients with metastatic prostate cancer and regional prostate cancer for alterations in homologous recombination DNA repair genes *BRCA1*, *BRCA2*, *ATM*, *PALB2*, *FANCA*, *RAD51D*, *CHEK2*, and *CDK12*; “Post-test genetic counseling is recommended if pathogenic/likely pathogenic somatic mutations in any gene that has clinical implications if also identified in germline (eg, *BRCA2*, *BRCA1*, *ATM*, *PALB2*, *CHEK2*, *MLH1*, *MSH2*, *MSH6*, *PMS2*) (NCCN, 2022).” The NCCN noted that somatic tumor testing of the aforementioned genes has potential for early use of platinum chemotherapy, use of PARP inhibitors, or eligibility for clinical trials. Lastly, they recommend that men with regional disease, metastatic castration-resistant disease, or castration-naïve metastatic disease should additionally consider tumor testing for microsatellite instability or mismatch repair deficiency. The NCCN also specifically does not recommend either Ki-67 or *PTEN* testing (NCCN, 2022).

The NCCN does include available tissue-based tests for prostate cancer prognosis within their table of possible testing as indicated in the Table below. Regarding Decipher testing, NCCN states that Decipher “may be considered to inform adjuvant treatment if adverse events are found to occur after radical prostatectomy”. Decipher testing can also be used to inform counseling for risk stratification in patients with PSA resistance or reoccurrence after radical prostatectomy. NCCN discourages repeat molecular tumor analysis (NCCN, 2021, 2022):

# Gene Expression Profiling and Protein Biomarkers for Prostate Cancer

## AHS - M2166

Test	Platform	Recommendation
<b>Decipher</b>	Whole-Transcriptome 1.4M RNA expression (46,050 genes), oligonucleotide microarray optimized for FFPE tissue	Cover post-biopsy for NCCN very-low- and low-risk prostate cancer in patients with at least 10 years life expectancy who have not yet received treatment for prostate cancer and are candidates for active surveillance or definitive therapy.  Cover post-RP for pT2 with positive margins, any pT3 disease, or rising PSA (above nadir)
<b>KI-67</b>	IHC	Not recommended
<b>Oncotype DX</b>	Quantitative RT-PCR for 12 prostate cancer-related genes and 5 housekeeping controls	Cover post-biopsy for NCCN very-low- and low-risk prostate cancer in patients with at least 10 years life expectancy who have not yet received treatment for prostate cancer and are candidates for active surveillance or definitive therapy.
<b>Prolaris</b>	Quantitative RT-PCR for 31 prostate cell cycle-related and 15 housekeeping controls	Cover post-biopsy for NCCN very-low- and low-risk prostate cancer in patients with at least 10 years life expectancy who have not yet received treatment for prostate cancer and are candidates for active surveillance or definitive therapy.
<b>PTEN</b>	Fluorescence in situ hybridization or IHC	Not recommended

The NCCN, within the algorithm for the indications for prostate biopsy, says to “consider biomarkers that improve the specificity of screening” for individuals who have had elevated levels of PSA (above 3 ng/mL for those ages 45 – 75 years or 4 ng/mL or higher for those individuals over the age of 75 years. The NCCN goes on to state, “Biomarkers that improve the specificity of detection are not, as yet, mandated as first-line screening tests in conjunction with serum PSA. However, there may be some patients who meet PSA standards for consideration of prostate biopsy, but for whom the patient and/or the physician wish to further define risk. Percent-free may improve cancer detection. The probability of high-grade cancer (Gleason score > 3+4, Grade Group 2, or higher) may be further defined utilizing the SelectMDx, 4Kscore, Prostate Health Index (PHI), and ExoDx Prostate test. Extent of validation of these tests across diverse populations is variable. It is not known how such tests could be applied in the optimal combination with MRI as yet (NCCN, 2021).” The NCCN notes that these tests- including percent-free PSA, 4Kscore, PHI, PCA3, and ConfirmMDx-improve specificity in the post-biopsy setting and it should be considered in patients who are thought to be at higher risk despite a negative prostate biopsy (NCCN, 2021).

The NCCN panel remarks that 4Kscore “can be considered for patients prior to biopsy and for those with a negative biopsy who are thought to be at higher risk for clinically significant prostate cancer” The NCCN further remarks that SelectMDx is “potentially informative” in patients who have never undergone biopsy, and can therefore be “considered” in these patients. The NCCN also acknowledged that ConfirmMDX and PCA3 can be considered an option for men contemplating repeat biopsy and is approved for limited coverage by MoIDX to reduce unnecessary repeat biopsies. Further, ExoDx *Prostate (IntelliScore)*, also called EPI, “can be considered as an option for men contemplating initial or repeat biopsy (NCCN, 2021).”

# Gene Expression Profiling and Protein Biomarkers for Prostate Cancer

## AHS - M2166

### **American Association of Clinical Urologists Inc.**

The AACU recommends use of tissue-based molecular testing to assess risk stratification in prostate cancer treatment decision making. The AACU states pursuing germline testing when appropriate is encouraged and support any further research into these tests. The AACU specifically recommends, “Tissue-based molecular testing should be considered for low and favorable intermediate risk men with life expectancy  $\geq 10$  years.” **The Large Urology Group Practice Association (LUGPA)** endorses this position statement by the AACU (AACU, 2018; LUGPA, 2018).

### **American Society of Clinical Oncology (ASCO)**

The ASCO released a guideline stating that they endorsed the non-cryotherapy 2017 joint guidelines from the American Urological Association (AUA)/American Society for Radiation Oncology (ASTRO)/Society of Urologic Oncology (SUO) (Bekelman et al., 2018).

Guideline 32 stated “tissue based genomic biomarkers have not shown a clear role in active surveillance for localized prostate cancer and are not necessary for follow up.” These joint guidelines also state that several genomic assays were validated in the pre-MRI era and that their clinical utility “remains to be established” (Sanda et al., 2018).

The AUA has also noted a “lack of predictive biomarkers to help better personalize therapy” in drug development for prostate cancer patients (AUA, 2018).

In 2019, an ASCO multidisciplinary panel published guidelines on molecular biomarkers in localized prostate cancer. These guidelines are below:

- “Commercially available molecular biomarkers (ie, Oncotype Dx Prostate, Prolaris, Decipher, and ProMark) may be offered in situations in which the assay result, when considered as a whole with routine clinical factors, is likely to affect management. Routine ordering of molecular biomarkers is not recommended
- Any additional molecular biomarkers evaluated do not have sufficient data to be clinically actionable or are not commercially available and thus should not be offered
- The Expert Panel recommends consideration of a commercially available molecular biomarker (eg, Decipher Genomic Classifier) in situations in which the assay result, when considered as a whole with routine clinical factors, is likely to affect management. In the absence of prospective clinical trial data, routine use of genomic biomarkers in the postprostatectomy setting to determine adjuvant versus salvage radiation or to initiate systemic therapies should not be offered (S. E. Eggener et al., 2019).”

In 2020, an ASCO panel published guidelines on the use of molecular biomarkers in localized prostate cancer. In concordance with the 2018 and 2019, ASCO recommends the use of commercially available tests (Oncotype Dx Prostate, Prolaris, Decipher, and ProMark) when the assay result “is likely to have an impact on patient management. Examples include select men with high-volume low-risk or favorable intermediate-risk prostate cancer who are considering active surveillance or in men with high-risk features for treatment intensification. While testing may influence management decisions, there is no high-level evidence that the results from these panels will improve quality of life or cancer-specific outcomes (Scott E. Eggener et al., 2020).

### **European Association of Urology (EAU), European Society for Radiotherapy and Oncology (ESTRO), European Society of Urogenital Radiology (ESUR), and the International Society of Geriatric Oncology (SIOG)**

The EAU, ESTRO, ESUR and SIOG released joint guidelines on prostate cancer. These guidelines stated that “To avoid unnecessary biopsies, offer further risk-assessment to asymptomatic men with a normal digital rectal examination (DRE) and a prostate-specific antigen (PSA) level between 2-10 ng/mL prior to performing a

# Gene Expression Profiling and Protein Biomarkers for Prostate Cancer AHS - M2166

prostate biopsy. Use one of the following tools:

- risk-calculator (Strong);
- imaging (Strong);
- an additional serum or urine-based test (Weak).”

These joint guidelines acknowledged both SelectMDX and ConfirmMDX as tests to select for repeat biopsies, but the guidelines noted SelectMDX as having an “uncertain role” and “probably not cost-effective”. No recommendation could be made for ConfirmMDX. Prolaris and OncoType DX were also recognized as tests that have been used to evaluate prostate cancer, but no recommendation could be made at this time (EAU, 2021).

In 2021, updated joint guidelines acknowledged five commercially available tests (Oncotype Dx, Prolaris, Decipher, Decipher PORTOS and ProMark). Since the long-term impact of the use of these tests is unproven, the panel concluded that “these tests should not be offered routinely, but only in subsets of patients where the test result provides clinically actionable information, such as for instance in men with favourable intermediate-risk PCa who might opt for AS or men with unfavourable intermediate-risk PCa at RP to decide on treatment intensification with hormonal therapy (HT) (EAU, 2021).”

## **Public Health England (PHE)**

PHE notes PCA3 as a “promising urinary RNA biomarker” (PHE, 2016). The source guideline for this information was withdrawn on July 27, 2021.

## **European Society for Medical Oncology (ESMO)**

ESMO provided recommendations on the use of precision medicine in providing prognostic information for prostate cancer. These are the following recommendations provided:

- ESMO does not recommend the use of AR-V7 testing, stating that the test is of limited value in therapy selection.
- Other tissue-based molecular assays may be used on conjunction with clinicopathological factors to make treatment decision.
- Germline testing for *BRCA2* and other DDR [DNA damage and repair] genes is recommended in patients with a family history of cancer and should be considered in patients with metastatic cancer (Parker et al., 2020).

## **State and Federal Regulations, as applicable**

### **Food and Drug Administration (FDA)**

The FDA has approved 3 tests for evaluation of gene expression profiles of prostate cancer as of February 1, 2021 (FDA, 2021).

On November 6, 2020, the FDA approved FoundationOne CDx, by Foundation Medicine, Inc. This device is a next generation sequencing based in vitro diagnostic device for detection of substitutions, insertion and deletion alterations (indels), and copy number alterations (CNAs) in 311 genes, rearrangements in 3 genes, and copy number alterations in 3 genes. FoundationOne CDx also utilizes circulating cell-free DNA collected in FoundationOne® Liquid CDx Blood Sample Collection Kit to identify patients with non-small cell lung cancer, prostate cancer, ovarian cancer, or breast cancer who may benefit from treatment with the targeted therapies. This test provides tumor mutation profiling of *BRCA1*, *BRCA2*, and ATM alterations for prostate cancer diagnosis (FDA, 2020).

On February 13, 2012, the FDA approved the PROGNSA PCA3 Assay created by Gen-Probe Inc. From the FDA website: “The PROGNSA PCA3 Assay is an in vitro nucleic acid amplification test. The assay measures the concentration of prostate cancer gene 3 (PCA3) and prostate-specific antigen (PSA) RNA molecules and calculates the ratio of PCA3 RNA molecules to PSA RNA molecules (PCA3 Score) in post-digital rectal exam (DRE) first catch male urine specimens. The PROGNSA PCA3 Assay is indicated for use in conjunction with other

# Gene Expression Profiling and Protein Biomarkers for Prostate Cancer AHS - M2166

patient information to aid in the decision for repeat biopsy in men 50 years of age or older who have had one or more previous negative prostate biopsies and for whom a repeat biopsy would be recommended by a urologist based on current standard of care, before consideration of PROGENSA PCA3 Assay results” (FDA, 2012).

On December 19, 2014, the FDA approved the BRACAnalysis CDx™ created by Myriad Genetics. From the FDA website: BRACAnalysis CDx™ is an in vitro diagnostic device intended for the qualitative detection and classification of variants in the protein coding regions and intron/exon boundaries of the BRCA1 and BRCA2 genes using genomic DNA obtained from whole blood specimens collected in EDTA. Single nucleotide variants and small insertions and deletions (indels) are identified by polymerase chain reaction (PCR) and Sanger sequencing. Large deletions and duplications in BRCA1 and BRCA2 are detected using multiplex PCR. Results of the test are used as an aid in identifying ovarian cancer patients with deleterious or suspected deleterious germline BRCA variants eligible for treatment with Lynparza™ (olaparib). This assay is for professional use only and is to be performed only at Myriad Genetic Laboratories, a single laboratory site located at 320 Wakara Way, Salt Lake City, UT 84108” (FDA, 2014) This test is commonly known as Prolaris.

Many labs have developed specific tests that they must validate and perform in house. These laboratory-developed tests (LDTs) are regulated by the Centers for Medicare and Medicaid (CMS) as high-complexity tests under the Clinical Laboratory Improvement Amendments of 1988 (CLIA '88). As an LDT, the U. S. Food and Drug Administration has not approved or cleared this test; however, FDA clearance or approval is not currently required for clinical use.

## **Billing/Coding/Physician Documentation Information**

---

This policy may apply to the following codes. Inclusion of a code in this section does not guarantee that it will be reimbursed. For further information on reimbursement guidelines, please see Administrative Policies on the Blue Cross Blue Shield of North Carolina web site at [www.bcbsnc.com](http://www.bcbsnc.com). They are listed in the Category Search on the Medical Policy search page.

*Applicable service codes: 81313, 81479, 81539, 81541, 81542, 81551, 0005U, 0021U, 0047U, 0053U, 0228U*

BCBSNC may request medical records for determination of medical necessity. When medical records are requested, letters of support and/or explanation are often useful, but are not sufficient documentation unless all specific information needed to make a medical necessity determination is included.

## **Scientific Background and Reference Sources**

---

### **For Policy Titled: Gene Expression Profiling and Protein Biomarkers for Prostate Cancer Management**

AACU. (2018). Retrieved from [https://aacuweb.org/docs/position-statements/ps\\_genomic-testing-in-prostate-cancer.aspx](https://aacuweb.org/docs/position-statements/ps_genomic-testing-in-prostate-cancer.aspx)

ACS. (2021). Cancer Facts & Figures 2020. Retrieved from <https://www.cancer.org/content/dam/cancer-org/research/cancer-facts-and-statistics/annual-cancer-facts-and-figures/2021/cancer-facts-and-figures-2021.pdf>

Alford, A. V., Brito, J. M., Yadav, K. K., Yadav, S. S., Tewari, A. K., & Renzulli, J. (2017). The Use of Biomarkers in Prostate Cancer Screening and Treatment. *Rev Urol*, 19(4), 221-234.  
doi:10.3909/riu0772

# Gene Expression Profiling and Protein Biomarkers for Prostate Cancer AHS - M2166

- AMA. (2019). CPT® Proprietary Laboratory Analyses (PLA) Codes: Long Descriptors. Retrieved from [https://www.ama-assn.org/system/files/2018-11/cpt-pla-codes-long\\_0.pdf](https://www.ama-assn.org/system/files/2018-11/cpt-pla-codes-long_0.pdf)
- AUA. (2018). Castration-Resistant Prostate Cancer (2018).
- Balducci, L., Pow-Sang, J., Friedland, J., & Diaz, J. I. (1997). Prostate cancer. *Clin Geriatr Med*, 13(2), 283-306.
- Bekelman, J. E., Rumble, R. B., Chen, R. C., Pisansky, T. M., Finelli, A., Feifer, A., . . . Freedland, S. J. (2018). Clinically Localized Prostate Cancer: ASCO Clinical Practice Guideline Endorsement of an American Urological Association/American Society for Radiation Oncology/Society of Urologic Oncology Guideline. *Journal of Clinical Oncology*, 36(32), 3251-3258. doi:10.1200/JCO.18.00606
- Bell, K. J., Del Mar, C., Wright, G., Dickinson, J., & Glasziou, P. (2015). Prevalence of incidental prostate cancer: A systematic review of autopsy studies. *Int J Cancer*, 137(7), 1749-1757. doi:10.1002/ijc.29538
- Benedettini, E., Nguyen, P., & Loda, M. (2008). The pathogenesis of prostate cancer: from molecular to metabolic alterations. *Diagn Histopathol (Oxf)*, 14(5), 195-201. doi:10.1016/j.mpdhp.2008.03.001
- Blume-Jensen, P., Berman, D. M., Rimm, D. L., Shipitsin, M., Putzi, M., Nifong, T. P., . . . Saad, F. (2015). Development and clinical validation of an in situ biopsy-based multimarker assay for risk stratification in prostate cancer. *Clin Cancer Res*, 21(11), 2591-2600. doi:10.1158/1078-0432.Ccr-14-2603
- Carneiro, A., Priante Kayano, P., Gomes Barbosa, Á. R., Langer Wroclawski, M., Ko Chen, C., Cavlini, G. C., . . . Bianco, B. (2018). Are localized prostate cancer biomarkers useful in the clinical practice? *Tumor Biology*, 40(9), 1010428318799255. doi:10.1177/1010428318799255
- Chen, X., Bernemann, C., Tolkach, Y., Heller, M., Nientiedt, C., Falkenstein, M., . . . Duensing, S. (2018). Overexpression of nuclear AR-V7 protein in primary prostate cancer is an independent negative prognostic marker in men with high-risk disease receiving adjuvant therapy. *Urol Oncol*, 36(4), 161.e119-161.e130. doi:10.1016/j.urolonc.2017.11.001
- Couñago, F., López-Campos, F., Díaz-Gavela, A. A., Almagro, E., Fenández-Pascual, E., Henríquez, I., . . . Del Cerro, E. (2020). Clinical Applications of Molecular Biomarkers in Prostate Cancer. *Cancers (Basel)*, 12(6). doi:10.3390/cancers12061550
- Cullen, J., Rosner, I. L., Brand, T. C., Zhang, N., Tsiatis, A. C., Moncur, J., . . . McLeod, D. G. (2015). A Biopsy-based 17-gene Genomic Prostate Score Predicts Recurrence After Radical Prostatectomy and Adverse Surgical Pathology in a Racially Diverse Population of Men with Clinically Low- and Intermediate-risk Prostate Cancer. *Eur Urol*, 68(1), 123-131. doi:10.1016/j.eururo.2014.11.030
- Cuzick, J., Stone, S., Fisher, G., Yang, Z. H., North, B. V., Berney, D. M., . . . Scardino, P. (2015). Validation of an RNA cell cycle progression score for predicting death from prostate cancer in a conservatively managed needle biopsy cohort. *Br J Cancer*, 113(3), 382-389. doi:10.1038/bjc.2015.223
- Dalela, D., Löppenber, B., Sood, A., Sammon, J., & Abdollah, F. (2016). Contemporary Role of the Decipher® Test in Prostate Cancer Management: Current Practice and Future Perspectives. *Rev Urol*, 18(1), 1-9.
- Davis, J. W. (2014). Novel commercially available genomic tests for prostate cancer: a roadmap to understanding their clinical impact. *BJU International*, 114(3), 320-322. doi:10.1111/bju.12695
- Duffy, M. J. (2020). Biomarkers for prostate cancer: prostate-specific antigen and beyond. *Clin Chem Lab Med*, 58(3), 326-339. doi:10.1515/cclm-2019-0693

# Gene Expression Profiling and Protein Biomarkers for Prostate Cancer AHS - M2166

EAU. (2021). EAU Guidelines. Edn. presented at the EAU Annual Congress Amsterdam 2020. Retrieved from <https://uroweb.org/guideline/prostate-cancer/#11>

Eggener, S. E., Rumble, R. B., Armstrong, A. J., Morgan, T. M., Crispino, T., Cornford, P., . . . Beltran, H. (2019). Molecular Biomarkers in Localized Prostate Cancer: ASCO Guideline. *J Clin Oncol*, Jco1902768. doi:10.1200/jco.19.02768

Eggener, S. E., Rumble, R. B., & Beltran, H. (2020). Molecular Biomarkers in Localized Prostate Cancer: ASCO Guideline Summary. *JCO Oncology Practice*, 16(6), 340-343. doi:10.1200/jop.19.00752

ExoSome. (2019). Prostate Cancer. Retrieved from <http://www.exosomedx.com/prostate-cancer-0>

ExoSome. (2021). ExoDx™ Prostate Test. Retrieved from <http://www.exosomedx.com/physicians/exodx-prostate-test>

FDA. (2012). PROGENSA PCA3 ASSAY. *Summary of Safety and Effectiveness Data (SSED)*. Retrieved from [https://www.accessdata.fda.gov/cdrh\\_docs/pdf10/P100033B.pdf](https://www.accessdata.fda.gov/cdrh_docs/pdf10/P100033B.pdf)

FDA. (2014). BRACANALYSIS CDX. *Summary of Safety and Effectiveness Data (SSED)*. Retrieved from [https://www.accessdata.fda.gov/cdrh\\_docs/pdf14/P140020B.pdf](https://www.accessdata.fda.gov/cdrh_docs/pdf14/P140020B.pdf) FDA. (2019).

FDA. (2020). Retrieved from <https://www.accessdata.fda.gov/scripts/cdrh/devicesatfda/index.cfm>

FDA. (2021). Retrieved from <https://www.accessdata.fda.gov/scripts/cdrh/devicesatfda/index.cfm>

Gittelman, M. C., Hertzman, B., Bailen, J., Williams, T., Koziol, I., Henderson, R. J., . . . Ward, J. F. (2013). PCA3 molecular urine test as a predictor of repeat prostate biopsy outcome in men with previous negative biopsies: a prospective multicenter clinical study. *J Urol*, 190(1), 64-69. doi:10.1016/j.juro.2013.02.018

Graf, R. P., Hullings, M., Barnett, E. S., Carbone, E., Dittamore, R., & Scher, H. I. (2020). Clinical Utility of the Nuclear-localized AR-V7 Biomarker in Circulating Tumor Cells in Improving Physician Treatment Choice in Castration-resistant Prostate Cancer. *Eur Urol*, 77(2), 170-177. doi:<https://doi.org/10.1016/j.eururo.2019.08.020>

Hoffman, R. (2020). Screening for prostate cancer - UpToDate. Retrieved from [https://www.uptodate.com/contents/screening-for-prostate-cancer?source=see\\_link#H30](https://www.uptodate.com/contents/screening-for-prostate-cancer?source=see_link#H30). [https://www.uptodate.com/contents/screening-for-prostate-cancer?source=see\\_link#H30](https://www.uptodate.com/contents/screening-for-prostate-cancer?source=see_link#H30)

Hologic. (2017). LEARN ABOUT PCA3. Retrieved from <http://www.pca3.org/learn-about-pca3>

Hu, J. C., Tosoian, J. J., Qi, J., Kaye, D., Johnson, A., Linsell, S., . . . Morgan, T. M. (2018). Clinical Utility of Gene Expression Classifiers in Men With Newly Diagnosed Prostate Cancer. (2), 1-15. doi:10.1200/po.18.00163

Jia, W., Wu, B., Shao, Y., Cao, X., & Wang, D. (2020). Diagnostic performance of prostate cancer antigen 3 and the Prostate Health Index in detecting overall and clinically significant prostate cancer in men at first biopsy: A meta-analysis. *International Journal of Urology*, n/a(n/a). doi:<https://doi.org/10.1111/iju.14464>

Lin, D. W., Crawford, E. D., Keane, T., Evans, B., Reid, J., Rajamani, S., . . . Cuzick, J. (2018). Identification of men with low-risk biopsy-confirmed prostate cancer as candidates for active surveillance. *Urologic*

# Gene Expression Profiling and Protein Biomarkers for Prostate Cancer AHS - M2166

*Oncology: Seminars and Original Investigations*, 36(6), 310.e317-310.e313.  
doi:<https://doi.org/10.1016/j.urolonc.2018.03.011>

Lotan, T. L., Gurel, B., Sutcliffe, S., Esopi, D., Liu, W., Xu, J., . . . De Marzo, A. M. (2011). PTEN protein loss by immunostaining: analytic validation and prognostic indicator for a high risk surgical cohort of prostate cancer patients. *Clin Cancer Res*, 17(20), 6563-6573. doi:10.1158/1078-0432.Ccr-11-1244

LUGPA. (2018). Retrieved from <https://lugpa.org/lugpa-endorses-and-supports-new-cancer-guidelines-from-nccn/>

Marascio, J., Spratt, D. E., Zhang, J., Trabulsi, E. J., Le, T., Sedzorme, W. S., . . . Den, R. B. (2020). Prospective study to define the clinical utility and benefit of Decipher testing in men following prostatectomy. *Prostate Cancer and Prostatic Diseases*, 23(2), 295-302. doi:10.1038/s41391-019-0185-7

Matuszczak, M., Schalken, J. A., & Salagierski, M. (2021). Prostate Cancer Liquid Biopsy Biomarkers' Clinical Utility in Diagnosis and Prognosis. *Cancers (Basel)*, 13(13). <https://doi.org/10.3390/cancers13133373>

McKiernan, J., Donovan, M. J., Margolis, E., Partin, A., Carter, B., Brown, G., . . . Carroll, P. (2018). A Prospective Adaptive Utility Trial to Validate Performance of a Novel Urine Exosome Gene Expression Assay to Predict High-grade Prostate Cancer in Patients with Prostate-specific Antigen 2-10ng/ml at Initial Biopsy. *Eur Urol*, 74(6), 731-738. doi:10.1016/j.eururo.2018.08.019

McKiernan, J., Donovan, M. J., O'Neill, V., Bentink, S., Noerholm, M., Belzer, S., . . . Carroll, P. (2016). A Novel Urine Exosome Gene Expression Assay to Predict High-grade Prostate Cancer at Initial Biopsy Urine Exosome Signature to Predict High-Grade Prostate Cancer Urine Exosome Signature to Predict High-Grade Prostate Cancer. *JAMA Oncology*, 2(7), 882-889. doi:10.1001/jamaoncol.2016.0097

MDx. (2018a). ConfirmMDx for Prostate Cancer.

MDx. (2018b). SelectMDx for Prostate Cancer.

MDx\_Health. (2020a). ConfirmMDx for Prostate Cancer. Retrieved from <https://mdxhealth.com/confirmmdx-physician/#overview>

MDx\_Health. (2020b). SelectMDx for Prostate Cancer. Retrieved from <https://mdxhealth.com/selectmdx-physician/>

Murphy, A. B., Carbanaru, S., Nettey, O. S., Gornbein, C., Macias, V., Sharifi, R., . . . Gann, P. (2020). A 17- Gene Panel Genomic Prostate Score has Similar Predictive Accuracy for Adverse Pathology at Radical Prostatectomy in African American and European American Men. *Urology*. doi:10.1016/j.urology.2020.01.052

NCCN. (2021, 07/14/2021). *Prostate Cancer Early Detection Version 2.2021*. Retrieved 02/01/2021 from [https://www.nccn.org/professionals/physician\\_gls/pdf/prostate\\_detection.pdf](https://www.nccn.org/professionals/physician_gls/pdf/prostate_detection.pdf)

NCCN. (2022, 01/10/2022). *NCCN Clinical Practice Guidelines in Oncology: Prostate Cancer, Version 3.2022*. Retrieved 2/8/21 from [https://www.nccn.org/professionals/physician\\_gls/pdf/prostate.pdf](https://www.nccn.org/professionals/physician_gls/pdf/prostate.pdf)

OncoType. (2019). Leverage the predictive power of the Oncotype DX AR-V7 Nucleus Detect test. Retrieved from <https://www.oncotypeiq.com/en-US/announcements/AR-V7>

OncoType. (2021). Leverage the predictive power of the Oncotype DX AR-V7 Nucleus Detect test. Retrieved from <https://www.oncotypeiq.com/en-US/announcements/AR-V7>

# Gene Expression Profiling and Protein Biomarkers for Prostate Cancer AHS - M2166

- OPKO. (2021). WHAT IS THE 4KSCORE®? Retrieved from <https://4kscore.com/what-is-the-4kscore.html>
- Parker, C., Castro, E., Fizazi, K., Heidenreich, A., Ost, P., Procopio, G., . . . Gillessen, S. (2020). Prostate cancer: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up<sup>&#x2020;</sup>. *Annals of Oncology*, 31(9), 1119-1134. doi:10.1016/j.annonc.2020.06.011
- PHE. (2016). Prostate cancer risk management programme (PCRM): benefits and risks of PSA testing. Retrieved from <https://www.gov.uk/government/publications/prostate-cancer-risk-management-programme-psa-test-benefits-and-risks/prostate-cancer-risk-management-programme-pcrmp-benefits-and-risks-of-psa-testing#the-future-of-prostate-cancer-detection>
- Proveri. (2013). TECHNOLOGY. Retrieved from <http://www.provericorp.com/technology/>
- Rodríguez, S. V. M., & García-Perdomo, H. A. (2020). Diagnostic accuracy of prostate cancer antigen 3 (PCA3) prior to first prostate biopsy: A systematic review and meta-analysis. *Can Urol Assoc J*, 14(5), E214-e219. <https://doi.org/10.5489/cuaj.6008>
- Ross, A. E., D'Amico, A. V., & Freedland, S. J. (2021, 02/08/2021). Molecular prognostic tests for prostate cancer. Wolters Kluwer. Retrieved 04/15/2019 from <https://www.uptodate.com/contents/molecular-prognostic-tests-for-prostate-cancer>
- Sanda, M. G., Cadeddu, J. A., Kirkby, E., Chen, R. C., Crispino, T., Fontanarosa, J., . . . Treadwell, J. R. (2018). Clinically Localized Prostate Cancer: AUA/ASTRO/SUO Guideline. Part I: Risk Stratification, Shared Decision Making, and Care Options. *J Urol*, 199(3), 683-690. doi:10.1016/j.juro.2017.11.095
- Sathianathen Niranjani, J., Kuntz Karen, M., Alarid-Escudero, F., Lawrentschuk Nathan, L., Bolton Damien, M., Murphy Declan, G., . . . Konety Badrinath, R. (2018). Incorporating Biomarkers into the Primary Prostate Biopsy Setting: A Cost-Effectiveness Analysis. *Journal of Urology*, 200(6), 1215-1220. doi:10.1016/j.juro.2018.06.016
- Scher, H. I., Lu, D., Schreiber, N. A., Louw, J., Graf, R. P., Vargas, H. A., . . . Dittamore, R. (2016). Association of AR-V7 on Circulating Tumor Cells as a Treatment-Specific Biomarker With Outcomes and Survival in Castration-Resistant Prostate CancerAR-V7 Expression in Circulating Tumor Cells and Castration-Resistant Prostate Cancer OutcomesAR-V7 Expression in Circulating Tumor Cells and Castration-Resistant Prostate Cancer Outcomes. *JAMA Oncology*, 2(11), 1441-1449. doi:10.1001/jamaoncol.2016.1828
- Shore, N. (2018). SelectMDx Impacts Prostate Biopsy Decision-making in Routine Clinical Practice.
- Siegel, R. L., Miller, K. D., & Jemal, A. (2018). Cancer statistics, 2018. *CA Cancer J Clin*, 68(1), 7-30. doi:10.3322/caac.21442
- Song, C. J., Chen, H., Chen, L. Z., Ru, G. M., Guo, J. J., & Ding, Q. N. (2018). The potential of microRNAs as human prostate cancer biomarkers: A meta-analysis of related studies. *J Cell Biochem*, 119(3), 2763-2786. doi:10.1002/jcb.26445
- Steiling, K., Christenson, Stephanie. (2019). Tools for genetics and genomics: Gene expression profiling. Retrieved from [https://www.uptodate.com/contents/tools-for-genetics-and-genomics-gene-expression-profiling?source=search\\_result&search=gene%20expression%20array%20cancer&selectedTitle=2~150](https://www.uptodate.com/contents/tools-for-genetics-and-genomics-gene-expression-profiling?source=search_result&search=gene%20expression%20array%20cancer&selectedTitle=2~150)
- Tabayoyong, W., & Abouassaly, R. (2015). Prostate Cancer Screening and the Associated Controversy. *Surg Clin North Am*, 95(5), 1023-1039. doi:10.1016/j.suc.2015.05.001

# Gene Expression Profiling and Protein Biomarkers for Prostate Cancer AHS - M2166

- Taplin, M.-E., & Smith, J. A. (2022, 01/03/2022). *Clinical presentation and diagnosis of prostate cancer*. <https://www.uptodate.com/contents/clinical-presentation-and-diagnosis-of-prostate-cancer>
- Tutrone, R., Donovan, M. J., Torkler, P., Tadiogola, V., McLain, T., Noerholm, M., . . . McKiernan, J. (2020). Clinical utility of the exosome based ExoDx Prostate(IntelliScore) EPI test in men presenting for initial Biopsy with a PSA 2-10 ng/mL. *Prostate Cancer Prostatic Dis*. doi:10.1038/s41391-020-0237-z
- Tward, J. D., Schlomm, T., Bardot, S., Freedland, S. J., Lenz, L., Cohen, T., . . . Bishoff, J. (2020). Ability of the combined clinical cell-cycle risk score to identify patients that benefit from multi versus single modality therapy in NCCN intermediate and high-risk prostate cancer. *Journal of Clinical Oncology*, 38(6\_suppl), 346-346. doi:10.1200/JCO.2020.38.6\_suppl.346
- Van den Broeck, T., Moris, L., Gevaert, T., Tosco, L., Smeets, E., Fishbane, N., . . . Joniau, S. (2019). Validation of the Decipher Test for Predicting Distant Metastatic Recurrence in Men with High-risk Nonmetastatic Prostate Cancer 10 Years After Surgery. *Eur Urol Oncol*, 2(5), 589-596. doi:10.1016/j.euo.2018.12.007
- Van Neste, L., Hendriks, R. J., Dijkstra, S., Trooskens, G., Cornel, E. B., Jannink, S. A., . . . Schalken, J. A. (2016). Detection of High-grade Prostate Cancer Using a Urinary Molecular Biomarker-Based Risk Score. *Eur Urol*, 70(5), 740-748. doi:10.1016/j.eururo.2016.04.012
- Van Neste, L., Partin, A. W., Stewart, G. D., Epstein, J. I., Harrison, D. J., & Van Criekinge, W. (2016). Risk score predicts high-grade prostate cancer in DNA-methylation positive, histopathologically negative biopsies. *Prostate*, 76(12), 1078-1087. doi:10.1002/pros.23191
- White, J., Shenoy, B. V., Tutrone, R. F., Karsh, L. I., Saltzstein, D. R., Harmon, W. J., Broyles, D. L., Roddy, T. E., Lofaro, L. R., Paoli, C. J., Denham, D., & Reynolds, M. A. (2018). Clinical utility of the Prostate Health Index (phi) for biopsy decision management in a large group urology practice setting. *Prostate Cancer Prostatic Dis*, 21(1), 78-84. <https://doi.org/10.1038/s41391-017-0008-7>
- Wojno, K. J., Costa, F. J., Cornell, R. J., Small, J. D., Pasin, E., Van Criekinge, W., . . . Van Neste, L. (2014). Reduced Rate of Repeated Prostate Biopsies Observed in ConfirmMDx Clinical Utility Field Study. *Am Health Drug Benefits*, 7(3), 129-134.
- Wysock, J. S., Becher, E., Persily, J., Loeb, S., & Lepor, H. (2020). Concordance and Performance of 4Kscore and SelectMDx for Informing Decision to Perform Prostate Biopsy and Detection of Prostate Cancer. *Urology*, 141, 119-124. doi:10.1016/j.urology.2020.02.032
- Zappala, S. M., Scardino, P. T., Okrongly, D., Linder, V., & Dong, Y. (2017). Clinical performance of the 4Kscore Test to predict high-grade prostate cancer at biopsy: A meta-analysis of us and European clinical validation study results. *Rev Urol*, 19(3), 149-155. doi:10.3909/riu0776

Medical Director review 4/2020

Medical Director review 7/2020

Specialty Matched Consultant Panel 8/2020

Medical Director review 1/2021

## **For Policy Titled: Gene Expression Profiling and Protein Biomarkers for Prostate Cancer**

Medical Director review 4/2021

# Gene Expression Profiling and Protein Biomarkers for Prostate Cancer AHS - M2166

Specialty Matched Consultant Panel 8/2021

Medical Director review 4/2022

## **Policy Implementation/Update Information**

---

### **For Policy Titled: Gene Expression Profiling and Protein Biomarkers for Prostate Cancer Management**

- 10/1/19 New policy developed. Gene Expression Profiling and Protein Biomarkers for Prostate Cancer Management is considered **investigational**. See **Related Policy: Prostate Cancer Screening AHS-G2008**. Medical Director review 8/2019. (lpr)
- 10/29/19 No change to policy statements. Minor reformatting and edits. (hb)
- 5/12/20 Off cycle review to align with CMP Prostate Cancer Screening AHS-G2008. Removed references to the following tests and transferred them to AHS-G2008: ExoDX Prostate, Intelliscore, Select MDX, PCA3, KLK3, ConfirmDX, PPCA. Removed CPT codes 0005U, 81313, 81551 and transferred to AHS-G2008. Updated Policy Guidelines. Medical Director review 4/2020. (lpr)
- 7/28/20 Reviewed by Avalon 2<sup>nd</sup> Quarter 2020 CAB. References updated. Medical Director review 7/2020. (lpr)
- 9/8/20 Specialty Matched Consultant Advisory Panel review 8/19/2020. No change to policy statement. (lpr)
- 2/9/21 Added PLA code 0005U to Billing/Coding section. Off-cycle review per Avalon, No change to policy statement. (lpr)

### **For Policy Titled: Gene Expression Profiling and Protein Biomarkers for Prostate Cancer**

- 5/18/21 Reviewed by Avalon 1<sup>st</sup> Quarter 2021 CAB. Medical Director review 4/2021. Revised “When Covered” section to include tests to assess and/or monitor prostate cancer which were relocated from AHS-G2008 Prostate Specific Antigen (PSA) Testing policy. These tests remain investigational. Updated Policy Guidelines, Billing/Coding sections as well as References. Changed related policy AHS-G2008 Prostatic Specific Antigen (PSA) Testing title. **Policy Title changed from: Gene Expression Profiling and Protein Biomarkers for Prostate Cancer Management to: Gene Expression Profiling and Protein Biomarkers for Prostate Cancer**. (lpr)
- 9/7/21 Specialty Matched Consultant Advisory Panel review 8/18/2021. No change to policy statement. (lpr)
- 2/8/22 Reviewed by Avalon Q4 2021 CAB. Removed PLA code 0244U from Billing/Coding section. Medical Director review 1/2022. No change to policy statement. (lpr)
- 5/31/22 Reviewed by Avalon Q1 2022 CAB. Extensive revisions to policy. Added coverage criteria under “When Covered” section. Updated policy guidelines and references. Medical Director review 4/2022. (lpr)

---

Medical policy is not an authorization, certification, explanation of benefits or a contract. Benefits and eligibility are determined before medical guidelines and payment guidelines are applied. Benefits are determined by the group contract and subscriber certificate that is in effect at the time services are rendered. This document is solely provided for informational purposes only and is based on research of current medical literature and review of common medical practices in the treatment and diagnosis of disease.

# Gene Expression Profiling and Protein Biomarkers for Prostate Cancer

## AHS - M2166

Medical practices and knowledge are constantly changing and BCBSNC reserves the right to review and revise its medical policies periodically.