

## Corporate Medical Policy

### Genetic Testing for Alpha- and Beta- Thalassemia AHS – M2131

<b>File Name:</b>	genetic_testing_for_alpha_and_beta_thalassemia
<b>Origination:</b>	01/2019
<b>Last CAP Review:</b>	07/2020
<b>Next CAP Review:</b>	07/2021
<b>Last Review:</b>	07/2020

#### Description of Procedure or Service

---

Alpha-thalassemia is characterized by impaired production of the alpha globin chains of hemoglobin, leading to a relative excess of gamma globin chains (fetus and newborn), or excess beta globin chains (children and adults) mainly due to deletion or mutation of the alpha globin genes. There are four alpha thalassemia syndromes, reflecting the loss of function of one, two, three, or all four of these alpha chain genes varying in severity from non-symptomatic to incompatibility with extrauterine life (Benz, 2018b, 2020; Martin & Thompson, 2013).

Beta-thalassemia is similarly characterized by impaired production of hemoglobin components but affects the beta chains instead of the alpha chains. This creates excess alpha globin chains, leading to hemolytic anemia, impaired iron handling, and other clinical symptoms (Schrier, 2018d).

#### Related Policies

General Genetic Testing, Germline Disorders AHS-M2145

General Genetic Testing, Somatic Disorders AHS-M2146

***\*\*\*Note: This Medical Policy is complex and technical. For questions concerning the technical language and/or specific clinical indications for its use, please consult your physician.***

#### Policy

---

**BCBSNC will provide coverage for genetic testing for alpha- and beta- thalassemia when it is determined the medical criteria or reimbursement guidelines below are met.**

#### Benefits Application

---

This medical policy relates only to the services or supplies described herein. Please refer to the Member's Benefit Booklet for availability of benefits. Member's benefits may vary according to benefit design; therefore member benefit language should be reviewed before applying the terms of this medical policy.

#### When Genetic Testing for Alpha- and Beta- Thalassemia is covered

---

1. Reimbursement for genetic counseling for alpha- or beta-thalassemia genetic testing is allowed and is recommended.
2. Preconception (carrier) testing for alpha- or beta- thalassemia in prospective parents is considered medically necessary when either parent has evidence of possible alpha- thalassemia (including alpha thalassemia minor, hemoglobin H disease [alpha thalassemia intermedia], or alpha thalassemia major) or beta-thalassemia (including beta thalassemia minor, beta thalassemia intermedia, or beta thalassemia major) based on biochemical testing.

# Genetic Testing for Alpha- and Beta- Thalassemia AHS – M2131

3. Reimbursement is allowed for genetic testing to confirm a diagnosis of alpha- or beta-thalassemia when one of the parents is a known carrier or when other testing to diagnose cause of microcytic anemia has been inconclusive.

## **When Genetic Testing for Alpha- and Beta- Thalassemia is not covered**

---

Genetic testing for alpha- or beta- thalassemia in other clinical situations (recognizing that prenatal testing is not addressed in this policy) is investigational.

## **Policy Guidelines**

---

### **Background**

Thalassemias result from deficiencies in hemoglobin biosynthesis due to mutations in or near the two globin gene clusters which encode the globin polypeptide subunits of hemoglobin (Benz, 2018b). Normal hemoglobin is a heterotetramer of two alpha globin chains and two beta globin chains (hemoglobin A) or two gamma globin chains (hemoglobin F). Well over 100 mutations have been documented to affect the biosynthesis or post-translational stability of the globin subunits needed for successful production of the large amounts of Hb needed for normal red cell homeostasis. Globin chain synthesis is very tightly controlled, such that the ratio of production of alpha to non-alpha chains is almost exactly 1:1 (Benz, 2018c).

Alpha thalassemia refers to thalassemias that result from impaired or absent production of alpha globin, leading to a relative excess of gamma globin (fetus and newborn), or excess beta globin (children and adults). Excess beta globin chains are capable of forming soluble homotetramers, but they are nonfunctional, unstable. This may lead to increased hemolysis and a variety of clinical manifestations, such as anemia, thrombosis, and skeletal changes. A diagnosis of alpha-thalassemia is often confirmed by genetic testing, as assessment of the hemoglobin gene is inexpensive and convenient (Benz, 2018c).

The clinical severity is directly attributable to the net deficit of alpha globin synthesis but is complicated by the number of alpha globin genes affected, which of the two alpha globin loci is affected, and the degree to which the mutation blocks gene expression. In addition, combinations of defects in both alpha and beta globulins can balance each other out. Thus, understanding the broad spectrum of clinical severity in alpha thalassemia requires a detailed knowledge of the underlying genetic defect and the impact of these defects on the overall levels and balance of globin chain synthesis (Benz, 2019).

The majority of cases of alpha thalassemia are attributable to deletion of alpha globin alleles, especially in Asia and Africa (Steinberg, 1999). However, more detailed analysis of globin gene sequences suggests that some fairly common forms of alpha thalassemia that appear to arise from a deletion of one copy of an alpha globin gene are actually due to unequal crossover and recombination events that fuse the two alpha globin genes together into one (Benz, 2018c). Additionally, non-deletion alleles are also common, especially in the Mediterranean area, which contain mutations producing highly unstable alpha globin variants unable to produce intact hemoglobin (Benz, 2018b). Current research continues to identify novel mutations and improve thalassemia screening (S. He et al., 2018).

Beta-thalassemia is similar to alpha-thalassemia, with the beta chains of hemoglobin affected instead of the alpha chains. However, excess alpha globin chains do not form soluble homotetramers, causing them to aggregate when they accumulate in erythroid precursors. This causes clinical symptoms to be more severe, although the symptoms themselves are similar to alpha-thalassemia (anemia, iron overload, and so on) (Benz, 2018b; Schrier, 2017, 2018). There are two beta globin genes compared to four for the alpha chain. As with alpha-thalassemia, the severity of clinical presentation depends on the genotype of the beta globin genes (i.e. the ratio of beta to alpha globin chains). Mutations may result in a reduced expression ( $\beta^+$ ) or absent expression ( $\beta^0$ ).

# Genetic Testing for Alpha- and Beta- Thalassemia AHS – M2131

$\beta^0$  phenotypes are generally transfusion-dependent as they produce very little (if any) adult hemoglobin (Schrier, 2018d).

Due to the frequency of thalassemias worldwide, carrier screening may be useful, particularly in areas such as Southeast Asia, Africa, and the Indian subcontinent. Both primary thalassemias are autosomal recessive genetic disorders so parents who are heterozygous carriers would have a 25% chance to have an affected child despite being asymptomatic themselves. Identification of an affected fetus could alter decisions during the pregnancy (Yates, 2019).

Below is a table summarizing the clinical genotypes and phenotypes of both thalassemia syndromes (Benz, 2018a, 2018b; Steinberg, 2018) (figure from Benz).

Severity	Genotype	Anemia	Hemoglobin Analysis
<b>Alpha Thalassemias</b>			
Silent carrier	$\alpha\alpha / \alpha-$	None	Normal <3% Hb Barts (gamma globin tetramer) at birth
Minor	$\alpha\alpha / --$ or	Mild Microcytic	Normal, 3 to 8% Hb Barts at birth
	$\alpha- / \alpha-$		
Hb H disease	$\alpha- / --$	Moderate Microcytic	Up to 30% HbH (beta globin tetramer), present in adults, up to 4% HbA2 (alpha and delta globin)
Major (fetal hydrops)	$-- / --$	Severe Microcytic, usually fatal	Hb Barts, Hb Portland (zeta and gamma globin), and HbH present, Hba, HbF, and HbA2 absent
<b>Beta Thalassemias</b>			
Minor (trait or carrier)	$\beta / \beta^0$ or $\beta / \beta^+$	Mild Microcytic	HbA2 (4% or more); HbF (up to 5%)
Intermedia (non-transfusion-dependent)	$\beta^+ / \beta^+$	Moderate Microcytic	HbA2 (4% or more); HbF (up to 50%)
Major (transfusion-dependent)	$\beta^0 / \beta^0$ or $\beta^0 / \beta^+$	Severe microcytic with target cells (typical Hb 3 to 4g/dL)	HbA2 (5% or more); HbF (up to 95%); no HbA

## Clinical Validity and Utility

He et al examined a next-generation sequencing (NGS) panel's utility for thalassemia screening in Southwestern China. 951 individuals were tested, and the NGS screen found 471 carriers (49.5%)

# Genetic Testing for Alpha- and Beta- Thalassemia AHS – M2131

of thalassemia. In comparison, traditional methods (defined as “red cell indexes and hemoglobin electrophoresis, then DNA sequencing”) identified only 209 carriers (22%) of thalassemia, missing 217 alpha-thalassemia carriers and 47 beta-thalassemia carriers (J. He et al., 2017).

Nosheen et al evaluated a preliminary screening program for beta-thalassemia. The screening program focused on families of beta-thalassemia major children. 98 samples were taken, and 57 were found to have a beta-thalassemia trait with elevated hemoglobin alpha 2. The mean hemoglobin alpha 2 level of the carriers was  $5.2 \pm 0.56\%$  compared to  $2.34 \pm 0.57\%$  in normal subjects. The authors suggested that screening programs and counseling for carriers could decrease incidence of beta-thalassemia major (Nosheen et al., 2015).

Satirapod et al. (2019) evaluated the clinical outcomes of using preimplantation genetic testing (PGT) in couples at risk of passing on beta thalassemia. Two components of PGT were used, PGT for monogenic disease (used for diagnosis) and PGT for aneuploidy (intended to identify chromosomal aberrations) A total of 15 couples were included and a total of 106 embryos were tested. After preimplantation testing, 12 of 15 women were able to obtain satisfactory genetic testing results (defined as non-disease affected embryos without chromosomal aberration and transfer within first two cycles). Of these, 9 women had successful implantations and 8 women had successful pregnancies with live births (deemed a 53.33% success rate). PGT assessment of genetic status was confirmed by pre- and post-natal genetic testing. Overall, the authors concluded that combined PGT-A and PGT-M was a useful technology to prevent beta-thalassemia in the offspring of recessive carriers.

Shook, Haygood, and Quinn (2020) et al. evaluated the accuracy of a specific pattern in hemoglobin separation tests. The authors desired to find if an “FSA” pattern corresponded to a final diagnosis of the sickle cell trait (HbAS), or a final diagnosis of sickle beta-thalassemia (HbS $\beta^+$ ). Traditionally, the FSA pattern has indicated a diagnosis of HbS $\beta^+$ ; however, the authors hypothesized that the FSA pattern truly indicates a diagnosis of HbAS instead. 31 newborns with an initial screening result of the FSA pattern (a suspected diagnosis of HbS $\beta^+$ ) were included. 30 of these newborns underwent protein-base confirmatory testing and 17 underwent confirmatory genetic testing. Of the newborns undergoing protein confirmatory testing, 23 had an “FSA” pattern, establishing a diagnosis of HbAS. Of the 8 remaining newborns with an FSA pattern, 7 underwent genetic testing which identified HbAS as well. Genetic testing also confirmed positive HbAS results in 10 newborns that tested initially positive by protein testing. The authors concluded that genetic testing had utility in newborn screening for hemoglobinopathies.

## Guidelines and Recommendations

### Canadian College of Medical Geneticists (CCMG) and Society of Obstetricians and Gynaecologists of Canada (SOGC) (Wilson et al., 2016)

The CCMG and SOGC published a joint guideline titled “Joint SOGC–CCMG Opinion for Reproductive Genetic Carrier Screening: An Update for All Canadian Providers of Maternity and Reproductive Healthcare in the Era of Direct-to-Consumer Testing” in 2016. Their recommendations addressing thalassemias/hemoglobinopathies are listed below:

- "Carrier screening for hemoglobinopathies should be offered to women/families from ethnic backgrounds with a reported increased carrier frequency, when red blood cell indices reveal a mean cellular volume  $< 80$  fl, or electrophoresis reveals an abnormal hemoglobin type. However, the use of ethnicity alone in the carrier risk identification process may create screening inconsistency and missed opportunity for carrier identification, with both obstetrical and fetal implications. High clinical suspicion is

# Genetic Testing for Alpha- and Beta- Thalassemia AHS – M2131

required as well. Screening should be done in the pre-conception period or as early into the pregnancy as possible. (II-2A) (GRADE moderate/moderate)”

- “Carrier screening for thalassemia/hemoglobinopathies should be offered by the most responsible health care provider or reproductive genetic provider and include: complete blood count, hemoglobin (Hb) electrophoresis (HE) or Hb high performance liquid chromatography (HHPLC), quantification of Hb alpha 2 and fetal Hb, and serum ferritin/H bodies (blood smear stain using brilliant cresyl blue) if microcytosis (mean cellular volume < 80 fl) and/or hypochromia (mean cellular Hb < 27 pg) in the presence of a normal HE or HHPLC assessment. (II-2A) (GRADE moderate/moderate)”
- “If the female thalassemia screening results are abnormal, a hemoglobinopathy screening protocol should be undertaken for the male partner. (III-A) (GRADE low/moderate)”
- “If both reproductive partners are found to be carriers of thalassemia or a combination of thalassemia and hemoglobin variant, they should be referred for formal genetic counselling (reproductive risks, recommended prenatal testing, and diagnostic management). (II-3A) (GRADE moderate/moderate)” (Wilson et al., 2016)

## **The Thalassemia Longitudinal Cohort (Tubman et al., 2015)**

The report on the Thalassemia Longitudinal Cohort (Tubman et al., 2015) recommends: “Obtaining genotyping to confirm the diagnosis and HLA typing for transplant evaluation for all patients who require chronic transfusion is strongly recommended. For pediatric patients, annual comprehensive follow up should include assessment of the availability of a related donor as well as a recommendation to bank cord blood and obtain HLA typing on all subsequently born full siblings.”

## **American College of Obstetrics and Gynecology (ACOG) (ACOG, 2007, 2018a, 2018b)**

The ACOG Committee Opinion #691 states that: “Couples at risk of having a child with a hemoglobinopathy may benefit from genetic counseling to review their risk, the natural history of these disorders, prospects for treatment and cure, availability of prenatal genetic testing, and reproductive options. Prenatal diagnostic testing for the mutation responsible for sickle cell disease is widely available. Testing for  $\alpha$ -thalassemia and  $\beta$ -thalassemia is possible if the mutations and deletions have been previously identified in both parents. These DNA-based tests can be performed using chorionic villi obtained by chorionic villus sampling or using cultured amniotic fluid cells obtained by amniocentesis. For some couples, preimplantation genetic diagnosis in combination with in vitro fertilization may be a desirable alternative to avoid termination of an affected pregnancy. Preimplantation genetic diagnosis has been successfully performed for sickle cell disease and most types of  $\beta$ -thalassemia (ACOG, 2018).

In 2018, ACOG reaffirmed its practice bulletin regarding hemoglobinopathies in pregnancy (originally published in 2007) (ACOG, 2018b). The following recommendations are considered “Level A” and based on “good and consistent scientific evidence.”

- “Individuals of African, Southeast Asian, and Mediterranean descent are at increased risk for being carriers of hemoglobinopathies and should be offered carrier screening and, if both parents are determined to be carriers, genetic counseling.”
- “Couples at risk for having a child with sickle cell anemia or thalassemia should be offered genetic counseling to review prenatal testing and reproduction options. Prenatal diagnosis of hemoglobinopathies is best accomplished by DNA analysis of cultured amniocytes or chorionic villi.”

ACOG also writes that “if the MCV [mean corpuscular volume] is below normal, iron deficiency anemia has been excluded, and the hemoglobin electrophoresis is not consistent with  $\beta$ -thalassemia trait (ie, there is no elevation of Hb A<sub>2</sub> or Hb F), then DNA-based testing should be used to detect

# Genetic Testing for Alpha- and Beta- Thalassemia AHS – M2131

$\alpha$ -globin gene deletions characteristic of  $\alpha$ -thalassemia.” ACOG remarks that neither hemoglobin electrophoresis nor solubility testing can identify individuals with the  $\alpha$ -thalassemia trait, only molecular genetic testing can (ACOG, 2007).

## **The Association of Public Health Laboratories (APHL) (APHL, 2015)**

Molecular testing (APHL, 2015) can be added to resolve cases when the newborn has been transfused with packed red blood cells. Since the newborn’s phenotype is masked by the donor, DNA testing can be used to identify any abnormal hemoglobins.

## **National Health Service (NHS, 2017) (NHS, 2017)**

NHS states that first, the mean cell hemoglobin (MCH) must be measured in a screening for thalassemia. The NHS presents this decision tree of when to test for thalassemia:

“1. Is the MCH <25pg?

2. Is the woman’s family origin identified as high risk from the Family Origin Questionnaire: China (including Hong Kong), Southeast Asia (especially Thailand, Taiwan, Cambodia, Laos, Vietnam, Burma, Malaysia, Singapore, Indonesia or Philippines), Cyprus, Greece, Sardinia, Turkey, or unknown? If the answer to both questions is yes, testing of the baby’s biological father must be offered if he is also from a high-risk area or unknown.

If the baby’s biological father is suspected of having  $\alpha^0$  thalassaemia, the samples on both biological parents must be sent for DNA analysis for  $\alpha^0$  thalassaemia mutations.

If one biological parent is a suspected carrier of  $\alpha^0$  thalassaemia and the other is a carrier of a thalassaemia and is also from one of the high-risk groups for  $\alpha^0$  thalassaemia with an MCH < 25pg, both biological parents should be screened for  $\alpha^0$  thalassaemia by DNA analysis.”

The NHS states that 99% of  $\alpha^0$  thalassemia cases have an MCH of < 25 pg. However, the NHS acknowledges that this algorithm may miss some beta-thalassemia carriers (NHS, 2017).

## **Public Health England (PHE) (PHE, 2019)**

The PHE highlights the importance of antenatal screening. If the baby’s mother is identified as a carrier, the biological father should also be tested. Both prenatal diagnosis and genetic counselling are recommended by the PHE (PHE, 2019).

## **British Society for Haematology (BSH) (Ryan et al., 2010)**

The BSH provides the following recommendations:

- “Antenatal screening/testing of pregnant women should be carried out according to the guidelines of the NHS Sickle Cell and Thalassaemia Screening programme.
- Laboratories performing antenatal screening should utilize methods capable of detecting significant variants and be capable of quantitating haemoglobins A<sub>2</sub> and F at the cut-off points required by the national antenatal screening program (Ryan et al., 2010).”

Genetic counseling is also permitted for prospective parents.

## **Applicable Federal Regulations**

A search on the FDA website of “thalassemia” on May 17, 2020, did not yield any genetic resting results. Additionally, many labs have developed specific tests that they must validate and perform

# Genetic Testing for Alpha- and Beta- Thalassemia AHS – M2131

in house. These laboratory-developed tests (LDTs) are regulated by the Centers for Medicare and Medicaid (CMS) as high-complexity tests under the Clinical Laboratory Improvement Amendments of 1988 (CLIA '88). As an LDT, the U. S. Food and Drug Administration has not approved or cleared this test; however, FDA clearance or approval is not currently required for clinical use.

## Billing/Coding/Physician Documentation Information

---

This policy may apply to the following codes. Inclusion of a code in this section does not guarantee that it will be reimbursed. For further information on reimbursement guidelines, please see Administrative Policies on the Blue Cross Blue Shield of North Carolina web site at [www.bcbsnc.com](http://www.bcbsnc.com). They are listed in the Category Search on the Medical Policy search page.

*Applicable service codes: 81257, 81258, 81259, 81269, 81361, 81363, 81364, 96040, S3845, S3846, S0265*

BCBSNC may request medical records for determination of medical necessity. When medical records are requested, letters of support and/or explanation are often useful, but are not sufficient documentation unless all specific information needed to make a medical necessity determination is included.

## Scientific Background and Reference Sources

---

### For policy titled: Genetic Testing for Alpha Thalassemia:

- ACOG. (2018). ACOG Committee Opinion 691; Carrier Screening for Genetic Conditions - ACOG. (1074-861X). <https://www.acog.org/Clinical-Guidance-and-Publications/Committee-Opinions/Committee-on-Genetics/Carrier-Screening-for-Genetic-Conditions>
- AHPL. (2015). *Hemoglobinopathies: Current Practices for Screening, Confirmation and Follow-up*. Association of Public Health Laboratories Retrieved from [Retrieved from https://www.cdc.gov/ncbddd/sicklecell/documents/nbs\\_hemoglobinopathy-testing\\_122015.pdf](https://www.cdc.gov/ncbddd/sicklecell/documents/nbs_hemoglobinopathy-testing_122015.pdf)
- Benz, E. (2017a). Clinical manifestations and diagnosis of the thalassemias. In S. Schrier (Ed.), UpToDate. Waltham, MA. Retrieved from [https://www.uptodate.com/contents/clinical-manifestations-and-diagnosis-of-the-thalassemias?source=search\\_result&search=alpha%20thalassemia&selectedTitle=1~47#H29](https://www.uptodate.com/contents/clinical-manifestations-and-diagnosis-of-the-thalassemias?source=search_result&search=alpha%20thalassemia&selectedTitle=1~47#H29).
- Benz, E. (2017b). Molecular pathology of the thalassemic syndromes - UpToDate. In S. Schrier (Ed.), UpToDate. Waltham, MA. Retrieved from [https://www.uptodate.com/contents/molecular-pathology-of-the-thalassemic-syndromes?source=search\\_result&search=alpha%20thalassemia&selectedTitle=4~47](https://www.uptodate.com/contents/molecular-pathology-of-the-thalassemic-syndromes?source=search_result&search=alpha%20thalassemia&selectedTitle=4~47).
- Fogel, B. N., Nguyen, H. L. T., Smink, G., & Sekhar, D. L. (2018). Variability in State-Based Recommendations for Management of Alpha Thalassemia Trait and Silent Carrier Detected on the Newborn Screen. *J Pediatr*, 195, 283-287. doi:10.1016/j.jpeds.2017.11.048
- Galanello, R., & Cao, A. (2011). Gene test review. Alpha-thalassemia. *Genet Med*, 13(2), 83-88. doi:10.1097/GIM.0b013e3181fcb468
- He, S., Li, J., Li, D. M., Yi, S., Lu, X., Luo, Y., . . . Qiu, X. (2018). Molecular characterization of alpha- and beta-thalassemia in the Yulin region of Southern China. *Gene*, 655, 61-64. doi:10.1016/j.gene.2018.02.058
- Kulozik, A. E., Thein, S. L., Wainscoat, J. S., Gale, R., Kay, L. A., Wood, J. K., . . . Huehns, E. R. (1987). Thalassaemia intermedia: interaction of the triple alpha-globin gene arrangement and heterozygous beta-thalassaemia. *Br J Haematol*, 66(1), 109-112.
- Landaw, S. (2017). Lab findings thalassemias. In UpToDate. Waltham, MA. Retrieved from [https://www.uptodate.com/contents/image?imageKey=HEME%2F50393&topicKey=HEME%2F7116&rank=1~47&source=see\\_link&search=alpha%20thalassemia](https://www.uptodate.com/contents/image?imageKey=HEME%2F50393&topicKey=HEME%2F7116&rank=1~47&source=see_link&search=alpha%20thalassemia).

# Genetic Testing for Alpha- and Beta- Thalassemia AHS – M2131

- Langlois, S., Ford, J. C., Chitayat, D., Desilets, V. A., Farrell, S. A., Geraghty, M., . . . Wyatt, P. (2008). Carrier screening for thalassemia and hemoglobinopathies in Canada. *J Obstet Gynaecol Can*, 30(10), 950-971. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/19038079>.
- Martin, A., & Thompson, A. A. (2013). Thalassemias. *Pediatr Clin North Am*, 60(6), 1383-1391. Retrieved from <http://dx.doi.org/10.1016/j.pcl.2013.08.008>.doi:10.1016/j.pcl.2013.08.008
- Origa, R., & Moi, P. (2016). Alpha-Thalassemia. *GeneReviews*. doi:<https://www.ncbi.nlm.nih.gov/books/NBK1435/>
- Schrier, S. (2017). Pathophysiology of alpha thalassemia - UpToDate. In J. Tirnauer (Ed.), *UpToDate*. Waltham, MA. Retrieved from [https://www.uptodate.com/contents/pathophysiology-of-alpha-thalassemia?source=see\\_link](https://www.uptodate.com/contents/pathophysiology-of-alpha-thalassemia?source=see_link).
- Steinberg, M. H. (1999). Management of sickle cell disease. *N Engl J Med*, 340(13), 1021-1030. Retrieved from <http://www.md.rcm.upr.edu/medschool/anesthesiology/residency/24-Suggested%20Readings/GA%20READINGS/General%20Operating%20Rooms/Steinberg,%202004%20Sickle%20Cell.pdf>.doi:10.1056/nejm199904013401307
- Tubman, V. N., Fung, E. B., Vogiatzi, M., Thompson, A. A., Rogers, Z. R., Neufeld, E. J., & Kwiatkowski, J. L. (2015). Guidelines for the Standard Monitoring of Patients with Thalassemia: Report of the Thalassemia Longitudinal Cohort. *J Pediatr Hematol Oncol*, 37(3), e162-169. Retrieved from <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4511957/>.doi:10.1097/MPH.0000000000000307

Specialty Matched Consultant Advisory Panel review 7/2019

Medical Director review 7/2019

## **For policy titled: Genetic Testing for Alpha- and Beta- Thalassemia:**

- ACOG. (2007). ACOG Practice Bulletin No. 78: hemoglobinopathies in pregnancy. *Obstet Gynecol*, 109(1), 229-237. doi:10.1097/00006250-200701000-00055
- ACOG. (2018a). ACOG Committee Opinion 691; Carrier Screening for Genetic Conditions - ACOG. Retrieved from <https://www.acog.org/Clinical-Guidance-and-Publications/Committee-Opinions/Committee-on-Genetics/Carrier-Screening-for-Genetic-Conditions>. (1074-861X). <https://www.acog.org/Clinical-Guidance-and-Publications/Committee-Opinions/Committee-on-Genetics/Carrier-Screening-for-Genetic-Conditions>
- ACOG. (2018b). ACOG Publications. *Obstetrics & Gynecology*, 131(1). Retrieved from [https://journals.lww.com/greenjournal/Fulltext/2018/01000/ACOG\\_Publications.31.aspx](https://journals.lww.com/greenjournal/Fulltext/2018/01000/ACOG_Publications.31.aspx)
- APHL. (2015). *Hemoglobinopathies: Current Practices for Screening, Confirmation and Follow-up*. Association of Public Health Laboratories Retrieved from [https://www.cdc.gov/ncbddd/sicklecell/documents/nbs\\_hemoglobinopathy-testing\\_122015.pdf](https://www.cdc.gov/ncbddd/sicklecell/documents/nbs_hemoglobinopathy-testing_122015.pdf)
- Benz, E. (2018a). Classical thalassemia syndromes (genotypes and laboratory findings). In *UpToDate*. Waltham, MA.
- Benz, E. (2018b). Clinical manifestations and diagnosis of the thalassemias. In S. Schrier (Ed.), *UpToDate*. Retrieved from [https://www.uptodate.com/contents/clinical-manifestations-and-diagnosis-of-the-thalassemias?search=alpha%20thalassemia&source=search\\_result&selectedTitle=1~97&usage\\_type=default&display\\_rank=1#H2871792528](https://www.uptodate.com/contents/clinical-manifestations-and-diagnosis-of-the-thalassemias?search=alpha%20thalassemia&source=search_result&selectedTitle=1~97&usage_type=default&display_rank=1#H2871792528)
- Benz, E. (2018c). Molecular pathology of the thalassemic syndromes - UpToDate. In S. Schrier (Ed.), *UpToDate*. Retrieved from [https://www.uptodate.com/contents/molecular-pathology-of-the-thalassemic-syndromes?source=search\\_result&search=alpha%20thalassemia&selectedTitle=4~47](https://www.uptodate.com/contents/molecular-pathology-of-the-thalassemic-syndromes?source=search_result&search=alpha%20thalassemia&selectedTitle=4~47)
- Benz, E. (2018d). Pathophysiology of beta thalassemia. In J. Tirnauer (Ed.), *UpToDate*. Retrieved from [https://www.uptodate.com/contents/pathophysiology-of-beta-thalassemia?topicRef=7103&source=see\\_link](https://www.uptodate.com/contents/pathophysiology-of-beta-thalassemia?topicRef=7103&source=see_link)
- Benz, E. (2019). Pathophysiology of alpha thalassemia - UpToDate. In J. Tirnauer (Ed.), *UpToDate*. Retrieved from [https://www.uptodate.com/contents/pathophysiology-of-alpha-thalassemia?source=see\\_link](https://www.uptodate.com/contents/pathophysiology-of-alpha-thalassemia?source=see_link)



# Genetic Testing for Alpha- and Beta- Thalassemia AHS – M2131

- Benz, E. (2020). Clinical manifestations and diagnosis of the thalassemias. In S. Schrier (Ed.), *UpToDate*. Retrieved from [https://www.uptodate.com/contents/clinical-manifestations-and-diagnosis-of-the-thalassemias?search=alpha%20thalassemia&source=search\\_result&selectedTitle=1~97&usage\\_type=default&display\\_rank=1#H2871792528](https://www.uptodate.com/contents/clinical-manifestations-and-diagnosis-of-the-thalassemias?search=alpha%20thalassemia&source=search_result&selectedTitle=1~97&usage_type=default&display_rank=1#H2871792528)
- FDA. (2020). Devices@FDA. Retrieved from <https://www.accessdata.fda.gov/scripts/cdrh/devicesatfda/index.cfm>
- He, J., Song, W., Yang, J., Lu, S., Yuan, Y., Guo, J., . . . Zhu, B. (2017). Next-generation sequencing improves thalassemia carrier screening among premarital adults in a high prevalence population: the Dai nationality, China. *Genet Med*, 19(9), 1022-1031. doi:10.1038/gim.2016.218
- He, S., Li, J., Li, D. M., Yi, S., Lu, X., Luo, Y., . . . Qiu, X. (2018). Molecular characterization of alpha- and beta-thalassemia in the Yulin region of Southern China. *Gene*, 655, 61-64. doi:10.1016/j.gene.2018.02.058
- Martin, A., & Thompson, A. A. (2013). Thalassemias. *Pediatr Clin North Am*, 60(6), 1383-1391. doi:10.1016/j.pcl.2013.08.008
- NHS. (2017). NHS Sickle Cell and Thalassaemia Screening Programme. Retrieved from [https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment\\_data/file/656094/Antenatal\\_Laboratory\\_Handbook.pdf](https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/656094/Antenatal_Laboratory_Handbook.pdf)
- Nosheen, A., Ahmad, H., Qayum, I., Siddiqui, N., Abbasi, F. M., & Iqbal, M. S. (2015). Premarital genetic screening for beta thalassemia carrier status of indexed families using HbA2 electrophoresis. *J Pak Med Assoc*, 65(10), 1047-1049.
- PHE. (2019). Sickle cell and thalassaemia screening: commission and provide. Retrieved from <https://www.gov.uk/government/collections/sickle-cell-and-thalassaemia-screening-commission-and-provide>
- Ryan, K., Bain, B. J., Worthington, D., James, J., Plews, D., Mason, A., . . . Streetly, A. (2010). Significant haemoglobinopathies: guidelines for screening and diagnosis. *Br J Haematol*, 149(1), 35-49. doi:10.1111/j.1365-2141.2009.08054.x
- Satirapod, C., Sukprasert, M., Panthan, B., Charoenyingwattana, A., Chitayanan, P., Chantratita, W., . . . Hongeng, S. (2019). Clinical utility of combined preimplantation genetic testing methods in couples at risk of passing on beta thalassemia/hemoglobin E disease: A retrospective review from a single center. *PLoS One*, 14(11), e0225457. doi:10.1371/journal.pone.0225457
- Shook, L. M., Haygood, D., & Quinn, C. T. (2020). Clinical Utility of Confirmatory Genetic Testing to Differentiate Sickle Cell Trait from Sickle-beta(+)-Thalassemia by Newborn Screening. *Int J Neonatal Screen*, 6(1). doi:10.3390/ijns6010007
- Steinberg, M. H. (1999). Management of sickle cell disease. *N Engl J Med*, 340(13), 1021-1030. doi:10.1056/nejm199904013401307
- Steinberg, M. H. (2018). Structure and function of normal hemoglobins. Retrieved from [https://www.uptodate.com/contents/structure-and-function-of-normal-hemoglobins?sectionName=HEMOGLOBIN%20A2&search=alpha%20thalassemia&topicRef=7116&anchor=H1062835623&source=see\\_link#H1062835623](https://www.uptodate.com/contents/structure-and-function-of-normal-hemoglobins?sectionName=HEMOGLOBIN%20A2&search=alpha%20thalassemia&topicRef=7116&anchor=H1062835623&source=see_link#H1062835623)
- Tubman, V. N., Fung, E. B., Vogiatzi, M., Thompson, A. A., Rogers, Z. R., Neufeld, E. J., & Kwiatkowski, J. L. (2015). Guidelines for the Standard Monitoring of Patients with Thalassemia: Report of the Thalassemia Longitudinal Cohort. *J Pediatr Hematol Oncol*, 37(3), e162-169. doi:10.1097/mp.0000000000000307
- Wilson, R. D., De Bie, I., Armour, C. M., Brown, R. N., Campagnolo, C., Carroll, J. C., . . . Zwingerman, R. (2016). Joint SOGC–CCMG Opinion for Reproductive Genetic Carrier Screening: An Update for All Canadian Providers of Maternity and Reproductive Healthcare in the Era of Direct-to-Consumer Testing. *Journal of Obstetrics and Gynaecology Canada*, 38(8), 742-762.e743. doi:<https://doi.org/10.1016/j.jogc.2016.06.008>
- Yates, A. (2019). Prenatal screening and testing for hemoglobinopathy. Retrieved from [https://www.uptodate.com/contents/prenatal-screening-and-testing-for-hemoglobinopathy?search=alpha%20thalassemia&topicRef=7116&source=see\\_link#H10](https://www.uptodate.com/contents/prenatal-screening-and-testing-for-hemoglobinopathy?search=alpha%20thalassemia&topicRef=7116&source=see_link#H10)

# Genetic Testing for Alpha- and Beta- Thalassemia AHS – M2131

Medical Director review 7/2020

## **Policy Implementation/Update Information**

---

### **For policy titled: Genetic Testing for Alpha Thalassemia**

1/1/2019 BCBSNC will provide coverage for genetic testing for alpha thalassemia when it is determined to be medically necessary because the criteria and guidelines are met. Medical Director review 1/1/2019. (jd)

8/13/2019 Specialty Matched Consultant Advisory Panel review 7/2019. Medical Director review 7/2019. (jd)

### **For policy titled: Genetic Testing for Alpha- and Beta- Thalassemia:**

9/10/2019 Reviewed by Avalon 2<sup>nd</sup> Quarter 2019 CAB with title change. Added characteristics of beta-thalassemia and Related Policies to the Description section. Indications for beta-thalassemia added to the policy statement along with both the When Covered and When Not Covered sections. Policy guidelines updated. Added the following codes to the Billing/Coding section: S3845, S3846, and removed code table. References updated. Medical Director review 8/2019. (jd)

10/29/19 Wording in the Policy, When Covered, and/or Not Covered section(s) changed from Medical Necessity to Reimbursement language, where needed. (hb)

7/28/20 Reviewed by Avalon 2<sup>nd</sup> Quarter 2020 CAB. Added item 1 reimbursement language for genetic counseling and revised item 3 from medically necessary to reimbursement language under the When Covered section. Policy guidelines and references updated. Added the following codes to the Billing/Coding section: 81361, 81362, 81363, 81364, 96040, S0265. Specialty Matched Consultant Advisory Panel review 7/2020. Medical Director review 7/2020. (jd)

---

Medical policy is not an authorization, certification, explanation of benefits or a contract. Benefits and eligibility are determined before medical guidelines and payment guidelines are applied. Benefits are determined by the group contract and subscriber certificate that is in effect at the time services are rendered. This document is solely provided for informational purposes only and is based on research of current medical literature and review of common medical practices in the treatment and diagnosis of disease. Medical practices and knowledge are constantly changing and BCBSNC reserves the right to review and revise its medical policies periodically.