

Corporate Medical Policy

Mutation Analysis in Myeloproliferative Neoplasms AHS - M2101

File Name: mutation_analysis_in_myeloproliferative_neoplasms
Origination: 1/1/2019
Last Review: 4/2022

Description of Procedure or Service

Myeloproliferative neoplasms (MPN) are a heterogeneous group of clonal disorders characterized by overproduction of one or more differentiated myeloid lineages (Grinfeld, Nangalia, & Green, 2017). These include polycythemia vera (PV), essential thrombocythemia (ET), and primary myelofibrosis (PMF). The majority of MPN result from somatic mutations in the 3 driver genes, *JAK2*, *CALR*, and *MPL*, which represent major diagnostic criteria in combination with hematologic and morphological abnormalities (Rumi & Cazzola, 2017).

Related Policies:

BCR-ABL 1 Testing for Chronic Myeloid Leukemia AHS-M2027

******Note: This Medical Policy is complex and technical. For questions concerning the technical language and/or specific clinical indications for its use, please consult your physician.***

Policy

BCBSNC will provide coverage for mutation analysis in myeloproliferative neoplasms when it is determined to be medically necessary because the medical criteria and guidelines shown below are met.

Benefits Application

This medical policy relates only to the services or supplies described herein. Please refer to the Member's Benefit Booklet for availability of benefits. Member's benefits may vary according to benefit design; therefore member benefit language should be reviewed before applying the terms of this medical policy.

When Mutation Analysis in Myeloproliferative Neoplasms is covered

1. *JAK2*, *CALR* or *MPL* mutation testing is considered medically necessary for the diagnosis of patients presenting with clinical, laboratory, or pathological findings suggesting classic forms of myeloproliferative neoplasms (MPN), that is, polycythemia vera (PV), essential thrombocythemia (ET), or primary myelofibrosis (PMF) when ordered by a hematology and/or oncology specialist in the following situations:
 - A. For patients suspected to have PV, *JAK2*, *CALR*, or *MPL* mutation testing is considered medically necessary only if one of the following testing criteria are met:
 - i. Hemoglobin >16.5 g/dL in men; Hemoglobin >16.0 g/dL in women, or Hematocrit >49% in men; Hematocrit >48% in women, on two separate occasions or

Mutation Analysis in Myeloproliferative Neoplasms AHS - M2101

Increased red cell mass (More than 25% above mean normal predicted value), and no other known cause of erythrocytosis, or

- ii. Bone marrow biopsy showing hypercellularity for age with trilineage hyperplasia including prominent erythroid, granulocytic, and megakaryocytic proliferation with pleomorphic, mature megakaryocytes (differences in size)

B. For patients suspected to have essential thrombocythemia (ET) testing for *JAK2*, *CALR* and *MPL* mutations is considered medically necessary only if one of the following testing criteria are met:

- i. Platelet count $\geq 450 \times 10^9/L$ greater than 3 months OR
- ii. Bone marrow biopsy showing proliferation mainly of the megakaryocyte lineage with increased numbers of enlarged, mature megakaryocytes with hyper-lobulated nuclei. No significant increase or left shift in neutrophil granulopoiesis or erythropoiesis and very rarely minor (grade 1) increase in reticulin fibers.

C. For patients suspected to have primary myelofibrosis (PMF), testing for *JAK2*, *CALR* and *MPL* mutations if the following testing criteria are met:

- i. Patient has demonstrated leukocytosis of greater or equal to 11×10^9 on two separate occasions in the absence of other conditions that can cause leukocytosis or
- ii. Enlarged spleen or
- iii. BM biopsy shows megakaryocytic proliferation and atypia, without r81120, reticulin fibrosis $>$ grade 1, accompanied by increased age-adjusted BM cellularity, granulocytic proliferation, and often decreased erythropoiesis or
- iv. BM biopsy shows presence of megakaryocytic proliferation and atypia, accompanied by either reticulin and/or collagen fibrosis grades 2 or 3

2. In patients with a clinical suspicion of prePMF or overt PMF who have already tested negative for mutations in *JAK2*, *CALR*, or *MPL* and who do not meet the WHO criteria for BCR-AB1⁺ CML, PV, ET, myelodysplastic syndromes, or other myeloid neoplasms, screening for mutations in clonal markers *ASXL1*, *EZH2*, *TET2*, *IDH1/IDH2*, *SRSF2*, and *SFS3B1* is considered medically necessary.

3. *JAK2*, *CALR*, or *MPL* mutation testing is considered medically necessary in individuals diagnosed with Budd-Chiari Syndrome.

4. *JAK2*, *CALR*, or *MPL* mutation testing is considered medically necessary in individuals with persistent and unexplained erythrocytosis or thrombocytosis.

5. Screening for *JAK2* V617F is considered medically necessary in patients with normal blood counts and unexplained splanchnic vein thrombosis.

6. For patients suspected to have chronic neutrophilic leukemia, testing for *CSF3R* mutations is considered medically necessary.

7. Screening for *KIT* D816V is considered medically necessary in patients with a clinical suspicion of mastocytosis.

Mutation Analysis in Myeloproliferative Neoplasms AHS - M2101

NOTE: For 5 or more gene tests being run on the same platform, such as multi-gene panel next generation sequencing, please refer to AHS-R2162 Reimbursement Policy.

When Mutation Analysis in Myeloproliferative Neoplasms is not covered

JAK2 tyrosine kinase, *CALR*, and *MPL* mutation testing is considered **investigational** in all other cases.

Policy Guidelines

Scientific Background

Myeloproliferative neoplasms, including polycythemia vera (PV), essential thrombocythemia (ET), and primary myelofibrosis (PMF), arise from somatic mutation in hematopoietic stem cell (HSC) that clonally expand resulting in single or multilineage hyperplasia (Vainchenker & Kralovics, 2017). They are relatively rare, affecting 0.84 (PV), 1.03 (ET), and 0.47 (PMF) people per 100,000 worldwide; however, these may not be reflective of its true incidence due to the high heterogeneity of MPN (Titmarsh et al., 2014).

MPNs share features of bone marrow hypercellularity, increased incidence of thrombosis or hemorrhage, and an increased rate of progression to acute myeloid leukemia. Abnormalities in cytokine signaling pathways are common and usually lead to increased JAK-STAT signaling (Grinfeld et al., 2017). PV is characterized by erythrocytosis with suppressed endogenous erythropoietin production, bone marrow panmyelosis, and *JAK2* mutation leading to constitutive activation. ET is defined by thrombocytosis, bone marrow megakaryocytic proliferation, and presence of *JAK2*, *CALR*, or *MPL* mutation. PMF is characterized by bone marrow megakaryocytic proliferation, reticulin and/or collagen fibrosis, and presence of *JAK2*, *CALR*, or *MPL* mutation (Rumi & Cazzola, 2017). Mutations in other genes involved in signal transduction (*CBL*, *LNK/SH2B3*), chromatin modification (*TET2*, *EZH2*, *IDH1/2*, *ASXL1*, *DNM3TA*), RNA splicing (*SF3B1*, *SRSF2*, *U2AF1*), and tumor suppressor function (*TP53*) have also been reported and are considered “high-risk” (NCCN, 2019).

JAK2, which stands for “Janus Kinase 2”, is a gene whose mutation is responsible for a significant amount of MPNs. It is a mutation that causes hypersensitivity of hematopoietic progenitor cells to other cytokines, and this mutation typically appears on red blood cells or bone marrow cells. This mutation is often found on exon 12 or 14, and the exon 14 mutation results in a cytokine-independent activation of several regulatory pathways. *JAK2* mutations contribute to at least 95% of PV cases, about 50-65% of ET cases, and 60-65% of PMF cases (Ayalew Tefferi, 2020a, 2020b, 2021a).

MPL, which encodes a thrombopoietin receptor, also contributes to MPNs. *MPL* mutations result in a similar phenotype to *JAK2* mutations; both result in cytokine-independent growth of their targets. However, *MPL* mutations are not nearly as common as *JAK2* and *CALR* mutations, casting doubt on the clinical utility for testing. *MPL* mutations comprise up to 4% of ET cases and 5% of PMF cases (Ayalew Tefferi, 2020a, 2020b, 2021a).

CALR is a gene that encodes calreticulin (or calregulin), which is a Ca²⁺ binding protein. The mutation typically involves the creation of the incorrect Ca²⁺ binding region, thereby not allowing the protein to perform its regular duties such as maintaining calcium homeostasis. This results in a similar phenotype to the *JAK2* mutation, which is the cytokine-independent activation of regulatory pathways. *CALR* mutations contribute to approximately 15-25% of ET cases and 20-25% of PMF cases, and about 70% of ET or PMF patients without a *JAK2* or *MPL* mutation have this mutation (Ayalew Tefferi, 2020a, 2020b, 2021a).

The significance of *JAK2*, *MPL*, *CALR* and other mutations in the genesis of the MPNs as well as their roles in determining phenotype are unclear (Ayalew Tefferi, 2021b). However, integrated genomic analyses suggest that regardless of diagnosis or *JAK2* mutational status, MPNs are characterized by upregulation of JAK-STAT target genes, demonstrating the central importance of this pathway in the pathogenesis (Rampal et al., 2014). An Argentinean study focusing on establishing the frequency of *JAK2*,

Mutation Analysis in Myeloproliferative Neoplasms AHS - M2101

MPL, and *CALR* mutations and comparing their clinical and hematological features corroborates this importance. Mutations of *JAK2*V617F, *JAK2* exon 12, *MPL* W515L/K and *CALR* were analyzed in 439 patients with *BCR-ABL1*-negative MPN, and it was demonstrated that these mutations were present in 94.9% of the cases of polycythemia vera (PV), 85.5% in patients with essential thrombocythemia (ET), and 85.2% with primary myelofibrosis, leading the researchers to conclude that “the combined genetic tests of these driver mutations are essential for accurate diagnoses of *BCR-ABL1*-negative MPN” (Ojeda et al., 2018). This may lead to development of novel *JAK2* therapeutics (Silvennoinen & Hubbard, 2015). Thus, mutation analysis at the time of diagnosis has value for determining prognosis as well as individual risk assessment and guide treatment-making decisions (Hussein, Granot, Shpilberg, & Kreipe, 2013; Ayalew Tefferi, 2018, 2021b).

Neutrophilia, an increase in peripheral blood neutrophils at least two standard deviations above the mean, can be associated with any the MPNs. In chronic neutrophilic leukemia (CNL), *CSF3R* mutations have been discovered in the majority of patients with CNL (Coates, 2021; A. Tefferi, Thiele, Vannucchi, & Barbui, 2014). A study released in 2013 reported 16 of 27 patients with CNL or atypical chronic myeloid leukemia (aCML) had activating mutations in *CSF3R* (Maxson et al., 2013). *SETBP1* has also been used as a part of comprehensive mutation profiling in distinguishing aCML and chronic myelomonocytic leukemia (CMML). A 2019 NGS study reports significant differences in the profiles of patients with aCML or CMML when comparing *TET2*, *SETBP1*, and *CSF3R*. The researchers conclude, “differential mRNA expression could be detected between both cohorts in a subset of genes (*FLT3*, *CSF3R*, and *SETBP1* showed the strongest correlation). However, due to high variances in the mRNA expression, the potential utility for the clinic is limited (Faisal et al., 2019).”

In 2017 the FDA approved ipsogen® *JAK2* RGQ PCR Kit (FDA, 2017b) to detect Janus Tyrosine Kinase 2 (*JAK2*) gene mutation G1849T (V617F) with an allele-specific, quantitative, polymerase chain reaction (PCR) using an amplification refractory mutation system (ARMS). The device marketing authorization was based on data from a clinical study of 473 suspected patients with MPNs, 276 with suspected PV, 98 with suspected ET, and 99 with suspected PMF. The study compared results from the ipsogen *JAK2* RGQ PCR Kit to results obtained with independently validated bi-directional sequencing. The study found that the ipsogen *JAK2* RGQ PCR Kit test was in 96.8% agreement with the reference method, 100% in positive agreement, and 95.1% in negative agreement, with 458 samples in agreement out of 473. The concordance with each condition was also high; agreement of 90.8% within the ET samples (89/98), 94.9% agreement within the PMF samples (94/99), and 99.6% within the PV samples (275/276). All three conditions had positive agreements of 100%. The authors went on to note that the 15 samples with disagreeing results had mutation levels under the detection capability of bi-directional sequencing. To validate these 15 samples, an independently validated NGS panel was used to compare results with the kit, and all 15 samples were found to test positive, thereby agreeing with the kit. The authors concluded that the kit was accurate for any mutation levels at or above 1% (FDA, 2017a).

Genoptix, Inc. offers multiple testing options for *JAK2* testing. One option is a myeloid molecular profile test of genomic DNA from either bone marrow aspirates or blood. These test uses NGS technology to sequence 44 different genes, including *JAK2*, *CLR*, *MPL*, *SETBP1*, and *CSF3R* in addition to *KRAS*, *NRAS*, and other genes. Genoptix claims to identify at least one somatic mutation in 80 – 90% of patients with a myelodysplastic syndrome, including MPN (Genoptix, 2018b). Genoptix also offers a test titled “MPN Targeted Profile”, an NGS test focused on targeted regions of *JAK2*, *CALR*, *MPL*, *CSF3R*, and *SETBP1*. They note that “the MPN Molecular Profile is intended as an aid in the diagnosis and subclassification of *BCR-ABL1*-negative myeloproliferative neoplasms (MPN) (Genoptix, 2018a).”

Many other proprietary tests are available for mutational analysis in MPN. IntelliGEN® Myeloid is a NGS assay that analyzes fifty genes for somatic mutations that could be useful in providing diagnostic or prognostic information for patients with MDS, AML, or MPN (Labcorp, 2022). The LeukoVantage® Myeloid Neoplasm Mutation Panel detects myeloid neoplasm-associated mutations in 48 genes associated with AML, MDS, and MPN. The LeukoVantage AML panel can be used to assess AML subclass and prognosis based on genetic abnormalities in *NPM1*, *CEBPA*, and *RUNX1* (Quest_Diagnostics, 2020). NeoGenomics offers two tests which include the MPN Reflex Test and NeoTYPE® Myeloid Disorders Profile. The MPN Reflex Test is a sequential testing panel for qualitative detection of *JAK2* V617F, *JAK2*

Mutation Analysis in Myeloproliferative Neoplasms AHS - M2101

Exon 12-14, CALR exon 9, and MPL exon 10 (NeoGenomics, 2022a). NeoTYPE® Myeloid Disorders Profile is a 63 gene panel that targets known mutations associated with AML, MPN, MDS, CML, chronic myelomonocytic leukemia (CMML) and juvenile myelomonocytic leukemia (JMML) (NeoGenomics, 2022b). Centogene has released a Myeloid Tumor Panel which targets 35 genes that are associated with myeloid malignancies which also include AML, MPN, MDS, CML, CMML, and JMML (Centogene, 2022).

Analytical Validity

Poluben et al. analyzed the characteristics of myeloproliferative neoplasms (MPN) in patients exposed to ionizing radiation (IR) from the 1986 Chernobyl accident. 281 patients (90 exposed to radiation, 181 unexposed) were included. *JAK2*, *MPL*, and *CALR* mutations were identified. IR-exposed patients had several different genetic features compared to the unexposed cohort: lower rate of *JAK2* V617F mutations (58.4% vs 75.4%), higher rate of type 1-like *CALR* mutations (12.2% vs 3.1%), higher rate of triple-negative cases (27.8% vs 16.2%), and higher rate of “potentially pathogenic” sequence variants (4.8 vs 3.1). The authors suggested IR-exposed patients as a cohort with “distinct” genomic characteristics (Poluben et al., 2019).

Rosenthal studied the analytical validity of a 48-gene NGS panel for detecting mutations in myeloid neoplasms. The panel detects single nucleotide variations (SNVs), insertions/deletions, and *FLT3* internal tandem duplications (*FLT3*-ITD). 184 samples were analyzed using the 48-gene panel and compared to those identified by a 35-gene hematologic neoplasms panel using an additional 137 samples. Analytical validation yielded 99.6% sensitivity and 100% specificity. Concordance of variants detected by the 2 tested panels was 100%. “Among patients with suspected myeloid neoplasms, 54.5% patients had at least one clinically significant mutation: 77% in AML patients, 48% in MDS, and 45% in MPN.” The authors conclude that “the assay can identify mutations associated with diagnosis, prognosis, and treatment options of myeloid neoplasms even in technically challenging genes” (Rosenthal et al., 2021).

Guidelines and Recommendations

World Health Organization (WHO)

The 2017 edition of the World Health Organization’s classification of myeloid neoplasm and acute leukemia proposed the following criteria for the diagnosis of PV, ET and PMF.

WHO Criteria for PV

Diagnosis of PV requires meeting either all 3 major criteria, or the first 2 major criteria and the minor criterion:

Major Criteria

1. Hemoglobin >16.5 g/dL in men; Hemoglobin >16.0 g/dL in women, or Hematocrit >49% in men; Hematocrit >48% in women, or Increased red cell mass (More than 25% above mean normal predicted value)
2. Bone marrow biopsy showing hypercellularity for age with trilineage growth (panmyelosis) including prominent erythroid, granulocytic, and megakaryocytic proliferation with pleomorphic, mature megakaryocytes (differences in size)
3. Presence of *JAK2V617F* or *JAK2* exon 12 mutation

Minor Criteria

Subnormal serum erythropoietin level

WHO Criteria for ET

Diagnosis of ET requires meeting all 4 major criteria or the first 3 major criteria and the minor criterion:

Major Criteria

1. Platelet count $\geq 450 \times 10^9/L$

Mutation Analysis in Myeloproliferative Neoplasms AHS - M2101

2. Bone marrow biopsy showing proliferation mainly of the megakaryocyte lineage with increased numbers of enlarged, mature megakaryocytes with hyperlobulated nuclei. No significant increase or left shift in neutrophil granulopoiesis or erythropoiesis and very rarely minor (grade 1) increase in reticulin fibers
3. Not meeting WHO criteria for BCR-ABL1+ CML, PV, PMF, myelodysplastic syndromes, or other myeloid neoplasms
4. Presence of *JAK2*, *CALR*, or *MPL* mutation

Minor Criteria

Presence of a clonal marker or absence of evidence for reactive thrombocytosis

WHO Criteria for PrePMF

Diagnosis of prePMF requires meeting all 3 major criteria, and at least 1 minor criterion:

Major Criteria

1. Megakaryocytic proliferation and atypia, without reticulin fibrosis >grade 1, accompanied by increased age-adjusted BM cellularity, granulocytic proliferation, and often decreased erythropoiesis
2. Not meeting the WHO criteria for BCR-ABL1+ CML, PV, ET, myelodysplastic syndromes, or other myeloid neoplasms
3. Presence of *JAK2*, *CALR*, or *MPL* mutation or in the absence of these mutations, presence of another clonal marker (e.g. *ASXL1*, *EZH2*, *TET2*, *IDH1/IDH2*, *SRSF2*, *SF3B1*), or absence of minor reactive BM reticulin fibrosis

Minor Criteria

1. Anemia not attributed to a comorbid condition
2. Leukocytosis $\geq 11 \times 10^9/L$
3. Palpable splenomegaly
4. LDH increased to above upper normal limit of institutional reference range

WHO Criteria for Overt PMF

Diagnosis of overt PMF requires meeting all 3 major criteria, and at least 1 minor criterion

Major Criteria

1. Presence of megakaryocytic proliferation and atypia, accompanied by either reticulin and/or collagen fibrosis grades 2 or 3
2. Not meeting WHO criteria for ET, PV, BCR-ABL1+ CML, myelodysplastic syndromes, or other myeloid neoplasms
3. Presence of *JAK2*, *CALR*, or *MPL* mutation or in the absence of these mutations, presence of another clonal marker (e.g. *ASXL1*, *EZH2*, *TET2*, *IDH1/IDH2*, *SRSF2*, *SF3B1*), or absence of reactive myelofibrosis

Minor Criteria

1. Anemia not attributed to a comorbid condition
2. Leukocytosis $\geq 11 \times 10^9/L$
3. Palpable splenomegaly
4. LDH increased to above upper normal limit of institutional reference range
5. Leukoerythroblastosis (T. Barbui et al., 2018)

These guidelines also list four additional “clinicopathologic entities” for MPNs: “chronic myeloid leukemia (CML), chronic neutrophilic leukemia (CNL), chronic eosinophilic leukemia, not otherwise specified (CELNOS) and MPN, unclassifiable (MPN-U)”. The guidelines note that although *CSF3R* mutations are “specific” to WHO-defined CNL, they also remark that “the presence of a membrane

Mutation Analysis in Myeloproliferative Neoplasms AHS - M2101

proximal *CSF3R* mutation in a patient with neutrophilic granulocytosis should be sufficient for the diagnosis of CNL, regardless of the degree of leukocytosis” (T. Barbui et al., 2018).

European LeukemiaNet (ELN)

ELN guidelines also recommend “strict adherence” to these guidelines for the three categories of Philadelphia-negative MPNs, (i.e. ET, PV, and MF) (Tiziano Barbui et al., 2018).

However, they also recommend “searching” for complementary clonal markers such as *ASXL1*, *EZH2*, *IDH1/2*, and *SRSF2* for patients that tested negative for the three driver mutations and have bone marrow features as well as a clinical phenotype consistent with myelofibrosis (Tiziano Barbui et al., 2018).

National Comprehensive Cancer Network

The NCCN Guidelines Version 2.2021 for Myeloproliferative Neoplasms recommends “molecular testing for *JAK2* V617F mutations as part of an initial workup for all patients molecular testing for *CALR* and *MPL* mutations should be performed for ET and PMF patients, and molecular testing for *JAK2* exon 12 should be done for patients who test negative for *JAK2* but are suspected for PV. An NGS panel including *JAK2*, *CALR*, and *MPL* may also be used. The NCCN follows the 2017 edition of the WHO diagnostic criteria for all three conditions. The NCCN does state that NGS “may be useful to establish clonality in selected circumstances (eg, triple negative non-mutated *JAK2*, *MPL*, and *CALR*). They include a list of somatic mutations with prognostic significance in individuals with MPN that includes the *ASXL1*, *EZH2*, *IDH1/2*, *SRSF2*, *TP53*, and *U2AF1 Q157*.”

Finally, the NCCN recommends following the 2017 WHO diagnostic criteria to diagnose MPNs (NCCN, 2021).

British Society for Hematology (BSH)

The BSH recommends testing for *CALR* for patients suspected of ET and PMR, as *CALR* mutations account for most patients without either a *JAK2* or *MPL* mutation. The authors found that as many as one third of ET and PMF patients had a mutation in exon 9 of the *CALR* gene (Harrison et al., 2014).

The BSH also published guidelines on the diagnosis of polycythemia vera (PV). In it, they divide PV into *JAK2*-positive and *JAK2*-negative PV. For *JAK2*-positive PV, the only two diagnostic criteria are as follows:

- “High hematocrit (>0.52 in men, >0.48 in women) OR raised red cell mass (>25% above predicted)”
- “Mutation in *JAK2*”

For *JAK2*-negative PV, the diagnostic criteria are as follows (requiring A1-A4, as well as another “A” criteria or two “B” criteria).

- “A1 Raised red cell mass (>25% above predicted) OR hematocrit ≥ 0.60 in men, ≥ 0.56 in women”
- “A2 Absence of mutation in *JAK2*”
- “A3 No cause of secondary erythrocytosis”
- “A4 Bone marrow histology consistent with polycythemia vera”
- “A5 Palpable splenomegaly”

Mutation Analysis in Myeloproliferative Neoplasms AHS - M2101

- “A6 Presence of an acquired genetic abnormality (excluding BCR-ABL1) in the hematopoietic cells”
- “B1 Thrombocytosis (platelet count $>450 \times 10^9 /l$)”
- “B2 Neutrophil leucocytosis (neutrophil count $>10 \times 10^9 /l$ in non-smokers, $\geq 12.5 \times 10^9 /l$ in smokers)”
- “B3 Radiological evidence of splenomegaly”
- “B4 Low serum erythropoietin”

The guidelines also note that investigation of erythrocytosis should be undertaken to properly identify the diagnosis. The BSH remarks that EPO receptor mutations may be a primary cause for erythrocytosis and that *EGNLI*, *VHL*, and *EPAS1* mutations may be a secondary cause. Other hemoglobinopathies caused by mutations in genes such as *HBA1*, *HBA2*, *HBB*, or *BGPM* may also be a factor (McMullin et al., 2019).

In 2021, the BSH published guidelines on the use of genetic tests to diagnose and manage patients with myeloproliferative neoplasms. The following recommendations were made:

1. “Molecular screening for *JAK2*, *CALR* and *MPL* variants as appropriate is recommended in patients with persistent erythrocytosis or thrombocytosis (GRADE 1B).
2. Screening for *JAK2* V617F is recommended in cases with normal blood counts and unexplained splanchnic vein thrombosis (GRADE 1B) and may be considered in selected patients with unexplained cerebral vein thrombosis (GRADE 2C).
3. Screening for *CALR* variants may be considered in patients with splanchnic vein thrombosis or cerebral vein thrombosis (GRADE 2C).
4. Screening for *JAK2*, *CALR* and *MPL* variants should be considered for patients with arterial or unprovoked venous thrombosis who have a mildly or variably elevated hematocrit or platelet count that persists for 2–3 months (GRADE 2C).
5. *BCR-ABL1* should be excluded in cases with persistent thrombocytosis negative for *JAK2*, *CALR* and *MPL* variants or with atypical features (GRADE 1B).
6. Younger patients (e.g. under 60 years) with bone marrow histology typical of ET [or myeloproliferative neoplasm, unclassifiable (MPN-U) or suspected prefibrotic MF] where confirmation of a clonal disorder would be useful in view of the patient’s likely long-term disease course and ideally where a broad panel that covers non-canonical variants in *JAK2* and *MPL* and a range of other driver genes is available.
7. Patients with significant thrombocytosis (e.g. platelet count $> 600 \times 10^9/l$), no reactive cause and borderline bone marrow histology, where cytoreduction would be indicated if there was convincing evidence of a clonal disorder. Examples would include those with an unexplained thrombotic event, particularly younger patients. For older patients without thrombosis, testing may be considered but results must be interpreted with caution in view of the possibility of incidental CH.
8. A myeloid gene panel and cytogenetic analysis (or equivalent) is recommended for patients with bone marrow histology and clinical features consistent with PMF (+/- suggestive features of MDS or MDS/MPN) who test negative for *JAK2/CALR/MPL* (GRADE 1B).
9. A myeloid gene panel and cytogenetic analysis (or equivalent) is not recommended for most patients with *JAK2/CALR/MPL*-negative erythrocytosis or thrombocytosis but may be considered in individual cases (GRADE 2C).
10. Myeloid gene panel testing is recommended for MPN cases who test positive for *JAK2/CALR/MPL* mutations and have additional cytopenias(s) at diagnosis, unexplained ring sideroblasts or other dysplasia, increased blasts (including blastic transformation), peripheral-blood monocytosis or atypical clinical features (GRADE 1B).
11. Myeloid gene panel testing and conventional karyotyping are recommended for all patients with PMF, post-PV or post-ET MF who are candidates for allogeneic stem cell transplant (GRADE 1B).
12. Myeloid gene panel testing should be considered for other patients if the additional genomic data will guide clinical management (GRADE 2C).

Mutation Analysis in Myeloproliferative Neoplasms AHS - M2101

13. High-sensitivity assays of mutant allele burden are recommended following post-allogeneic stem cell transplant to monitor for residual disease (GRADE 1C).
14. Quantitative assays of mutant allele burden are not recommended for most MPN patients but may be considered where demonstration of molecular response would influence clinical management (GRADE 2C).
15. Patients with persistent eosinophilia should be investigated initially for *FIP1L1–PDGFRA* by FISH and/or nested RT-PCR (GRADE 1B).
16. BM cytogenetics or FISH is recommended to screen for other fusion genes, which must then be confirmed by molecular methods (GRADE 1B).
17. Myeloid gene panel and *KIT* D816V testing should be considered for patients with persistent unexplained eosinophilia who test negative for fusion genes (GRADE 2B).
18. Testing for *CSF3R* variants, preferably as part of wider myeloid panel, is recommended for all patients with suspected CNL (Grade 2B).
19. Sensitive testing for *KIT* D816V is recommended for all patients with a clinical suspicion of mastocytosis (GRADE 1B).
20. If negative for *KIT* D816V, screening for other *KIT* mutations should be considered for adults (but is recommended for children) (GRADE 1B).
21. Myeloid panel analysis is recommended for patients with advanced SM who are candidates for allogeneic stem cell transplantation (GRADE 1B).
22. Myeloid panel analysis may be considered for other SM patients if the apparent aggressiveness of the disease might influence options for therapy (GRADE 2B).
23. Myeloid panel and/or BM cytogenetics should be considered to characterize the AHN component of SM-AHN (GRADE 2B).
24. *BCR–ABL1* should be excluded in all cases of suspected MDS/MPN, and rearrangements associated with MLN-eo should be excluded in cases with eosinophilia (GRADE 1B).
25. Myeloid gene panel analysis and BM cytogenetics or SNP array is recommended for patients diagnosed with MDS/MPN and for cases with suspected MDS/MPN but with indeterminate morphology (GRADE 1B)” (Cross et al., 2021).

European Association for the Study of the Liver

For myeloproliferative neoplasms, the EASL recommends testing for JAK2 V617F mutations in splanchnic vein thrombosis patients, as well as patients with normal peripheral blood cell counts. If the JAK2 mutation test is negative, a calreticulin mutation test should be performed, and if both are negative, a bone marrow histology analysis should be performed (EASL, 2016).

European Society of Medical Oncology

The ESMO recommends that anyone with a suspected MPN be tested for the three driver mutations (JAK2, CALR, MPL) and that genotyping should be obtained at diagnosis. However, the ESMO states that it is not recommended to repeat testing in follow-up or assessing response to treatment, except for “allogeneic stem-cell transplantation and possibly interferon treatment”. For these two assessments a detection limit of $\leq 1\%$ is recommended. The ESMO also notes that conventional sequencing methods (PCR, melting analysis) may be used for detecting mutations (Vannucchi et al., 2015).

State and Federal Regulations, as applicable

Food and Drug Administration (FDA)

On July 28, 2017 the FDA approved Ipsogen® *JAK2 RGQ* PCR Kit (FDA, 2017b) to detect Janus Tyrosine Kinase 2 (*JAK2*) gene mutation G1849T (V617F) with an allele-specific, quantitative, polymerase chain reaction (PCR) using an amplification refractory mutation system (ARMS). This is the first FDA-authorized test intended to help physicians in evaluating patients for suspected Polycythemia Vera (PV).

Mutation Analysis in Myeloproliferative Neoplasms AHS - M2101

However, the FDA specifically states that this test is not intended for a stand-alone diagnosis of an MPN, nor can it detect less common mutations for MPN such as an exon 12 mutation (FDA, 2017a).

Many labs have developed specific tests that they must validate and perform in house. These laboratory-developed tests (LDTs) are regulated by the Centers for Medicare and Medicaid (CMS) as high-complexity tests under the Clinical Laboratory Improvement Amendments of 1988 (CLIA '88). As an LDT, the U. S. Food and Drug Administration has not approved or cleared this test; however, FDA clearance or approval is not currently required for clinical use.

Billing/Coding/Physician Documentation Information

This policy may apply to the following codes. Inclusion of a code in this section does not guarantee that it will be reimbursed. For further information on reimbursement guidelines, please see Administrative Policies on the Blue Cross Blue Shield of North Carolina web site at www.bcbsnc.com. They are listed in the Category Search on the Medical Policy search page.

Applicable service codes: 81120, 81121, 81175, 81176, 81219, 81236, 81237, 81270, 81279, 81338, 81339, 81348, 81450, 81455, 81479, 0017U, 0027U

BCBSNC may request medical records for determination of medical necessity. When medical records are requested, letters of support and/or explanation are often useful, but are not sufficient documentation unless all specific information needed to make a medical necessity determination is included.

Scientific Background and Reference Sources

For Policy Titled: JAK2, CALR, MPL Mutation Analysis in Myeloproliferative Neoplasms

Barbui, T., Tefferi, A., Vannucchi, A. M., Passamonti, F., Silver, R. T., Hoffman, R., . . . Barosi, G. (2018). Philadelphia chromosome-negative classical myeloproliferative neoplasms: revised management recommendations from European LeukemiaNet. *Leukemia*, 32(5), 1057-1069. doi:10.1038/s41375-018-0077-1

Barbui, T., Thiele, J., Gisslinger, H., Kvasnicka, H. M., Vannucchi, A. M., Guglielmelli, P., . . . Tefferi, A. (2018). The 2016 WHO classification and diagnostic criteria for myeloproliferative neoplasms: document summary and in-depth discussion. *Blood Cancer J*, 8(2), 15. doi:10.1038/s41408-018-0054-y

Centogene. (2022). Myeloid Tumor Panel. Retrieved from <https://www.centogene.com/diagnostics/somatic-mutation-testing/somatic-mutation-testing-for-myeloid-tumors.html>

Coates, T. D. (2021, January 04, 2021). Approach to the patient with neutrophilia. Retrieved from <https://www.uptodate.com/contents/approach-to-the-patient-with-neutrophilia>

Cross, N. C. P., Godfrey, A. L., Cargo, C., Garg, M., Mead, A. J., & Paper, A. B. S. f. H. G. P. (2021). The use of genetic tests to diagnose and manage patients with myeloproliferative and myeloproliferative/myelodysplastic neoplasms, and related disorders. *British Journal of Haematology*, 195(3), 338-351. doi:<https://doi.org/10.1111/bjh.17766>

EASL. (2016). EASL Clinical Practice Guidelines: Vascular diseases of the liver. *J Hepatol*, 64(1), 179-202. doi:10.1016/j.jhep.2015.07.040

Mutation Analysis in Myeloproliferative Neoplasms AHS - M2101

Faisal, M., Stark, H., Busche, G., Schlue, J., Teiken, K., Kreipe, H. H., . . . Bartels, S. (2019). Comprehensive mutation profiling and mRNA expression analysis in atypical chronic myeloid leukemia in comparison with chronic myelomonocytic leukemia. *Cancer Med.* doi:10.1002/cam4.1946

FDA. (2017a). Retrieved from https://www.accessdata.fda.gov/cdrh_docs/pdf17/K172287.pdf

FDA. (2017b). Approved Drugs - Ipsogen JAK2 RGQ PCR Kit (WebContent). Retrieved from <https://www.fda.gov/Drugs/InformationOnDrugs/ApprovedDrugs/ucm551474.htm>. from Center for Drug Evaluation and Research
<https://www.fda.gov/Drugs/InformationOnDrugs/ApprovedDrugs/ucm551474.htm>

Genoptix. (2018a). MPN TARGETED PROFILE. *HEMATOLOGIC DISEASES*. Retrieved from <https://genoptix.com/test-menu/mpn-targeted-profile/>

Genoptix. (2018b). MYELOID MOLECULAR PROFILE. *HEMATOLOGIC DISEASES*. Retrieved from <https://genoptix.com/test-menu/myeloid-molecular-profile/>

Grinfeld, J., Nangalia, J., & Green, A. R. (2017). Molecular determinants of pathogenesis and clinical phenotype in myeloproliferative neoplasms. *Haematologica*, 102(1), 7-17. doi:10.3324/haematol.2014.113845

Harrison, C. N., Butt, N., Campbell, P., Conneally, E., Drummond, M., Green, A. R., . . . McMullin, M. F. (2014). Modification of British Committee for Standards in Haematology diagnostic criteria for essential thrombocythaemia. *British Journal of Haematology*, 167(3), 421-423. doi:10.1111/bjh.12986

Hussein, K., Granot, G., Shpilberg, O., & Kreipe, H. (2013). Clinical utility gene card for: familial polycythaemia vera. *Eur J Hum Genet*, 21(6). doi:10.1038/ejhg.2012.216

Labcorp. (2022). IntelliGEN® Myeloid. Retrieved from <https://oncology.labcorp.com/cancer-care-team/test-menu/intelligen-myeloid>

Maxson, J. E., Gotlib, J., Pollyea, D. A., Fleischman, A. G., Agarwal, A., Eide, C. A., . . . Tyner, J. W. (2013). Oncogenic *CSF3R* mutations in chronic neutrophilic leukemia and atypical CML. *N Engl J Med*, 368(19), 1781-1790. doi:10.1056/NEJMoa1214514

McMullin, M. F., Harrison, C. N., Ali, S., Cargo, C., Chen, F., Ewing, J., . . . the, B. S. H. C. (2019). A guideline for the diagnosis and management of polycythaemia vera. A British Society for Haematology Guideline. *British Journal of Haematology*, 184(2), 176-191. doi:10.1111/bjh.15648

NCCN. (2019). NCCN Clinical Practice Guidelines in Oncology; Myeloproliferative Neoplasms v3.2019. Retrieved from https://www.nccn.org/professionals/physician_gls/pdf/mpn.pdf
https://www.nccn.org/professionals/physician_gls/pdf/mpn.pdf

NCCN. (2021). NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines (R)): Myeloproliferative Neoplasms. Retrieved from https://www.nccn.org/professionals/physician_gls/pdf/mpn.pdf

NeoGenomics. (2022a). MPN JAK2 V617F with Sequential Reflex to JAK2 Exon 12-13, CALR, and MPL. Retrieved from <https://neogenomics.com/test-menu/mpn-jak2-v617f-sequential-reflex-jak2-exon-12-13-calr-and-mpl>

NeoGenomics. (2022b). NeoTYPE® Myeloid Disorders Profile. Retrieved from <https://neogenomics.com/test-menu/neotype-myeloid-disorders-profile>

Mutation Analysis in Myeloproliferative Neoplasms AHS - M2101

- Ojeda, M. J., Bragós, I. M., Calvo, K. L., Williams, G. M., Carbonell, M. M., & Pratti, A. F. (2018). CALR, JAK2 and MPL mutation status in Argentinean patients with BCR-ABL1- negative myeloproliferative neoplasms. *Hematology*, 23(4), 208-211. doi:10.1080/10245332.2017.1385891
- Poluben, L., Puligandla, M., Neuberger, D., Bryke, C. R., Hsu, Y., Shumeiko, O., . . . Fraenkel, P. G. (2019). Characteristics of myeloproliferative neoplasms in patients exposed to ionizing radiation following the Chernobyl nuclear accident. *Am J Hematol*, 94(1), 62-73. doi:10.1002/ajh.25307
- Quest_Diagnostics. (2020). LeukoVantage® Myeloid Neoplasm Mutation Panels. Retrieved from <https://www.questdiagnostics.com/healthcare-professionals/clinical-education-center/faq/faq208>
- Rampal, R., Al-Shahrour, F., Abdel-Wahab, O., Patel, J. P., Brunel, J. P., Mermel, C. H., . . . Levine, R. L. (2014). Integrated genomic analysis illustrates the central role of JAK-STAT pathway activation in myeloproliferative neoplasm pathogenesis. *Blood*, 123(22), e123-133. doi:10.1182/blood-2014-02-554634
- Rosenthal, S. H., Gerasimova, A., Ma, C., Li, H. R., Grupe, A., Chong, H., . . . Lachawan, F. (2021). Analytical validation and performance characteristics of a 48-gene next-generation sequencing panel for detecting potentially actionable genomic alterations in myeloid neoplasms. *PLoS One*, 16(4), e0243683. doi:10.1371/journal.pone.0243683
- Rumi, E., & Cazzola, M. (2017). Diagnosis, risk stratification, and response evaluation in classical myeloproliferative neoplasms. *Blood*, 129(6), 680-692. doi:10.1182/blood-2016-10-695957
- Silvennoinen, O., & Hubbard, S. R. (2015). Molecular insights into regulation of JAK2 in myeloproliferative neoplasms. *Blood*, 125(22), 3388-3392. doi:10.1182/blood-2015-01-621110
- Tefferi, A. (2018). Overview of the myeloproliferative neoplasms - UpToDate. In A. Rosmarin (Ed.), *UpToDate*. Retrieved from https://www.uptodate.com/contents/overview-of-the-myeloproliferative-neoplasms?search=myeloproliferative%20disorders&usage_type=default&source=search_result&selectedTitle=1~150&display_rank=1#H15.
- Tefferi, A. (2020a). Clinical manifestations and diagnosis of primary myelofibrosis. Retrieved from https://www.uptodate.com/contents/clinical-manifestations-and-diagnosis-of-primary-myelofibrosis?topicRef=4716&source=see_link
- Tefferi, A. (2020b). Diagnosis and clinical manifestations of essential thrombocythemia. Retrieved from https://www.uptodate.com/contents/diagnosis-and-clinical-manifestations-of-essential-thrombocythemia?topicRef=4716&source=see_link
- Tefferi, A. (2021a). Overview of the myeloproliferative neoplasms - UpToDate. In A. Rosmarin (Ed.), *UpToDate*. Retrieved from https://www.uptodate.com/contents/overview-of-the-myeloproliferative-neoplasms?search=myeloproliferative%20disorders&usage_type=default&source=search_result&selectedTitle=1~150&display_rank=1#H15
- Tefferi, A. (2021b). Overview of the myeloproliferative neoplasms - UpToDate. In A. Rosmarin (Ed.), *UpToDate*. Retrieved from https://www.uptodate.com/contents/overview-of-the-myeloproliferative-neoplasms?search=myeloproliferative%20disorders&usage_type=default&source=search_result&selectedTitle=1~150&display_rank=1#H15
- Tefferi, A., Thiele, J., Vannucchi, A. M., & Barbui, T. (2014). An overview on CALR and CSF3R mutations and a proposal for revision of WHO diagnostic criteria for myeloproliferative neoplasms. *Leukemia*, 28(7), 1407-1413. doi:10.1038/leu.2014.35

Mutation Analysis in Myeloproliferative Neoplasms AHS - M2101

Titmarsh, G. J., Duncombe, A. S., McMullin, M. F., O'Rorke, M., Mesa, R., De Vocht, F., . . . Anderson, L. A. (2014). How common are myeloproliferative neoplasms? A systematic review and meta-analysis. *Am J Hematol*, 89(6), 581-587.

Vainchenker, W., & Kralovics, R. (2017). Genetic basis and molecular pathophysiology of classical myeloproliferative neoplasms. *Blood*, 129(6), 667-679. doi:10.1182/blood-2016-10-695940

Vannucchi, A. M., on behalf of the, E. G. C., Barbui, T., on behalf of the, E. G. C., Cervantes, F., on behalf of the, E. G. C., . . . on behalf of the, E. G. C. (2015). Philadelphia chromosome-negative chronic myeloproliferative neoplasms: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up†. *Annals of Oncology*, 26(suppl_5), v85-v99. doi:10.1093/annonc/mdv203

For Policy Titled: Mutation Analysis in Myeloproliferative Neoplasms

Medical Director review 5/2019

Medical Director review 8/2019

Specialty Matched Consultant Advisory Panel review- 11/2019

Medical Director review 11/2019

Medical Director review 4/2020

Specialty Matched Consultant Advisory Panel review- 11/2020

Medical Director review 11/2020

Medical Director review 4/2021

Specialty Matched Consultant Advisory Panel review- 8/2021

Policy Implementation/Update Information

For Policy Titled: JAK2, CALR, MPL Mutation Analysis in Myeloproliferative Neoplasms

1/1/2019 New policy developed. BCBSNC will provide coverage for JAK2, CALR, MPL mutation analysis in myeloproliferative neoplasms when it is determined to be medically necessary and criteria are met. Medical Director review 1/1/2019. Policy noticed 1/1/2019 for effective date 4/1/2019. (lpr)

For Policy Titled: Mutation Analysis in Myeloproliferative Neoplasms

5/14/19 Reviewed by Avalon 1st Quarter 2019 CAB. Added related policies section. Under When Not Covered section B., removed statement: should first be tested for JAK2 mutations; if testing is negative, further testing to detect CALR and MPL mutation and for patient suspected to have ET.” Under When Not Covered section C, removed statement: should first be tested JAK2 mutations; if testing is negative further testing to detect CALR and MPL mutations, for patient suspected to have PMF.” Under When Not Covered section, added statement: “If testing five or more genes, refer to policy AHS-M2109 Molecular Panel Testing of Cancers to Identify Targeted Therapy.” Updated Policy Guidelines section. Added PLA codes 0017U and 0027U to Billing/Coding section. Title changed from “JAK2, CALR, MPL Mutation Analysis in Myeloproliferative Neoplasms” to “Mutation Analysis in Myeloproliferative Neoplasms.” Medical Director review 5/2019. (lpr)

Mutation Analysis in Myeloproliferative Neoplasms AHS - M2101

- 10/1/19 Reviewed by Avalon 2nd Quarter 2019 CAB. Under “When Covered” section added: NOTE: For 5 or more gene tests being run on the same platform, such as multi-gene panel next generation sequencing, please refer to AHS-R2162 Reimbursement Policy. Deleted coding table from Billing/Coding section. Medical Director review 8/2019. (lpr)
- 12/31/19 Specialty Matched Consultant Advisory Panel review 11/20/2019. No change to policy statement. (lpr)
- 5/12/20 Reviewed by Avalon 1st Quarter 2020 CAB. Medical Director review 4/2020. Updated Description, Policy Guidelines, References. (lpr)
- 12/8/20 Specialty Matched Consultant Advisory Panel review 11/18/2020. No change to policy statement. (lpr)
- 5/4/21 Reviewed by Avalon 1st Quarter 2021 CAB. Medical Director review 4/2021. Under “When Covered” section, removed statements: “Patients suspected to have polycythemia vera (PV) should first be tested for the most common finding JAK2V617F; and If testing for PV is negative, further testing to detect other JAK2 tyrosine kinase mutations, eg. in exon 12”; added CALR and MPL testing indication for Budd-Chiari Syndrome and PV. Under Billing/Coding section, removed CPT codes 81402, 81403; added CPT codes 81279, 81338, 81339. Updated Policy Guidelines and references. (lpr)
- 9/7/21 Specialty Matched Consultant Advisory Panel review 8/18/2021. No change to policy statement. (lpr)
- 5/17/22 Reviewed by Avalon 1st Quarter 2022 CAB. Medical Director review 4/2022. Under “When Covered” section: added coverage criteria #2 and #4-7. Added CPT codes 81120, 81121, 81236, 81237, 81348, 81479 to Billing/Coding section. Updated policy guidelines and references. (lpr)

Medical policy is not an authorization, certification, explanation of benefits or a contract. Benefits and eligibility are determined before medical guidelines and payment guidelines are applied. Benefits are determined by the group contract and subscriber certificate that is in effect at the time services are rendered. This document is solely provided for informational purposes only and is based on research of current medical literature and review of common medical practices in the treatment and diagnosis of disease. Medical practices and knowledge are constantly changing and BCBSNC reserves the right to review and revise its medical policies periodically.