

## Corporate Medical Policy

# Testing for Mosquito or Tick-Related Infections AHS – G2158

**File Name:** testing\_for\_mosquito\_or\_tick-related\_infections  
**Origination:** 1/1/2019  
**Last CAP Review:** 2/2021  
**Next CAP Review:** 2/2022  
**Last Review:** 2/2021

### Description of Procedure or Service

---

#### I. Policy Description

Arthropod vectors, including mosquitoes, ticks, fleas, and mites, that feed on vertebrate hosts can spread bacteria, protozoa, and viruses during feeding to their susceptible host, resulting in a variety of infections and diseases. Arboviruses (Arthropod-borne viruses) include Zika virus, West Nile virus (WNV), chikungunya virus, dengue virus (DENV), yellow fever virus (YFV), and Colorado tick fever virus (CTF) to name a few. Malaria and babesiosis are both conditions caused by arthropod-borne protozoan parasites, *Plasmodium* and *Babesia*, respectively. Conditions caused by arthropod-borne bacteria include rickettsial diseases, ehrlichiosis, anaplasmosis, and Lyme disease, as well as other *Borrelia*-associated disorders (Calisher, 1994; CDC, 2018a). Isolation, identification, and characterization of these various infections depend on the causative agent. Identification methods may include culture testing, microscopy, and staining techniques; moreover, molecular testing, such as nucleic acid amplification testing (NAAT), and serologic testing, including immunofluorescence antibody assays and enzyme-linked immunosorbent assays (ELISA), can be used for laboratory diagnosis (Miller et al., 2018).

For Lyme disease and testing for *Borrelia burgdorferi*, please see AHS-G2143 Lyme Disease.  
For Zika virus testing, please see AHS-G2133 Zika Virus Risk Assessment.

#### Related Policies

Lyme Disease AHS – G2143  
Zika Virus Risk Assessment AHS – G2133


#### II. Scientific Background

Hematophagous arthropods, such as mosquitoes, ticks, fleas, and mites, can spread opportunistic bacteria, protozoa, and viruses to host organisms when feeding. Numerous outbreaks of arthropod-borne disease have been documented, including plague, an acute febrile disease caused by *Yersinia pestis* through the bite of infected fleas, which resulted in more than 50 million deaths in Europe alone during the “Black Death” outbreak. More than 3000 cases of plague were reported to the World Health Organization (WHO) between 2010 and 2015 with 584 deaths. Today, most cases of plague occur in the Democratic Republic of Congo, Madagascar, and Peru (WHO, 2017).


The Centers for Disease Control and Prevention (CDC) reported a large increase in the number of vector-borne diseases within the United States and its territories between 2004 -2016. More than 640,000 cases were reported during that time; in fact, infections of tick-borne bacteria and protozoa more than doubled from 2004 to 2016. “In the United States, 16 vectorborne diseases are reportable to state and territorial health departments, which are encouraged to report them to the National Notifiable Disease Surveillance System (NNDSS). Among the diseases on the list that are caused by indigenous pathogens are Lyme disease (*Borrelia burgdorferi*); West Nile, dengue and Zika virus diseases; plague (*Yersinia pestis*); and

# Testing for Mosquito or Tick-Related Infections AHS – G2158

spotted fever rickettsioses (e.g., *Rickettsia rickettsii*). Malaria and yellow fever are no longer transmitted in the United States but have the potential to be reintroduced (Rosenberg et al., 2018).” New vector-borne infections are emerging; for example, two unknown, life-threatening RNA viruses spread by ticks have been identified in the U.S. since 2004. Although both tick- and mosquito-borne diseases are increasing across the U.S., the CDC reports that these two vectors are showing different trends. The mosquito-borne diseases are characterized by epidemics; for example, West Nile Virus is essentially limited to the continental U.S. but has spread rapidly since its introduction to New York in 1999, whereas chikungunya and dengue primarily occur within the U.S. territories. On the other hand, the tick-borne disease increase occurs in the continental U.S. and has been a gradual, steady rate increase with Lyme disease comprising 82% of all tick-borne diseases (Rosenberg et al., 2018). Figure 1 and 2 below, taken from Rosenberg et al. (2018), show the reported cases of tickborne and mosquito-borne disease in the United States from 2004-2016.



This figure shows the number of tick-borne diseases in the U.S. states and territories reported to the CDC, 2004-2016 (Rosenberg et al., 2018)



This figure shows the number of mosquito-borne diseases in the U.S. states and territories reported to the CDC, 2004-2016 (Rosenberg et al., 2018)

## ***Rickettsial infections***

Rocky Mountain spotted fever (RMSF) is the most common rickettsial infection in the U.S. with 2553 cases reported to the CDC alone in 2008. RMSF is caused by *Rickettsia rickettsia*, spread in the U.S. predominantly

# Testing for Mosquito or Tick-Related Infections AHS – G2158

by *Dermacentor variabilis* the American dog tick) and *D. andersoni* (the Rocky Mountain wood tick), and can be found throughout North America as well as parts of South America. The Council for State and Territorial Epidemiologists combined RMSF with other rickettsial diseases into the more broad “spotted fever rickettsiosis” designation in 2009. Besides the obligatory tick bite, typical symptoms of RMSF include fever, headache, and rash with the characteristic rash occurring in approximately 90% of patients within five days of illness. If left untreated, RMSF can be fatal but can easily be treated with antimicrobial therapy upon timely diagnosis. Definitive diagnosis of RMSF cannot usually be made via culture because *Rickettsia* cannot be grown in cell-free culture media since they are obligate intracellular bacteria requiring living host cells. RMSF diagnosis can be made via either skin biopsy prior to treatment with antibiotics or through serologic testing using indirect immunofluorescence assays (IFAs). Immunoglobulin G (IgG) antibodies are more specific than immunoglobulin M (IgM) antibodies since the latter can give false-positive results due to cross-reactivity with other bacterial pathogens. A drawback of IFA is that usually it is unreliable for the first five days of infection until antibody levels are high enough for detection. The CDC and major clinical labs do offer a polymerase chain reaction (PCR)-based assay for RMSF (Sexton & McClain, 2020).

Since 2001, thirteen other human rickettsiae belonging to the spotted fever group (SFG) have been identified. All SFGs can cause fever, headache, and myalgia and are arthropod-borne (primarily ticks and mites). Most patients with an SFG display a rash and/or a localized eschar. Rickettsialpox, caused by *R. akari*, is transmitted from the bite of a house mouse mite, usually after mouse extermination programs result in a decrease of the mite’s food supply. Rickettsialpox is typically a relatively mild disease that can resolve itself without treatment within three weeks, but treatment hastens improvement. Rickettsiosis can also be due to infection with *R. parkeri*, *R. amblyommii*, and *Rickettsia* species 364D (also called *R. philipii*). Isolation of SFG rickettsiae is rare in clinical practice due to the difficulty of obtaining culture; consequently, serology, immunologic detection from tissue, and PCR are more often used for diagnosis. Microimmunofluorescent (MIF) antibody tests, enzyme-linked immunosorbent assays (ELISAs), and Western blot immunoassays can be used to detect convalescent IgG and IgM antibodies, but these methods can only be used at least 10-14 days after the onset of illness when antibody concentrations are high enough for detection. PCR is a very specific technique. PCR using tissue samples has higher specificity than whole blood PCR. Immunologic detection from a tissue biopsy requires the use of special laboratory equipment so it is not as frequently used as either the serologic or PCR detection methods (Sexton & McClain, 2019b).

## ***Ehrlichiosis and Anaplasmosis***

Human ehrlichiosis was first reported in 1986, and the causative agent for human granulocytic anaplasmosis, *Anaplasma phagocytophilum*, was identified in 1994. Both ehrlichiosis and anaplasmosis are transmitted from the bite of infected ticks and have similar clinical and laboratory manifestations. Ehrlichiosis can be caused by *Ehrlichia chaffeensis*, *E. ewingii*, and *E. muris*. Typically, patients have a fever within an incubation period of one to two weeks. Other symptoms can include malaise, myalgia, headache, chills, gastrointestinal distress, and cough. Both leukopenia and thrombocytopenia can occur. Diagnosis via culture is extremely difficult. “Until 1995, only two isolates of *E. chaffeensis* had been recovered from humans; in both cases, this process required over 30 days of cultivation. The isolation of *A. phagocytophilum* from three additional patients has been accomplished using a cell culture system derived from human promyelocytic leukemia cells (Sexton & McClain, 2019a). IFA testing for bacteria-specific antibodies is the most common method for diagnosing ehrlichiosis and anaplasmosis, but similar to rickettsiae, ELISA, PCR, and immunochemical tissue staining can be used as well. Unlike rickettsiosis, ehrlichiosis and anaplasmosis can also be detected by the presence of characteristic intraleukocytic morulae in a peripheral blood smear or buffy coat smear (Sexton & McClain, 2016).

## ***Borrelia Infections***

Besides Lyme disease, caused by *Borrelia burgdorferi*, *Borrelia* can cause relapsing fever. Tick-borne relapsing fever (TBRF) in North America is primarily caused by *B. hermsii*, *B. turicatae*, *B. parkeri*, *B. miyamotoi*, and *B. mazzottii*, and louse-borne relapsing fever (LBRF) is an infection by *B. recurrentis* (Barbour, 2020; Miller et al., 2018). The characteristic feature of these infections is the relapsing fever due to cyclical spirochetemia caused by antigenic variation of the spirochetes. Each bout of fever lasts 3 to 12 days with temperatures ranged from 39°C to 43°C (102.2°F to 109.4°F). Visual analysis by Giemsa or Wright staining

# Testing for Mosquito or Tick-Related Infections AHS – G2158

blood smears taken during a febrile episode is common practice. PCR can also be on a variety of samples, including cerebrospinal fluid (CSF), blood, tissue, or even culture medium. Culture and serologic testing are not typically performed on the *Borrelia* that cause TBRF and LBRF due to the lack of facilities that grow these cultures and to cross-reactivity to other antibodies. One exception is using antibodies to the GlpQ protein characteristic of these *Borrelia* species but not to *B. burgdorferi* (Lyme disease) (Barbour, 2018).

## ***Protozoa infections***

Babesiosis is due to primarily *Babesia microti* in the U.S, but *B. divergens* and *B. venatorum* are the primary causative agents of babesiosis in Europe and China, respectively. The incubation period of Babesia depends on the mode of transfection: 1-4 weeks following a tick bite or up to 9 weeks following a blood transfusion, even though incubation periods up to six months have been reported. The most common symptoms of infection include a fever, fatigue, malaise, chills, sweats, headache, and myalgia. Immunocompromised individuals can develop relapsing babesiosis due to an absent or impaired production of antibodies with approximately 20% mortality rate for patients who develop relapsing babesiosis. A majority of patients with babesiosis are also co-infected with other tick-borne bacterial pathogens. “Preferred tools for diagnosis of babesiosis include blood smear for identification of Babesia organisms and polymerase chain reaction (PCR) for detection of Babesia DNA. Serology can be a useful adjunct to blood smear and PCR” (Krause & Vannier, 2019). Serology is not ideal in diagnosing an acute infection since antibody concentrations remain elevated post-recovery.

*Plasmodium falciparum*, *P. vivax*, and *P. ovale* are responsible for malaria. They are spread by the bite of an Anopheles mosquito where their sporozoites infect the liver within one to two hours. Within the hepatocyte, they form merozoites. Upon rupturing into the bloodstream, the merozoites infect red blood cells for trophozoite formation, causing the erythrocytic stage of the life cycle where additional merozoites are released. During this stage of the cycle, the symptoms of malaria, including fever, occur. This process usually takes 12 to 35 days, but clinical manifestations can be delayed in individuals with partial immunity or those who are taking ineffective prophylaxis. Other initial symptoms can include irregular heartbeat, cough, anorexia, gastrointestinal distress, sweating, chills, malaise, arthralgia, and myalgia. Malaria, if left untreated, can also include acidosis, hypoglycemia, severe anemia, renal and hepatic impairment, edema, and death (Breman, 2020). Parasite-based diagnosis may include microscopic examination of blood smears, which can often identify the species of Plasmodium as well as the parasite density, and antigen-based tests. Rapid diagnostic testing (RDT) of the antigens using immunochromatographic methods is available, but the accuracy of the RDT can vary considerably. NAATs can also be used to identify a malarial infection, and NAATs “are typically used as a gold standard in efficacy studies for antimalarial drugs, vaccines, and evaluation of other diagnostic agents” with a “theoretical limit of detection for PCR...estimated at 0.02 to 1 parasite/microL (Hopkins, 2020).”

## ***Viral infections***

Examples of arthropod-borne viruses (arboviruses) include West Nile virus (WNV), dengue, yellow fever virus (YFV), chikungunya, and Colorado tick fever virus. In the United States, WNV is the most common arbovirus reported to the CDC. In 2016, 96% of the reported 2,240 cases of domestic arboviruses were WNV with 61% of the WNV cases reported being neuroinvasive. Neuroinvasive WNV includes meningitis, encephalitis, and acute flaccid paralysis (Burakoff, Lehman, Fischer, Staples, & Lindsey, 2018). In general, most infected individuals are asymptomatic with only 20-40% of infected patients showing any characteristic symptoms of WNV, including fever, headache, malaise, myalgia, anorexia, and rash. Diagnosis of WNV of a symptomatic individual usually occurs with a WNV IgM antibody capture ELISA (MAC-ELISA) assay. A patient with symptoms of a neurologic infection does require a lumbar puncture. Confirmatory testing can include a plaque reduction neutralization test (PRNT). PCR testing is primarily used with immunocompromised patients who have delayed or absent antibody production, patients with a history of prior flavivirus infections, and blood donors who may be asymptomatic (L. Petersen, 2020).

Dengue virus (DENV) infection is a result of being bitten by an infected *Aedes aegypti* or *A. albopictus* mosquito. Four distinct DENV types of *Flavivirus* are known: DENV-1, DENV-2, DENV-3, and DENV-4. DENV is endemic throughout much of the tropical regions of the world, but the only region of the U.S. endemic for DENV is Puerto Rico. The last major outbreak occurred in Puerto Rico in 2010 where 26,766 cases of suspected DENV were reported and 47% of all laboratory tested specimen were positive (CDC, 2013, 2014).

# Testing for Mosquito or Tick-Related Infections AHS – G2158

“Dengue fever...is an acute febrile illness defined by the presence of fever and two or more of the following but not meeting the case definition of dengue hemorrhagic fever: headache, retro-orbital or ocular pain, myalgia and/or bone pain, arthralgia, rash, hemorrhagic manifestations...[and] leukopenia. The cardinal feature of dengue hemorrhagic fever is plasma leakage due to increased vascular permeability as evidenced by hemoconcentration ( $\geq 20$  percent rise in hematocrit above baseline), pleural effusion, or ascites. DHF [dengue hemorrhagic fever] is also characterized by fever, thrombocytopenia, and hemorrhagic manifestations.... (Thomas, Rothman, Srikiatkachorn, & Kalayanarooj, 2019).” Laboratory diagnostic testing includes direct detection of viral components in serum or indirect serologic assays. “Detection of viral nucleic acid or viral antigen has high specificity but is more labor intensive and costly; serology has lower specificity but is more accessible and less costly” (Thomas, Rothman, Srikiatkachorn, & Kalayanarooj, 2018). Culture testing as a diagnostic tool usually is time-prohibitive.

Colorado tick fever virus (CTFV) is a *Reoviridae* transmitted primarily by the Rocky Mountain wood tick (*Dermacentor andersoni*) in the western US and Canada. Transmission of CTFV has also been reported in blood transfusions. The incubation period can last up to 14 days, and symptoms include fever, headache, chills, myalgia, leukopenia, and prostration. Only 15% of symptomatic patients demonstrate a rash. Serologic tests are usually not helpful until at least 10-14 days for antibody production whereas RT-PCR can be used on the first day of symptoms (L. R. Petersen, 2019).

Yellow fever, occurring primarily in sub-Saharan Africa and South America, is a flavivirus spread by mosquitoes that causes hemorrhagic fever with a high fatality rate. An outbreak in Brazil in January-March 2018 resulted in 4 of 10 patients infected with YFV dying. None of those showing symptoms had been vaccinated against YFV. Yellow fever causes hemorrhagic diathesis due to decreased synthesis of vitamin K-dependent coagulation factors as well as hepatic dysfunction, renal failure, and coagulopathy. Yellow fever diagnosis is typically made by a serologic test using an ELISA-IgM assay; however, this assay does cross-react with other flaviviruses and with the YFV vaccination. Rapid diagnostic testing using either PCR or immunoassay is available. Viral isolation and culture can be performed, but it requires inoculation of mosquitoes or mammalian cell culture. Tissue biopsy, such as liver, cannot be performed on the living patient due to possible fatal hemorrhaging; biopsy would be performed during the post-mortem workup (Ryan, 2019).

Chikungunya virus, endemic in many tropical and subtropical regions of the world, is transmitted by the mosquitoes *Aedes aegypti* and *Aedes albopictus*. Within the U.S., chikungunya is prevalent in Puerto Rico where approximately 25% of blood donors were seropositive; it has also been reported in Florida. Both dengue and Zika are transmitted by the same vectors, so these viruses often co-circulate in a given geographic area. Chikungunya can cause acute febrile polyarthralgia and arthritis. The predominant testing method for diagnosis of chikungunya is the detection of viral RNA via either RT-PCR or virus serology using either ELISA or IFA. Viral culture is typically not used as a diagnostic tool but is used for epidemiologic research (Wilson & Lenschow, 2020).

## Types of Testing

Test	Description	Rationale
<b>Culture</b>	Culture growth depends on the pathogen being studied. If the pathogen is an obligate intracellular organism, then it must be isolated using more sophisticated cell culture techniques. In many circumstances, culture is used for research and/or epidemiology rather than as a diagnostic tool (Biggs et al., 2016; Miller et al., 2018).	At times, culture testing is not as sensitive as either NAAT or serologic testing and can be time-intensive when treatment should not be delayed. Depending on the organism, this may require high biosafety level laboratory for culture growth (Biggs et al., 2016).

# Testing for Mosquito or Tick-Related Infections AHS – G2158

<b>Indirect immunofluorescence antibody (IFA) assays</b>	IFA is a serologic assay that can be used to test for the presence of antibodies, such as IgG and IgM, reactive against the pathogen (Biggs et al., 2016).	Depending on the pathogen, IFA can be a useful tool. At times, though, it can cross-react with either a prior vaccination or infection (Ryan, 2018). An acute infection can often be determined by performing IFA in both the acute phase and convalescent phase where at least a fourfold increase in antibodies is indicative of an acute infection (Biggs et al., 2016).
<b>Darkfield microscopy</b>	Darkfield microscopy can be used to detect the presence of microorganisms, such as motile spirochetes (Miller et al., 2018).	This technique is not widely available, and transport of sample must be done immediately if testing of motile specimen is desired (Miller et al., 2018).
<b>Blood-smear microscopy</b>	Blood-smear microscopy can be either thick or thin and is typically performed on a sample stained with an eosin-azure-type dye, such as Giemsa, to look at intracellular structures or morphological features (Biggs et al., 2016).	This technique should be performed by an experienced microscopist since it can be inconsistent. As compared to other techniques, this technique is relatively inexpensive (Biggs et al., 2016).
<b>Nucleic acid amplification testing (NAAT)</b>	NAATs can include polymerase chain reaction (PCR), real-time PCR (RT-PCR), or other enzyme-dependent amplification testing for the presence of nucleic acids (DNA or RNA).	NAATs can be specific and sensitive; however, they may not be available at all laboratories and/or can be costly. Some NAATs are available as rapid diagnostic tools. NAATs have been used on serum, whole blood, tissue, CSF, and even formalin-fixed, paraffin-embedded biopsies from autopsy tissues. The sensitivity of the technique can vary depending on the sample; for example, whole blood PCR for <i>R. rickettsii</i> is less sensitive than a similar sample test for <i>E. chaffeensis</i> (Biggs et al., 2016).

## **Analytical and Clinical Validity**

The use of antibodies to detect and diagnose arthropod-associated infections and diseases is a common practice. Johnson, Martin, Karabatsos, and Roehrig (2000) first reported the use of monoclonal antibody-based capture ELISA testing for a variety of alphaviruses, including chikungunya, flaviviruses, including dengue and yellow fever, and bunyaviruses. The researchers concluded, “IgG ELISA results correlated with those of the standard plaque-reduction neutralization assays. As expected, some test cross-reactivity was encountered within the individual genera, and tests were interpreted within the context of these reactions. The tests were standardized for laboratory diagnosis of arboviral infections, with the intent that they be used in tandem with the corresponding IgM antibody-capture ELISAs” (Johnson et al., 2000). Kalish and associates also demonstrated



# Testing for Mosquito or Tick-Related Infections AHS – G2158

that IgG and/or IgM antibody responses can still occur up to 20 years post-infection; consequently, a rise in antibody titer does not necessarily indicate a current, acute infection (Kalish et al., 2001).

In 2013, Kato and colleagues tested the sensitivity of two different RT-PCR-based assays for Rickettsia—PanR8, an assay that tests for Rickettsia in general, and RRI6, an assay specific for *R. rickettsii*. Both of these methods were more sensitive in testing for Rickettsia than the nested PCR method of the CDC; moreover, both of these methods are faster than the nested PCR method (1 hr versus 1-2 days, respectively) (Kato et al., 2013). These results were corroborated in 2014 by Denison and colleagues. They used a multiplex PCR assay to correctly identify all cell controls for *R. rickettsii*, *R. parkeri*, and *R. akari*; moreover, no false-positive results were reported using this methodology. “This multiplex real-time PCR demonstrates greater sensitivity than nested PCR assays in FFPE [formalin-fixed, paraffin-embedded] tissues and provides an effective method to specifically identify cases of Rocky Mountain spotted fever, rickettsialpox, and *R. parkeri* rickettsiosis by using skin biopsy specimens” (Denison, Amin, Nicholson, & Paddock, 2014).

The FDA has approved the use of the BinaxNOW malaria test for screening and diagnosing malaria. Even though, this testing method is considerably faster than other methods (as low as 1.1-1.7 hours complete turnaround time (Ota-Sullivan & Blecker-Shelly, 2013)), the use of BinaxNOW in non-endemic areas is a point of controversy due to relatively low sensitivity (84.2%) and for misclassifying Plasmodium falciparum malaria as non-falciparum (Dimaio, Pereira, George, & Banaei, 2012). Moreover, it has been reported that Salmonella typhi can give a false-positive for malaria using the BinaxNOW test (Meatherall, Preston, & Pillai, 2014).

Akoolo et al. (2017) compared qPCR results in the detection of Babesia infection against currently available non-NAAT tests (FISH and microscopy). Blood samples were analyzed from 192 patients. The researchers report that “Of 28 samples that were positive by FISH, 27 (96%) were also positive by qPCR indicating high congruency between nucleic acid based tests. Interestingly, of 78 asymptomatic samples not tested by FISH, 22 were positive by our qPCR” (Akoolo et al., 2017). Overall, the qPCR method was found to have a sensitivity of 96.2% and a specificity of 70.5%. The authors conclude, “Robust qPCR using specific probes can be highly useful for efficient and appropriate diagnosis of babesiosis in patients in conjunction with conventional diagnostics, or as a stand-alone test, especially for donated blood screening” (Akoolo et al., 2017).

### III. State and Federal Regulations, as applicable

On 6/29/2017, the FDA approved the Rickettsia Real-Time PCR Assay (K170940) by the Centers for Disease Control and Prevention (CDC) with the following definition: “An in vitro diagnostic test for the detection of Rickettsia spp. nucleic acids in specimens from individuals with signs or symptoms of rickettsial infection and epidemiological risk factors consistent with potential exposure. Test results are used in conjunction with other diagnostic assays and clinical observations to aid in the diagnosis infection, in accordance with criteria defined by the appropriate public health authorities in the Federal government (FDA, 2018).”

On 9/1/2009, the FDA approved the BinaxNOW Malaria Positive Control Kit (K083744) rapid diagnostic test (RDT), an in vitro qualitative immunochromatographic assay, for use by hospital and commercial laboratories, but it is not approved for individual or physician offices (Arguin & Tan, 2017; FDA, 2018).

As of 8/7/2018, the FDA has approved the following assays for the detection of West Nile Virus (FDA, 2018): West Nile Virus ELISA IgG model EL0300G and West Nile Virus IgM Capture ELISA model EL0300M by Focus Technologies, Inc., West Nile Virus IgM Capture ELISA model E-WNV02M and West Nile Virus IgG Indirect ELISA by Panbio Limited, West Nile Detect IgM ELISA by Inbios Intl, Inc., Spectral West Nile Virus IgM Status Test by Spectral Diagnostics, Inc., and the EUROIMMUN Anti-West Nile Virus ELISA (IgG) and EUROIMMUN Anti-West Nile Virus ELISA (IgM) by Euroimmun US, Inc.

Additionally, many labs have developed specific tests that they must validate and perform in house. These laboratory-developed tests (LDTs) are regulated by the Centers for Medicare and Medicaid (CMS) as high-complexity tests under the Clinical Laboratory Improvement Amendments of 1988 (CLIA '88). As an LDT, the

# Testing for Mosquito or Tick-Related Infections AHS – G2158

U. S. Food and Drug Administration has not approved or cleared this test; however, FDA clearance or approval is not currently required for clinical use.

**\*\*\*Note: This Medical Policy is complex and technical. For questions concerning the technical language and/or specific clinical indications for its use, please consult your physician.**

## Policy

---

**BCBSNC will provide coverage for testing for mosquito or tick-related infections when it is determined the medical criteria or reimbursement guidelines below are met.**

## Benefits Application

---

This medical policy relates only to the services or supplies described herein. Please refer to the Member's Benefit Booklet for availability of benefits. Member's benefits may vary according to benefit design; therefore member benefit language should be reviewed before applying the terms of this medical policy.

## When testing for mosquito or tick-related infections is covered

---

1. Concerning suspected cases of rickettsial diseases (see signs and symptoms below), including Rocky Mountain spotted fever, *Rickettsia parkeri* rickettsiosis, *Rickettsia* species 364D rickettsiosis, *Rickettsia spp* (mild spotted fever), and *R. akari* (rickettsialpox):
  - a. Reimbursement is allowed for the following:
    - i. Indirect immunofluorescence antibody (IFA) assay for IgG antibodies
    - ii. A limit of one unit of IFA assay meets coverage criteria.
2. Concerning suspected cases of ehrlichiosis and/or anaplasmosis (see signs and symptoms below):
  - a. Reimbursement is allowed for the following:
    - i. NAAT, including PCR, of whole blood; AND
    - ii. Indirect immunofluorescence antibody (IFA) assay for IgG antibodies; AND
    - iii. Microscopy for morulae detection
3. Concerning suspected cases of tick-borne relapsing fever (TBRF) caused by *Borrelia hermsii*, *B. parkerii*, *B. mazzottii*, or *B. turicatae* (see signs and symptoms below):
  - a. Reimbursement is allowed for the following:
    - i. Dark-field microscopy of peripheral blood smear; OR
    - ii. Microscopy of Wright- or Giemsa-stained blood smear; OR
    - iii. PCR testing
4. Concerning suspected cases of babesiosis (see signs and symptoms below):
  - a. Reimbursement is allowed for the following:
    - i. Giemsa- or Wright-stain of blood smear
    - ii. NAAT, including PCR
5. Concerning suspected cases of malaria (see signs and symptoms below):
  - a. Reimbursement is allowed for the following:
    - i. Smear microscopy to diagnose malaria, determine the species of *Plasmodium*, identify the parasitic life-cycle stage, and/or quantify the parasitemia  
Can be repeated up to three times within three days if initial microscopy is negative in suspected cases of malaria; OR
    - ii. Rapid immunochromatographic diagnostic test
6. Concerning suspected cases of chikungunya virus (see signs and symptoms below):



# Testing for Mosquito or Tick-Related Infections AHS – G2158

- a. Reimbursement is allowed for the following:
  - i. Viral culture for diagnosis; OR
  - ii. NAAT, including PCR, for presence of chikungunya in serum sample; OR
  - iii. Indirect immunofluorescence antibody (IFA) assay for IgM antibodies during both the acute and convalescent phases
7. Concerning suspected cases of West Nile Virus (WNV) (see signs and symptoms below):
  - a. Reimbursement is allowed for the following:
    - i. IFA for WNV-specific IgM or IgG antibodies in either serum or CSF; AND
    - ii. Plaque reduction neutralization test for WNV
8. Concerning suspected cases of Yellow Fever Virus (YFV) (see signs and symptoms below):
  - a. Reimbursement is allowed for the following:
    - i. Plaque reduction neutralization test for YFV; AND
    - ii. NAAT, including PCR, for YFV; AND
    - iii. Serologic assays to detect virus-specific IgM and IgG antibodies
9. Concerning suspected cases of Dengue virus (DENV) (see signs and symptoms below):
  - a. Reimbursement is allowed for the following:
    - i. Plaque reduction neutralization test for DENV; AND
    - ii. NAAT, including PCR, for DENV; OR
    - iii. IgM antibody capture ELISA (MAC-ELISA); OR
    - iv. NS1 ELISA
10. Concerning suspected cases of Colorado Tick Fever (CTF) (see signs and symptoms below):
  - a. Reimbursement is allowed for the following:
    - i. Virus-specific IFA-stained blood smears; OR
    - ii. IFA for CTF-specific antibodies

## **When testing for mosquito or tick-related infections is not covered**

1. Concerning suspected cases of rickettsial diseases (see signs and symptoms below), including Rocky Mountain spotted fever, *Rickettsia parkeri* rickettsiosis, *Rickettsia* species 364D rickettsiosis, *Rickettsia spp* (mild spotted fever), and *R. akari* (rickettsialpox):
  - a. Reimbursement is not allowed for the following:
    - i. Standard blood culture; OR
    - ii. Nucleic acid amplification testing (NAAT), including PCR; OR
    - iii. IFA assay for IgM antibodies;
    - iv. More than one unit of IFA testing
2. Concerning suspected cases of ehrlichiosis and/or anaplasmosis (see signs and symptoms below):
  - a. Reimbursement is not allowed for the following:
    - i. IFA assay for IgM antibodies; OR
    - ii. Standard blood culture
3. Concerning suspected cases of tick-borne relapsing fever (TBRF) caused by *Borrelia hermsii*, *B. parkerii*, *B. mazzottii*, or *B. turicatae* (see signs and symptoms below):
  - a. Reimbursement is not allowed for the following:
    - i. IFA assay for either IgG or IgM for *Borrelia*; OR
    - ii. Culture testing for *Borrelia*
4. Concerning suspected cases of babesiosis (see signs and symptoms below):
  - a. Reimbursement is not allowed for the following:
    - i. IFA assay for either Babesia IgG or IgM
5. Concerning suspected cases of malaria (see signs and symptoms below):
  - a. Reimbursement is not allowed for the following:
    - i. NAAT, including PCR

# Testing for Mosquito or Tick-Related Infections AHS – G2158

- ii. IFA for *Plasmodium* antibodies
- 6. Concerning suspected cases of West Nile Virus (WNV) (see signs and symptoms below):
  - a. Reimbursement is not allowed for the following:
    - i. NAAT, including PCR, for WNV
- 7. Concerning suspected cases of Dengue virus (DENV) (see signs and symptoms below):
  - a. Reimbursement is not allowed for the following:
    - i. IgG-ELISA
    - ii. Hemagglutination testing

## Policy Guidelines

---

1. Typical signs and symptoms of rickettsial diseases usually begin 3 – 12 days after initial bite and can include (Biggs et al., 2016):
  - a. Fever
  - b. Headache
  - c. Chills
  - d. Malaise
  - e. Myalgia
  - f. Nausea
  - g. Vomiting
  - h. Abdominal pain
  - i. Photophobia
  - j. Anorexia
  - k. Skin rash
  - l. Ulcerative lesion with regional lymphadenopathy (for *Rickettsia* species 364D rickettsiosis)
2. Typical signs and symptoms of ehrlichiosis and/or anaplasmosis usually begin 5-14 days after an infected tick bite, and they include (Biggs et al., 2016):
  - a. Fever
  - b. Headache
  - c. Malaise
  - d. Myalgia
  - e. Shaking chills
  - f. Gastrointestinal issues, including nausea, vomiting, and diarrhea, in ehrlichiosis
3. Typical signs and symptoms of tick-borne relapsing fever include (CDC, 2018f):
  - a. Recurring febrile episodes that last approximately 3 days separated by approximately 7 days
  - b. Nonspecific symptoms that occur in at least 50% of cases include headache, myalgia, chills, nausea, arthralgia, and vomiting
4. Typical signs and symptoms of babesiosis can include (CDC, 2019a):
  - a. Hemolytic anemia
  - b. Splenomegaly
  - c. Hepatomegaly
  - d. Jaundice
  - e. Nonspecific flu-like symptoms such as fever, chills, body aches, weakness, and fatigue
5. Typical signs and symptoms of malaria can include (Arguin & Tan, 2019):
  - a. Fever
  - b. Influenza-like symptoms such as chills, headache, body aches, and so on
  - c. Anemia
  - d. Jaundice

# Testing for Mosquito or Tick-Related Infections AHS – G2158

- e. Seizures
  - f. Mental confusion
  - g. Kidney failure
  - h. Acute respiratory distress syndrome
6. Typical signs and symptoms of chikungunya include (Staples, Hill, & Powers, 2020):
- a. High fever (>102°F or 39°C)
  - b. Joint pains (usually multiple joints, bilateral, and symmetric)
  - c. Headache
  - d. Myalgia
  - e. Arthritis
  - f. Conjunctivitis
  - g. Nausea
  - h. Vomiting
  - i. Maculopapular rash
7. Typical signs and symptoms of WNV include (Nasci et al., 2013):
- a. Headache
  - b. Myalgia
  - c. Arthralgia
  - d. Gastrointestinal symptoms
  - e. Maculopapular rash
  - f. Less than 1% develop neuroinvasive WNV with symptoms of meningitis, encephalitis, or acute flaccid paralysis
8. Typical signs and symptoms of yellow fever include (Gershman & Staples, 2019):
- a. Nonspecific influenza like syndrome including fever, chills, headache, backache, myalgia, prostration, nausea, and vomiting in initial illness
  - b. Toxic form of disease symptoms includes jaundice, hemorrhagic symptoms, and multisystem organ failure
9. Typical signs and symptoms of dengue can include (CDC, 2014):
- a. Fever
  - b. Headache
  - c. Retro-orbital eye pain
  - d. Myalgia
  - e. Arthralgia
  - f. Erythematous maculopapular rash
  - g. Petechiae
  - h. Leukopenia
10. Typical signs and symptoms of CTF can include (CDC, 2015a):
- a. Fever
  - b. Chills
  - c. Headache
  - d. Myalgia
  - e. Malaise
  - f. Sore throat
  - g. Vomiting
  - h. Abdominal Pain
  - i. Maculopapular or petechial rash

Note: For Lyme disease and testing for *Borrelia burgdorferi*, please see AHS-G2143 Lyme Disease. For Zika virus testing, please see AHS-G2133 Zika Virus Risk Assessment.

# Testing for Mosquito or Tick-Related Infections AHS – G2158

## Guidelines and Recommendations

Centers for Disease Control and Prevention (CDC) (Arguin & Tan, 2019; Biggs et al., 2016; CDC, 2018a, 2018b, 2019a, 2019c; Gershman & Staples, 2019; Nasci et al., 2013; Staples et al., 2020)

*Diagnosis and Management of Tickborne Rickettsial Diseases* (Biggs et al., 2016): In 2016, the CDC released their guidelines and recommendations concerning Rickettsial diseases, including Rocky Mountain spotted fever, in the MMWR. The table below summarizes their recommended diagnostic tests for tickborne rickettsial diseases:

TABLE 4. Recommended diagnostic tests for tickborne rickettsial diseases

Disease	PCR			Microscopy for morulae detection	IFA assay for IgG antibodies (acute and convalescent)*
	Whole blood	Eschar biopsy or swab	Rash biopsy		
Rocky Mountain spotted fever	Yes†	—	Yes	—	Yes
<i>Rickettsia parkeri</i> rickettsiosis	—	Yes	Yes	—	Yes
<i>Rickettsia species 364D</i> rickettsiosis	—	Yes	—	—	Yes
<i>Ehrlichia chaffeensis</i> ehrlichiosis (human monocytic ehrlichiosis)	Yes	—	—	Yes	Yes
<i>Ehrlichia ewingii</i> ehrlichiosis	Yes	—	—	Yes	Yes
<i>Ehrlichia muris</i> -like agent ehrlichiosis	Yes	—	—	—	Yes
Human anaplasmosis (human granulocytic anaplasmosis)	Yes	—	—	Yes	Yes

Abbreviations: IFA = indirect immunofluorescence antibody; IgG = immunoglobulin G; PCR = polymerase chain reaction.

\* IFA assay is insensitive during the first week of illness for most tickborne rickettsial diseases; a sample should be collected during this interval (acute specimen), and a second sample should be collected 2-4 weeks later (convalescent specimen) for comparison. Elevated titers alone are not sufficient to diagnose infection with tickborne rickettsial diseases; serial titers are needed for confirmation. Demonstration of at least a fourfold rise in antibody titer is considered confirmatory evidence of acute infection.

† PCR of whole blood samples for *Rickettsia rickettsii* has low sensitivity; sensitivity increases in patients with severe disease.

To summarize their recommendations, even though indirect immunofluorescence antibody assays (IFAs) are insensitive typically during the first week of an acute infection, they are the standard reference for tickborne rickettsial infections and that a minimum of two tests are to be performed for diagnosis. Usually, one sample is taken early after the initial symptoms are present, and a second sample is taken 2-4 weeks later. A minimum of a fourfold rise in antibody titer is required to confirm diagnosis. In cases of ehrlichiosis and anaplasmosis, during the first week, PCR amplification can be used on whole blood for diagnosis, but PCR has low sensitivity in Rocky Mountain spotted fever except in patients with severe disease. Morulae detection via either blood-smear or buffy-coat preparation microscopy can also be indicative of ehrlichiosis or anaplasmosis. “Rickettsiae cannot be isolated with standard blood culture techniques because they are obligate intracellular pathogens; specialized cell culture methods are required. Because of limitations in availability and facilities, culture is not often used as a routine confirmatory diagnostic method for tickborne rickettsial diseases” (Biggs et al., 2016).

*Tick-borne relapsing fever (TBRF)* (CDC, 2018f): In the U.S., TBRF can be caused by *Borrelia hermsii*, *B. parkerii*, and *B. turicatae* with *B. hermsii* being the most common causative agent. TBRF often presents with a relapsing nature (usually ~3 days per febrile episode followed by an afebrile period of approximately one week). “Spirochetemia (spirochetes in blood) in TBRF patients often reaches high concentrations (>10<sup>6</sup> spirochetes/ml). Thus, microscopy is a useful diagnostic tool for TBRF. The diagnosis of TBRF may be based on direct microscopic observation of relapsing fever spirochetes using dark field microscopy or stained peripheral blood smears. Spirochetes are more readily detected by microscopy in symptomatic, untreated patients early in the course of infection. Other bacteria, such as *Helicobacter*, may appear morphologically similar, so it is important to consider clinical and geographical characteristics of the case when making a diagnosis of TBRF based on microscopy. Additional testing, such as serology or culture, is recommended.”

“Serologic testing for TBRF is not standardized and results may vary by laboratory. Serum taken early in infection may be negative, so it is important to also obtain a serum sample during the convalescent period (at least 21 days after symptom onset). A change in serology results from negative to positive, or the development of an IgG response in the convalescent sample, is supportive of a TBRF diagnosis. However, early antibiotic

# Testing for Mosquito or Tick-Related Infections AHS – G2158

treatment may limit the antibody response. Patients with TBRF may have false-positive tests for Lyme disease because of the similarity of proteins between the causative organisms. A diagnosis of TBRF should be considered for patients with positive Lyme disease serology who have not been in areas endemic for Lyme disease. Incidental laboratory findings include normal to increased white blood cell count with a left shift towards immature cells, a mildly increased serum bilirubin level, mild to moderate thrombocytopenia, elevated erythrocyte sedimentation rate (ESR), and slightly prolonged prothrombin time (PT) and partial thromboplastin time (PTT) (CDC, 2018f).”

*Colorado Tick Fever (CTF) (CDC, 2018d):* As of 2015, CTF was reportable in Arizona, Colorado, Montana, Oregon, Utah, and Wyoming. “Laboratory diagnosis of CTF is generally accomplished by testing of serum to detect viral RNA or virus-specific immunoglobulin (Ig) M and neutralizing antibodies. Antibody production can be delayed with CTF, so tests that measure antibodies may not be positive for 14–21 days after the onset of symptoms. RT-PCR (reverse-transcriptase polymerase chain reaction) is a more sensitive test early in the course of disease. CTF testing is available at some commercial and state health department laboratories and at CDC. Contact your state or local health department for assistance with diagnostic testing. They can help you determine if samples should be sent to the CDC Arbovirus Diagnostic Laboratory for further testing (CDC, 2018d).”

*Babesiosis (CDC, 2018c, 2019a):* According to the CDC website, the most recent update about babesiosis for health professionals is from 2012 (with revision in 2018). Diagnosis can be challenging due to the nonspecific clinical manifestations of the disease. “For acutely ill patients, the findings on routine laboratory testing frequently include hemolytic anemia and thrombocytopenia. Additional findings may include proteinuria, hemoglobinuria, and elevated levels of liver enzymes, blood urea nitrogen, and creatinine. If the diagnosis of babesiosis is being considered, manual (non-automated) review of blood smears should be requested explicitly. In symptomatic patients with acute infection, *Babesia* parasites typically can be detected by light-microscopic examination of blood smears, although multiple smears may need to be examined. Sometimes it can be difficult to distinguish between *Babesia* and *Plasmodium* (especially *P. falciparum*) parasites and even between parasites and artifacts (such as stain or platelet debris). Consider having a reference laboratory confirm the diagnosis—by blood-smear examination and, if indicated, by other means, such as molecular and/or serologic methods tailored to the setting/species (CDC, 2018c, 2019a).”

*Malaria (Arguin & Tan, 2017, 2019):* The CDC considers smear microscopy as the gold standard in diagnosing malaria since it can determine the species, identify the stage of parasitic life-cycle, and quantify the parasitemia. The CDC states, “Blood smear microscopy remains the most important method for malaria diagnosis. Microscopy can provide immediate information about the presence of parasites, allow quantification of the density of the infection, and allow determination of the species of the malaria parasite—all of which are necessary for providing the most appropriate treatment. Microscopy results should ideally be available within a few hours. These tests should be performed immediately when ordered by a health care provider. They should not be saved for the most qualified staff to perform or batched for convenience. In addition, these tests should not be sent out to reference laboratories with results available only days to weeks later. Assistance with speciation of malaria on smears is available from CDC” (Arguin & Tan, 2019). The CDC also notes that rapid diagnostic tests (RDTs) for malaria can detect malaria parasitic antigens. However, “RDTs offer a useful alternative to microscopy in situations where reliable microscopic diagnosis is not immediately available. Although RDTs can detect malaria antigens within minutes, they have several limitations. RDTs cannot distinguish between all of the *Plasmodium* species that affect humans, they may be less sensitive than expert microscopy or PCR for diagnosis, they cannot quantify parasitemia, and an RDT-positive test result may persist for days or weeks after an infection has been treated and cleared. Thus, RDTs are not useful for assessing response to therapy. Both positive and negative RDT results must always be confirmed by microscopy. Microscopy confirmation of the RDT result should occur as soon as possible because the information on the presence, density, and parasite species is critical for optimal management of malaria” (Arguin & Tan, 2019). Regarding PCR, the CDC states that “Although these tests are more sensitive than routine microscopy, results are not usually available as quickly as microscopy results, thus limiting the utility of this test for acute diagnosis and initial clinical management. Use of PCR testing is encouraged to confirm the species of malaria parasite and detect mixed infections” (Arguin & Tan, 2019).

The CDC also provided an update to malaria diagnosis in 2018. Although microscopy remained the “gold standard” for confirmation of malaria, other tests such as RDTs and PCR-based tests remained useful in certain

# Testing for Mosquito or Tick-Related Infections AHS – G2158

situations (namely if microscopy is unavailable). PCR is considered most useful for “confirming the species of malarial parasite after the diagnosis has been established by either smear microscopy or RDT.” Finally, the CDC recommends that all cases of malaria be evaluated for drug resistance, typically through molecular characterization (PCR, gene sequencing) or in vitro tests (CDC, 2018e).

*Chikungunya* (Staples, Hills, & Powers, 2017; Staples et al., 2020): In the CDC Yellow Book, concerning the chikungunya virus, they recommend that “the differential diagnosis of chikungunya virus infection depends on the clinical signs and symptoms as well as where the person was suspected of being infected.” The other diseases to consider include: Zika, malaria, leptospirosis, parvovirus, group A *Streptococcus*, rubella, measles, dengue, enterovirus, adenovirus, alphavirus infections, post-infectious arthritis, and rheumatic conditions. Laboratory diagnosis is done by serum testing for detection of virus, viral nucleic acids, or virus-specific IgM and neutralizing antibodies. “During the first week after onset of symptoms, chikungunya can often be diagnosed by performing viral culture or nucleic acid amplification on serum. Virus-specific IgM and neutralizing antibodies normally develop toward the end of the first week of illness. Therefore, to definitively rule out the diagnosis, convalescent-phase samples should be obtained from patients whose acute-phase samples test negative. Testing for chikungunya virus is performed at CDC, several state health department laboratories, and several commercial laboratories (Staples et al., 2017, 2020).”

*West Nile Virus (WNV)* (Nasci et al., 2013): “WNV infections are most frequently confirmed by detection of anti-WNV immunoglobulin (Ig) M antibodies in serum or cerebrospinal fluid (CSF). The presence of anti-WNV IgM is usually good evidence of recent WNV infection, but may indicate infection with another closely related flavivirus (e.g., St. Louis encephalitis). Because anti-WNV IgM can persist in some patients for >1 year, a positive test result occasionally may reflect past infection unrelated to current disease manifestations. Serum collected within 8 days of illness onset may lack detectable IgM, and the test should be repeated on a convalescent-phase sample. IgG antibody generally is detectable shortly after the appearance of IgM and persists for years. Plaque-reduction neutralization tests (PRNT) can be performed to measure specific virus-neutralizing antibodies. A fourfold or greater rise in neutralizing antibody titer between acute- and convalescent-phase serum specimens collected 2 to 3 weeks apart may be used to confirm recent WNV infection and to discriminate between cross-reacting antibodies from closely related flaviviruses.” NAAT may not be suitable in most cases since the concentrations of WNV RNA are so low by the time a patient begins to show symptoms of infection; however, NAAT may be suitable in immunocompromised individuals who have either delayed or absent antibody development.

*Yellow Fever Virus (YFV)* (Gershman & Staples, 2019): Isolation of the virus or NAAT should be performed as early as possible in suspected cases of YFV. “By the time more overt symptoms are recognized, the virus or viral RNA may no longer be detectable. Therefore, virus isolation and nucleic acid amplification should not be used to rule out a diagnosis of yellow fever... Serologic assays to detect virus-specific IgM and IgG antibodies (sic). Because of cross-reactivity between antibodies raised against other flaviviruses, more specific antibody testing, such as a plaque reduction neutralization test, should be done to confirm the infection (Gershman & Staples, 2019).” Since YFV is a nationally notifiable disease, clinicians should contact their state and/or local health departments according to their respective local, state, and/or federal guidelines.

*Dengue* (CDC, 2019b, 2020): Diagnosis of dengue can be via isolation of virus, serological tests such as immunoassays, and molecular methods, including RT-PCR. The CDC recommends testing symptomatic individuals that have recently traveled to or lives in areas where dengue is transmitted, as well as symptomatic pregnant women with possible dengue or Zika exposure. The CDC’s testing algorithm for dengue is as follows: “Patients with symptoms consistent with dengue can be tested with both molecular and serologic diagnostic tests during the first 7 days of illness. After the first 7 days of illness, test only with serologic diagnostic tests” (CDC, 2020).

Finally, the CDC does not recommend testing asymptomatic patients or preconception screening for dengue (CDC, 2019b, 2020).

**Infectious Diseases Society of America (IDSA) and the American Society for Microbiology (ASM) (Miller et al., 2018)**

# Testing for Mosquito or Tick-Related Infections AHS – G2158

*Laboratory Diagnosis of Tick-borne Infections:* The information given below outlines the diagnostic procedures for tick-borne infections and is taken from Table 47 of the 2018 IDSA/ASM guidelines.

Bacteria		
Etiologic Agents	Diagnostic Procedures	Optimum Specimens
<p>Relapsing fever borreliae</p> <p><i>Borrelia hermsii</i> (western US)</p> <p><i>Borrelia parkeri</i> (western US)</p> <p><i>Borrelia turicatae</i> (southwestern US)</p> <p><i>Borrelia mazzottii</i> (southern US)</p>	<p>Primary test: Darkfield microscopy or Wright, Giemsa, or Diff-Quik stains of peripheral thin or/ and thick blood smears. Can be seen in direct wet preparation of blood in some cases.</p> <p>Other testing: NAAT, Culture, Serologic testing</p>	<p>Blood or bone marrow for the primary test. Blood or body fluids for NAAT. Serum for culture or serologic testing.</p>
<p><i>Borrelia miyamotoi</i> (<i>B. miyamotoi</i> infection, hard tick-borne relapsing fever)</p>	<p>Primary test: NAAT</p> <p>Serology: EIA for detection of antibodies to recombinant GlpQ antigen</p>	<p>Blood for NAAT and serum for serology.</p>
<p><i>Anaplasma phagocytophilum</i> (human granulocytotropic anaplasmosis)</p>	<p>Primary test: NAAT</p> <p>Alternate Primary (if NAAT is unavailable): Wright or Giemsa stain of peripheral blood or buffy coat leukocytes during week first week of infection.</p> <p>Serology: Acute and convalescent IFA titers for IgG-class antibodies to <i>A. phagocytophilum</i> antibodies</p> <p>Immunohistochemical staining of <i>Anaplasma</i> antigens in formalin-fixed, paraffin-embedded specimens</p>	<p>Blood for both NAAT and Wright or Giemsa stain</p> <p>Serum for Serology</p> <p>Bone marrow biopsies or autopsy tissues (spleen, lymph nodes, liver, and lung) for IHC staining</p>
<p><i>Ehrlichia chaffeensis</i> (human monocytotropic ehrlichiosis)</p> <p><i>Ehrlichia muris</i></p> <p><i>Ehrlichia ewingii</i></p>	<p>Primary test: NAAT (Only definitive diagnostic assay for <i>E. ewingii</i>)</p> <p>Wright or Giemsa stain of peripheral blood or buffy</p>	<p>Whole blood for NAAT</p> <p>Blood for Wright or Giemsa stain</p>



# Testing for Mosquito or Tick-Related Infections AHS – G2158

	<p>coat leukocytes smear during first week of infection</p> <p>Serology: acute and convalescent IFA titers for Ehrlichia IgG-class antibodies</p> <p>Immunohistochemical staining of Ehrlichia antigens in formalin-fixed, paraffin-embedded specimens</p>	<p>Serum for serology</p> <p>Bone marrow biopsies or autopsy tissues (spleen, lymph nodes, liver and lung) for IHC staining</p>
<p><i>Rickettsia rickettsii</i> (RMSF)</p> <p>Other spotted fever group <i>Rickettsia</i> spp (mild spotted fever)</p> <p><i>R. typhi</i> (murine typhus)</p> <p><i>R. akari</i> (rickettsialpox)</p> <p><i>R. prowazekii</i> (epidemic typhus)</p>	<p>Serology: acute and convalescent IFA for <i>Rickettsia</i> sp IgM and IgG antibodies</p> <p>NAAT</p> <p>Immunohistochemical staining of spotted fever group rickettsiae antigens (up to first 24 h after antibiotic therapy initiated) in formalin-fixed, paraffin-embedded specimens</p>	<p>Serum for serology</p> <p>Skin biopsy (preferably a maculopapule containing petechiae or the margin of an eschar) or autopsy tissues (liver, spleen, lung, heart, and brain) for NAAT</p> <p>Skin biopsy (preferably a maculopapule containing petechiae or the margin of an eschar) or autopsy tissues (liver, spleen, lung, heart, and brain) for IHC staining</p>
Protozoa		
Etiologic Agents	Diagnostic Procedures	Optimum Specimens
<p><i>Babesia microti</i></p> <p><i>Babesia</i> spp</p>	<p>Primary test: Giemsa, Wright, Wright-Giemsa stains of peripheral thin and thick blood smears (Giemsa preferred)</p> <p>Primary test for acute infection: NAAT</p> <p>Serology: acute and convalescent IFA titers for Babesia IgG-class antibodies</p> <p><b>NOTE: Not recommended for acute infection.</b></p>	<p>Whole blood (EDTA vacutainer tube is a second choice) for staining</p> <p>Blood for NAAT and serum for serology</p>
Virus		

# Testing for Mosquito or Tick-Related Infections AHS – G2158

Etiologic Agents	Diagnostic Procedures	Optimum Specimens
Colorado tick fever virus	Virus-specific IFA-stained blood smears  Serology: IFA titers or complement fixation	Blood  Serum
Powassan/deer tick virus	Primary test: IgM capture EIA (available only through state departments of public health)  NAAT	Serum for EIA  Blood, CSF, brain (biopsy or autopsy) for NAAT

The IDSA/ASM does note that most PCR-based assays for babesiosis only detect *B. microti* even though there are at least three other species of *Babesia* that can cause the infection. “Real time PCR available from CDC and reference labs... Serology does not distinguish between acute and past infection” (Miller et al., 2018).

Their recommendation for the main diagnostic testing for malaria due to *Plasmodium falciparum*, *P. ovale*, *P. vivax*, *P. malariae*, and *P. knowlesi* is “Stat microscopic examination of Giemsa-stained thick and thin blood films (repeat testing every 12–24 h for a total of 3 exams before ruling out malaria); rapid antigen detection tests followed by confirmatory blood films within 12–24 h.” They make the following special remark: “Antigen tests lack sensitivity with low parasitemia and non-falciparum malaria and do not differentiate all species. PCR from some reference laboratories will detect and differentiate all species. Calculation of percentage parasitemia (using thick or thin blood films) is required for determining patient management and following response to therapy (Miller et al., 2018).” Concerning dengue virus DENV, “Plaque reduction neutralization tests (PRNTs) are considered the reference standard for detection of antibodies to arthropod-borne viruses (arboviruses) and provide improved specificity over commercial serologic assays; however, due to the complexity of testing, PRNT is currently only available at select public health laboratories and the CDC.” They note that false-positives for antibodies to DENV may not necessarily indicate DENV infection since it can also be indicative of a prior flavivirus infection, such as West Nile virus or Zika virus. They do state the following concerning the use of NAAT, “Detection of DENV RNA by NAAT is preferred for acutely ill patients. Recently, detection of the DENV NS1 antigen, which is secreted from infected host cells as early as 1 day after symptom onset and up to 10 days thereafter, has become an acceptable alternative to NAAT for diagnosis of acute DENV infection (Miller et al., 2018).”

For West Nile Virus (WNV), they state: “Laboratory diagnosis of WNV, and most other arboviruses, is typically accomplished by detecting virus-specific IgM- and/or IgG-class antibodies in serum and/or CSF.” Possible false-positives can occur if a patient has been vaccinated against yellow fever or if they have had a previous infection of another flavivirus. They do note that WNV RNA detection via NAAT can be performed on either the serum or CSF for immunosuppressed patients.

## American Academy of Pediatrics 2018 Redbook (AAP, 2018)

*Babesiosis* (AAP, 2018a): “Acute, symptomatic cases of babesiosis typically are diagnosed by microscopic identification of *Babesia* parasites on Giemsa- or Wright-stained blood smears... PCR assay is particularly recommended for use in early infection, when parasites are more difficult to visualize on blood smear. However, PCR assay should be used with caution when monitoring response to therapy, because *B. microti* can be detected for weeks and months after parasites no longer are visualized on blood smear.” They do state that antibody testing can be useful in distinguishing between *Babesia* and *Plasmodium* infections whenever blood smear examinations and travel histories are inconclusive or for detecting individuals with very low levels of parasitemia.

# Testing for Mosquito or Tick-Related Infections AHS – G2158

*Non-Lyme Borrelia Infections (AAP, 2018b)*: Dark-field microscopy and Wright-, Giemsa-, or acridine orange-stained preparations of blood smears can be used to observe the presence of spirochetes in the initial febrile episode, but their presence is more difficult to determine in future recurrences. Both enzyme immunoassay and Western immunoblot analysis can detect serum antibodies; however, “these antibody tests are not standardized and are affected by antigenic variations among and within *Borrelia* species and strains.” As of publication, PCR and antibody-based testing were still under development and were not widely available.

*Ehrlichia, Anaplasma, and Related Infections (AAP, 2018e)*: PCR testing should be performed within the first week of illness to diagnose anaplasmosis, ehrlichiosis, and other *Anaplasmataceae* infections because doxycycline treatment rapidly decreases the sensitivity of PCR. Consequently, negative PCR results do not necessarily indicate a lack of infection. Occasionally, Giemsa- or Wright staining of blood smears can be performed to identify the presence of the morulae of *Anaplasma* in the first week of illness. Culture testing for isolation is not performed. “Immunoglobulin (Ig) M serologic assays are prone to false-positive reactions, and IgM... can remain elevated for lengthy periods of time, reducing its diagnostic utility. Serologic testing may be used to demonstrate a fourfold change in IgG-specific antibody titer by indirect immunofluorescence antibody (IFA) assay between paired acute and convalescent specimens taken 2 to 4 weeks apart.”

*Rocky Mountain Spotted Fever (RMSF) (AAP, 2018h)*: “The gold standard for serologic diagnosis of RMSF is the indirect immunofluorescence antibody (IFA) test. A negative serologic test result from the acute phase does not rule out a diagnosis of RMSF in any case... A fourfold or greater increase in antigen-specific IgG between acute and convalescent sera obtained 2 to 4 weeks apart confirms the diagnosis (6 weeks for convalescent serum for *Rickettsia africae*).”

*Rickettsialpox (AAP, 2018g)*: Rickettsialpox can be mistaken for other rickettsial infections. Ideally, the use of *R. akari*-specific antigen is recommended for serologic diagnosis, but it has limited availability. Otherwise, indirect IFA for *R. rickettsia*, the causative agent of RMSF, since *R. akari* has extensive cross-reactivity. Again, a demonstration of at least a fourfold increase in antibody titers taken 2-6 weeks apart is indicative of infection.

*Chikungunya (AAP, 2018c)*: “Laboratory diagnosis generally is accompanied by testing serum to detect virus, viral nucleic acid, or virus-specific immunoglobulin (Ig) M and neutralizing antibodies.” RT-PCR can be used to diagnose chikungunya during the first week after onset of symptoms since chikungunya-specific antibodies have not formed at that time. After the first week, serum testing of IgM or a plaque-reduction neutralization test can be performed.

*Dengue (AAP, 2018d)*: “Dengue virus is detectable by RT-PCR or NS1 antigen EIAs from the beginning of the febrile phase until day 7 to 10 after illness onset.” Cross-reactivity occurs between anti-dengue virus IgM and other flaviviruses, including Zika. IgG EIA and hemagglutination testing is not specific for diagnosis of dengue, and IgG antibodies remain elevated for life; consequently, a fourfold increase in IgG between the acute and convalescent phase can confirm recent infection. “Reference testing is available from the Dengue Branch of the Centers for Disease Control and Prevention.”

*Malaria (AAP, 2018f)*: Microscopic identification of *Plasmodium* on both thick and thin blood films should be performed. “If initial blood smears test negative for *Plasmodium* species but malaria remains a possibility, the smear should be repeated every 12 to 24 hours during a 72-hour period... Serologic testing generally is not helpful, except in epidemiologic surveys... Species confirmation and antimalarial drug resistance testing are available free of charge at the Centers for Disease Control and Prevention (CDC) for all cases of malaria diagnosed in the United States.” One FDA-approved RADT is available in the U.S. to hospitals and commercial labs; however, both positive and negative test results must be corroborated by microscopic examination.

*West Nile Virus (WNV) (AAP, 2018i)*: PCR is not recommended for diagnosis of WNV in immunocompetent patients since WNV RNA is usually no longer detectable by the initial onset of symptoms. “Detection of anti-WNV immunoglobulin (Ig) M antibodies in serum or CSF is the most common way to diagnose WNV infection.” Anti-WNV IgM levels can remain elevated for longer than 1 year so a positive test result may be indicative of a prior infection. “Plaque-reduction neutralization tests can be performed to measure virus-specific neutralizing antibodies and to discriminate between cross-reacting antibodies from closely related flaviviruses.

# Testing for Mosquito or Tick-Related Infections AHS – G2158

A fourfold or greater increase in virus-specific neutralizing antibodies between acute-and convalescent-phase serum specimens collected 2 or 3 weeks apart may be used to confirm recent WNV infection.”

## **International Encephalitis Consortium (IEC) (Venkatesan et al., 2013)**

In 2013, the IEC released their *Case Definitions, Diagnostic Algorithms, and Priorities in Encephalitis*. Concerning arboviruses, they state the following: “For most arboviruses, serologic testing of serum and CSF is preferred to molecular testing, since the peak of viremia typically occurs prior to symptom onset. For example, in patients with West Nile virus (WNV) associated with neuroinvasive disease, CSF PCR is relatively insensitive (57%) compared with detection of WNV IgM in CSF. The cumulative percentage of seropositive patients increases by approximately 10% per day during the first week of illness suggesting the need for repeat testing if the suspicion for disease is strong in those with initially negative results. Notably, arbovirus IgM antibodies may be persistently detectable in the serum and, less commonly, in the CSF, for many months after acute infection, and therefore may not be indicative of a current infection. Therefore, if possible, documentation of acute infection by seroconversion and/or 4-fold or greater rises in titre using paired sera is recommended (Venkatesan et al., 2013).”

## **Billing/Coding/Physician Documentation Information**

---

This policy may apply to the following codes. Inclusion of a code in this section does not guarantee that it will be reimbursed. For further information on reimbursement guidelines, please see Administrative Policies on the Blue Cross Blue Shield of North Carolina web site at [www.bcbsnc.com](http://www.bcbsnc.com). They are listed in the Category Search on the Medical Policy search page.

*Applicable Service Codes: 86757, 87040, 87798, 86280, 86666, 86619, 85060, 87207, 87254, 86750, 86753, 86788, 86789, 86790, 87449, 87899, 0043U, 0044U*

BCBSNC may request medical records for determination of medical necessity. When medical records are requested, letters of support and/or explanation are often useful, but are not sufficient documentation unless all specific information needed to make a medical necessity determination is included.

## **Scientific Background and Reference Sources**

---

AAP, A. A. o. P. (2018a). Babesiosis. In D. Kimberlin, M. Brady, M. Jackson, & S. Long (Eds.), Red Book: 2018 Report of the Committee on Infectious Diseases (pp. 235-237): American Academy of Pediatrics.

AAP, A. A. o. P. (2018b). Borrelia Infections Other Than Lyme Disease. In D. Kimberlin, M. Brady, M. Jackson, & S. Long (Eds.), Red Book: 2018 Report of the Committee on Infectious Diseases (pp. 252-255): American Academy of Pediatrics.

AAP, A. A. o. P. (2018c). Chikungunya. In D. Kimberlin, M. Brady, M. Jackson, & S. Long (Eds.), Red Book: 2018 Report of the Committee on Infectious Diseases (pp. 271-272): American Academy of Pediatrics.

AAP, A. A. o. P. (2018d). Dengue. In D. Kimberlin, M. Brady, M. Jackson, & S. Long (Eds.), Red Book: 2018 Report of the Committee on Infectious Diseases (pp. 317-319): American Academy of Pediatrics.

AAP, A. A. o. P. (2018e). Ehrlichia, Anaplasma, and Related Infections. In D. Kimberlin, M. Brady, M. Jackson, & S. Long (Eds.), Red Book: 2018 Report of the Committee on Infectious Diseases (pp. 323-328): American Academy of Pediatrics.

AAP, A. A. o. P. (2018f). Malaria. In D. Kimberlin, M. Brady, M. Jackson, & S. Long (Eds.), Red Book: 2018 Report of the Committee on Infectious Diseases (pp. 527-537): American Academy of Pediatrics.

# Testing for Mosquito or Tick-Related Infections AHS – G2158

AAP, A. A. o. P. (2018g). Rickettsialpox. In D. Kimberlin, M. Brady, M. Jackson, & S. Long (Eds.), *Red Book: 2018 Report of the Committee on Infectious Diseases* (pp. 696-697): American Academy of Pediatrics.

AAP, A. A. o. P. (2018h). Rocky Mountain Spotted Fever. In D. Kimberlin, M. Brady, M. Jackson, & S. Long (Eds.), *Red Book: 2018 Report of the Committee on Infectious Diseases* (pp. 697-700): American Academy of Pediatrics.

AAP, A. A. o. P. (2018i). West Nile Virus. In D. Kimberlin, M. Brady, M. Jackson, & S. Long (Eds.), *Red Book: 2018 Report of the Committee on Infectious Diseases* (pp. 888-891): American Academy of Pediatrics.

Akoolo, L., Schlachter, S., Khan, R., Alter, L., Rojzman, A. D., Gedroic, K., . . . Parveen, N. (2017). A novel quantitative PCR detects Babesia infection in patients not identified by currently available non-nucleic acid amplification tests. *BMC Microbiol*, 17(1), 16. doi:10.1186/s12866-017-0929-2

Arguin, P., & Tan, K. (2017). Chapter 3 Infectious Diseases Related to Travel: Malaria. In G. Brunette (Ed.), *CDC Yellow Book 2018: Health Information for International Travel*. New York: Oxford University Press.

Arguin, P., & Tan, K. (2019). Chapter 3 Infectious Diseases Related to Travel: Malaria. In G. Brunette (Ed.), *CDC Yellow Book 2018: Health Information for International Travel*. New York: Oxford University Press.

Barbour, A. G. (2018, 03/12/2018). Clinical features, diagnosis, and management of relapsing fever. UpToDate. Retrieved from <https://www.uptodate.com/contents/clinical-features-diagnosis-and-management-of-relapsing-fever>

Barbour, A. G. (2020, 03/12/2018). Clinical features, diagnosis, and management of relapsing fever. UpToDate. Retrieved from <https://www.uptodate.com/contents/clinical-features-diagnosis-and-management-of-relapsing-fever>

Biggs, H. M., Behravesh, C. B., Bradley, K. K., Dahlgren, F. S., Drexler, N. A., Dumler, J. S., . . . Traeger, M. S. (2016). Diagnosis and Management of Tickborne Rickettsial Diseases: Rocky Mountain Spotted Fever and Other Spotted Fever Group Rickettsioses, Ehrlichioses, and Anaplasmosis - United States. *MMWR Recomm Rep*, 65(2), 1-44. doi:10.15585/mmwr.rr6502a1

Breman, J. G. (2020, 07/24/2018). Clinical manifestations of malaria in nonpregnant adults and children. UpToDate. Retrieved from <https://www.uptodate.com/contents/clinical-manifestations-of-malaria-in-nonpregnant-adults-and-children>

Burakoff, A., Lehman, J., Fischer, M., Staples, J. E., & Lindsey, N. P. (2018). West Nile Virus and Other Nationally Notifiable Arboviral Diseases - United States, 2016. *MMWR Morb Mortal Wkly Rep*, 67(1), 13-17. doi:10.15585/mmwr.mm6701a3

Calisher, C. H. (1994). Medically important arboviruses of the United States and Canada. *Clin Microbiol Rev*, 7(1), 89-116.

CDC. (2013, 12/14/2015). Dengue in Puerto Rico. Retrieved from <https://www.cdc.gov/dengue/about/inpuerto.html>

CDC. (2014, 09/06/2014). Dengue: Clinical Guidance. Retrieved from <https://www.cdc.gov/dengue/clinicallab/clinical.html>

# Testing for Mosquito or Tick-Related Infections AHS – G2158

CDC. (2015, 01/13/2015). Colorado Tick Fever Health Care Providers: Clinical & Laboratory Evaluation. Retrieved from <https://www.cdc.gov/coloradotickfever/clinicallybeval.html>

CDC. (2018a, 05/17/2018). About the Division of Vector-Borne Diseases. Retrieved from <https://www.cdc.gov/ncezid/dvbd/index.html>

CDC. (2018b, 02/20/2018). Algorithm to Assist in Medical Office Telephone Evaluation of Patients with Possible Influenza. Retrieved from <https://www.cdc.gov/flu/professionals/antivirals/office-evaluation.htm>

CDC. (2018c, 06/06/2018). Babesiosis: Resources for Health Professionals. Retrieved from [https://www.cdc.gov/parasites/babesiosis/health\\_professionals/index.html](https://www.cdc.gov/parasites/babesiosis/health_professionals/index.html)

CDC. (2018d, 12/03/2018). Colorado Tick Fever Health Care Providers: Diagnostic Testing. Retrieved from <https://www.cdc.gov/coloradotickfever/diagnostic-testing.html>

CDC. (2018e, 07/23/2018). Protect Your Baby from Group B Strep. Retrieved from <https://www.cdc.gov/features/groupbstrep/index.html>

CDC. (2018f, 11/26/2018). Tick-borne Relapsing Fever (TBRF): Information for Clinicians. Retrieved from <https://www.cdc.gov/relapsing-fever/clinicians/>

CDC. (2019a, 06/06/2018). Babesiosis: Resources for Health Professionals. Retrieved from [https://www.cdc.gov/parasites/babesiosis/health\\_professionals/index.html](https://www.cdc.gov/parasites/babesiosis/health_professionals/index.html)

CDC. (2019b, 05/03/2019). Dengue: Testing Guidance. Retrieved from <https://www.cdc.gov/dengue/healthcare-providers/testing/testing-guidance.html>

CDC. (2019c). Influenza virus testing in investigational outbreaks in institutional or other closed settings. Retrieved from <https://www.cdc.gov/flu/professionals/diagnosis/guide-virus-diagnostic-tests.htm>

CDC. (2020, 05/03/2019). Dengue: Testing Guidance. Retrieved from <https://www.cdc.gov/dengue/healthcare-providers/testing/testing-guidance.html>

Denison, A. M., Amin, B. D., Nicholson, W. L., & Paddock, C. D. (2014). Detection of *Rickettsia rickettsii*, *Rickettsia parkeri*, and *Rickettsia akari* in skin biopsy specimens using a multiplex real-time polymerase chain reaction assay. *Clin Infect Dis*, 59(5), 635-642. doi:10.1093/cid/ciu358

Dimairo, M. A., Pereira, I. T., George, T. I., & Banaei, N. (2012). Performance of BinaxNOW for diagnosis of malaria in a U.S. hospital. *J Clin Microbiol*, 50(9), 2877-2880. doi:10.1128/jcm.01013-12

FDA. (2018). Devices@FDA. Retrieved from <https://www.accessdata.fda.gov/scripts/cdrh/devicesatfda/index.cfm>

Gershman, M., & Staples, J. (2019). Chapter 3 Infectious Diseases Related to Travel: Yellow Fever. In G. Brunette (Ed.), *CDC Yellow Book 2018: Health Information for International Travel*. New York: Oxford University Press.

Hopkins, H. (2020). Diagnosis of malaria. UpToDate. Retrieved from <https://www.uptodate.com/contents/diagnosis-of-malaria>

Johnson, A. J., Martin, D. A., Karabatsos, N., & Roehrig, J. T. (2000). Detection of anti-arboviral immunoglobulin G by using a monoclonal antibody-based capture enzyme-linked immunosorbent assay. *J Clin Microbiol*, 38(5), 1827-1831.

# Testing for Mosquito or Tick-Related Infections AHS – G2158

Kalish, R. A., McHugh, G., Granquist, J., Shea, B., Ruthazer, R., & Steere, A. C. (2001). Persistence of immunoglobulin M or immunoglobulin G antibody responses to *Borrelia burgdorferi* 10-20 years after active Lyme disease. *Clin Infect Dis*, 33(6), 780-785. doi:10.1086/322669

Kato, C. Y., Chung, I. H., Robinson, L. K., Austin, A. L., Dasch, G. A., & Massung, R. F. (2013). Assessment of real-time PCR assay for detection of *Rickettsia* spp. and *Rickettsia rickettsii* in banked clinical samples. *J Clin Microbiol*, 51(1), 314-317. doi:10.1128/jcm.01723-12

Krause, P. J., & Vannier, E. G. (2019, 11/13/2017). Babesiosis: Clinical manifestations and diagnosis. UpToDate. Retrieved from <https://www.uptodate.com/contents/babesiosis-clinical-manifestations-and-diagnosis>

Meatherall, B., Preston, K., & Pillai, D. R. (2014). False positive malaria rapid diagnostic test in returning traveler with typhoid fever. *BMC Infect Dis*, 14, 377. doi:10.1186/1471-2334-14-377

Miller, J. M., Binnicker, M. J., Campbell, S., Carroll, K. C., Chapin, K. C., Gilligan, P. H., . . . Yao, J. D. (2018). A Guide to Utilization of the Microbiology Laboratory for Diagnosis of Infectious Diseases: 2018 Update by the Infectious Diseases Society of America and the American Society for Microbiology. *Clinical Infectious Diseases*, ciy381-ciy381. doi:10.1093/cid/ciy381

Nasci, R., Fischer, M., Lindsey, N., Lanciotti, R., Savage, H., Komar, N., . . . Petersen, L. (2013). West Nile Virus in the United States: Guidelines for Surveillance, Prevention, and Control. Ft. Collins, CO: CDC Retrieved from <https://www.cdc.gov/westnile/resources/pdfs/wnvguidelines.pdf>

Ota-Sullivan, K., & Blecker-Shelly, D. L. (2013). Use of the rapid BinaxNOW malaria test in a 24-hour laboratory associated with accurate detection and decreased malaria testing turnaround times in a pediatric setting where malaria is not endemic. *J Clin Microbiol*, 51(5), 1567-1569. doi:10.1128/jcm.00293-13

Petersen, L. (2020, 07/30/2018). Clinical manifestations and diagnosis of West Nile virus infection. UpToDate. Retrieved from <https://www.uptodate.com/contents/clinical-manifestations-and-diagnosis-of-west-nile-virus-infection>

Petersen, L. R. (2019). Arthropod-borne encephalitides. UpToDate. Retrieved from <https://www.uptodate.com/contents/arthropod-borne-encephalitides>

Rosenberg, R., Lindsey, N. P., Fischer, M., Gregory, C. J., Hinckley, A. F., Mead, P. S., . . . Petersen, L. R. (2018). Vital Signs: Trends in Reported Vectorborne Disease Cases - United States and Territories, 2004-2016. *MMWR Morb Mortal Wkly Rep*, 67(17), 496-501. doi:10.15585/mmwr.mm6717e1

Ryan, E. T. (2018, 03/30/2018). Yellow Fever. UpToDate. Retrieved from <https://www.uptodate.com/contents/yellow-fever>

Ryan, E. T. (2019, 03/30/2018). Yellow Fever. UpToDate. Retrieved from <https://www.uptodate.com/contents/yellow-fever>

Sexton, D. J., & McClain, M. T. (2016, 08/15/2016). Human ehrlichiosis and anaplasmosis. UpToDate. Retrieved from <https://www.uptodate.com/contents/human-ehrlichiosis-and-anaplasmosis>

Sexton, D. J., & McClain, M. T. (2019a, 08/15/2016). Human ehrlichiosis and anaplasmosis. UpToDate. Retrieved from <https://www.uptodate.com/contents/human-ehrlichiosis-and-anaplasmosis>

Sexton, D. J., & McClain, M. T. (2019b, 09/12/2018). Other spotted fever group rickettsial infections. UpToDate. Retrieved from <https://www.uptodate.com/contents/other-spotted-fever-group-rickettsial-infections>



# Testing for Mosquito or Tick-Related Infections AHS – G2158

Sexton, D. J., & McClain, M. T. (2020, 05/29/2018). Clinical manifestations and diagnosis of Rocky Mountain spotted fever. UpToDate. Retrieved from <https://www.uptodate.com/contents/clinical-manifestations-and-diagnosis-of-rocky-mountain-spotted-fever>

Staples, J., Hills, S., & Powers, A. (2017). Chapter 3 Infectious Diseases Related to Travel: Chikungunya. In G. Brunette (Ed.), *CDC Yellow Book 2018: Health Information for International Travel*. New York: Oxford University Press.

Staples, J., Hills, S., & Powers, A. (2020). Chapter 3 Infectious Diseases Related to Travel: Chikungunya. In G. Brunette (Ed.), *CDC Yellow Book 2018: Health Information for International Travel*. New York: Oxford University Press.

Thomas, S., Rothman, A., Srikiatkachorn, A., & Kalayanarooj, S. (2018, 07/24/2018). Dengue virus infection: Clinical manifestations and diagnosis. UpToDate. Retrieved from <https://www.uptodate.com/contents/dengue-virus-infection-clinical-manifestations-and-diagnosis>

Thomas, S., Rothman, A., Srikiatkachorn, A., & Kalayanarooj, S. (2019, 07/24/2018). Dengue virus infection: Clinical manifestations and diagnosis. UpToDate. Retrieved from <https://www.uptodate.com/contents/dengue-virus-infection-clinical-manifestations-and-diagnosis>

Venkatesan, A., Tunkel, A. R., Bloch, K. C., Laming, A. S., Sejvar, J., Bitnun, A., . . . International Encephalitis, C. (2013). Case definitions, diagnostic algorithms, and priorities in encephalitis: consensus statement of the international encephalitis consortium. *Clin Infect Dis*, 57(8), 1114-1128. doi:10.1093/cid/cit458

WHO. (2017, 10/31/2017). Plague. News-Room. Retrieved from <http://www.who.int/news-room/fact-sheets/detail/plague>

Wilson, M. E., & Lenschow, D. J. (2020). Chikungunya fever. UpToDate. Retrieved from <https://www.uptodate.com/contents/chikungunya-fever>

## Policy Implementation/Update Information

---

- 1/1/19 New policy developed. BCBSNC will provide coverage for testing for mosquito or tick-related infections when it is determined to be medically necessary because the medical criteria and guidelines are met. Medical Director review 1/1/2019. Policy noticed 1/1/2019 for effective date 4/1/2019. (sk)
- 10/29/19 Wording in the Policy, When Covered, and/or Not Covered section(s) changed from Medical Necessity to Reimbursement language, where needed. (gm)
- 12/10/19 Reviewed by Avalon 3rd Quarter CAB. Added PCR testing for suspected cases of tick-borne relapsing fever to When Covered section. Removed NAAT, including PCR for suspected cases of tick-borne relapsing fever from When Not Covered section. Added NAAT, including PCR for suspected cases of babesiosis to When Covered section. Removed NAAT, including PCR for suspected cases of babesiosis from When Not Covered section. (sk)
- 3/10/20 Specialty Matched Consultant Advisory Panel review 2/19/2020. (sk)
- 11/10/20 Reviewed by Avalon 3rd Quarter 2020 CAB. Description section updated. Related policies added. When Covered section updated. When Not Covered section updated. Billing/Coding section updated with addition of the following codes: 0043U, 0044U. Policy Guidelines updated. References updated. (sk)

# Testing for Mosquito or Tick-Related Infections AHS – G2158

3/9/21 Specialty Matched Consultant Advisory Panel review 2/17/2021. (sk)

---

Medical policy is not an authorization, certification, explanation of benefits or a contract. Benefits and eligibility are determined before medical guidelines and payment guidelines are applied. Benefits are determined by the group contract and subscriber certificate that is in effect at the time services are rendered. This document is solely provided for informational purposes only and is based on research of current medical literature and review of common medical practices in the treatment and diagnosis of disease. Medical practices and knowledge are constantly changing and BCBSNC reserves the right to review and revise its medical policies periodically.