

Corporate Medical Policy

Transcranial Magnetic Stimulation as a Treatment of Depression and Other Psychiatric/Neurologic Disorders

File Name: transcranial_magnetic_stimulation_tms
Origination: 10/2019
Last CAP Review: 6/2021
Next CAP Review: 6/2022
Last Review: 12/2021

Description of Procedure or Service

Transcranial magnetic stimulation (TMS), was introduced in 1985 as a method of noninvasive stimulation of the brain. It involves placement of a small coil over the scalp, passing a rapidly alternating current through the coil wire, which produces a magnetic field that passes unimpeded through the scalp and bone, resulting in electrical stimulation of the cortex. TMS was initially used to investigate nerve conduction; eg, TMS over the motor cortex will produce a contralateral muscular-evoked potential. The motor threshold, which is the minimum intensity of stimulation required to induce a motor response, is empirically determined for each person by localizing the site on the scalp for optimal stimulation of a hand muscle, then gradually increasing the intensity of stimulation. The stimulation site for treatment of depression is usually 5 cm anterior to the motor stimulation site.

In contrast to electroconvulsive therapy, TMS does not require general anesthesia and does not generally induce a convulsion. Interest in the use of TMS as a treatment for depression was augmented by the development of a device that could deliver rapid, repetitive stimulation. Imaging studies had shown a decrease in activity of the left dorsolateral prefrontal cortex (DLPFC) in depressed patients, and early studies suggested that high-frequency (eg, 5-10 Hz) TMS of the left DLPFC had antidepressant effects. Low-frequency (1-2 Hz) stimulation of the right DLPFC has also been investigated. The rationale for low-frequency TMS is inhibition of right frontal cortical activity to correct the interhemispheric imbalance. A combination approach (bilateral stimulation), or deep stimulation with a specialized H1 coil attachment may also be used.

Repetitive TMS (rTMS) is also being tested as a treatment for a variety of other disorders including alcohol dependence, Alzheimer disease, neuropathic pain, obsessive-compulsive disorder, postpartum depression, Parkinson disease, stroke, posttraumatic stress disorder, panic disorder, epilepsy, dysphagia, Tourette syndrome, schizophrenia, migraine, spinal cord injury, fibromyalgia, and tinnitus. In addition to the potential for altering interhemispheric imbalance, it has been proposed that high-frequency rTMS may facilitate neuroplasticity.

Regulatory Status

Devices for transcranial stimulation have been cleared for marketing by the U.S. Food and Drug Administration (FDA) for diagnostic uses. A number of devices subsequently received the FDA clearance for the treatment of major depressive disorder in adults who have failed to achieve satisfactory improvement from prior antidepressant medication in the current episode. Indications were expanded to include treating pain associated with certain migraine headaches in 2013, and obsessive-compulsive disorder in 2018. US Food and Drug Administration (FDA) approves TMS device therapy for repeated daily use over 4–6 weeks (20–30 sessions).

Listed below are FDA approved TMS devices for depression and their manufacturers.

Transcranial Magnetic Stimulation as a Treatment of Depression and Other Psychiatric/Neurologic Disorders

Device	Manufacturer	Indication
Neurostar	Neuronetics	Major Depressive Disorder
Brainsway Deep TMS System	Brainsway	Major Depressive Disorder
Rapid Therapy System	Magstim	Major Depressive Disorder
Magvita	Tonica Elektronik	Major Depressive Disorder
Neurosoft	TeleEMG	Major Depressive Disorder
Horizon	Magstim	Major Depressive Disorder
Nexstim	Nexstim	Major Depressive Disorder
Apollo	Mag & More	Major Depressive Disorder
Intermittent Theta Burst Stimulation (iTBS)	Magstim	Major Depressive Disorder
Theta Burst Stimulation (TBS)	Magventure	Major Depressive Disorder

*****Note: This Medical Policy is complex and technical. For questions concerning the technical language and/or specific clinical indications for its use, please consult your physician.**

Policy

BCBSNC will provide coverage for Transcranial Magnetic Stimulation (TMS) when it is determined to be medically necessary because the medical criteria and guidelines shown below are met.

Benefits Application

This medical policy relates only to the services or supplies described herein. Please refer to the Member's Benefit Booklet for a availability of benefits. Member's benefits may vary according to benefit design; therefore member benefit language should be reviewed before applying the terms of this medical policy.

When Transcranial Magnetic Stimulation (TMS) is covered

Transcranial Magnetic Stimulation as a Treatment of Depression and Other Psychiatric/Neurologic Disorders

All types of repetitive transcranial magnetic stimulation (rTMS) of the brain may be considered medically necessary as a treatment of major depressive disorder in adults age 18 years and older when **all** of the following conditions (1-4) have been met:

1. Confirmed diagnosis of severe major depressive disorder (single or recurrent) documented by standardized rating scales that reliably measure depressive symptoms; and
2. Any one of the following (a, b, or c):
 - a. Failure of 4 trials of psychopharmacologic agents including 2 different agent classes and 2 augmentation trials; or
 - b. Inability to tolerate a therapeutic dose of medications as evidenced by 4 trials of psychopharmacologic agents with distinct side effects; or
 - c. Is a candidate for electroconvulsive therapy; further, electroconvulsive therapy would not be clinically superior to rTMS (eg, in cases with psychosis, a acute suicidal risk, catatonia or life-threatening inanition rTMS should NOT be used); and
3. Failure of a trial of a psychotherapy known to be effective in the treatment of major depressive disorder of an adequate frequency and duration, without significant improvement in depressive symptoms, as documented by standardized rating scales that reliably measure depressive symptoms; and
4. Device is FDA approved and utilized in accordance with the FDA labeled indication of major depressive disorder.

Repeat rTMS (retreatment) of the brain may be considered medically necessary as a treatment of major depressive disorder in adults age 18 years and older when all of the following have been met:

- Initial criteria for rTMS treatment were met for previous depressive episode, and
- After initial rTMS treatment course, member achieved remission or at least a 50% reduction in depressive symptoms as measured by a standardized rating scale for depression, and
- Member is experiencing relapse or recurrence of a severe major depressive episode, and the recurrent episode is considered severe, as measured by a standardized rating scale for depression, and
- At least 3 months has lapsed since prior treatment with rTMS, and
- Retreatment is not requested as maintenance therapy or continuous therapy

For initial/repeat therapy, the following may be considered medically necessary:

- 9 week course of treatment
 - 1 unit of 90867
 - 36 units of 90868
 - 5 daily treatments over 6 weeks
 - Taper of twice weekly treatments over 3 weeks
 - Redetermination, 1 unit of 90869

For extension of initial therapy, the following may be considered medically necessary:

- 4 week course of treatment
 - 8 units of 90868
 - Twice weekly treatments over 4 weeks
 - Redetermination, 1 unit of 90869

When Transcranial Magnetic Stimulation (TMS) is not covered

Repetitive TMS for major depressive disorder that does not meet the criteria listed above is considered investigational.

Continued treatment with rTMS of the brain as maintenance therapy is considered investigational

Transcranial Magnetic Stimulation as a Treatment of Depression and Other Psychiatric/Neurologic Disorders

Repetitive TMS of the brain is considered investigational as a treatment of all other psychiatric and neurologic disorders, including but not limited to bipolar disorder, schizophrenia, obsessive-compulsive disorder, or migraine headaches.

Contraindications to repetitive TMS include:

- a. Seizure Disorder or any history of seizure with increased risk of future seizure; or
- b. Presence of acute or chronic psychotic symptoms or disorders (eg, schizophrenia, schizophreniform or schizoaffective disorder) in the current depressive episode; or
- c. Neurologic conditions that include epilepsy, cerebrovascular disease, dementia, increased intracranial pressure, having a history of repetitive or severe head trauma, or with primary or secondary tumors in the central nervous system; or
- d. Presence of an implanted magnetic-sensitive medical device located within 30 centimeters from the TMS magnetic coil or other implanted items including but not limited to a cochlear implant, implanted cardioverter defibrillator, pacemakers, vagus nerve stimulator or metal aneurysm clips or coils, staples, or stents.

Policy Guidelines

Repetitive TMS includes, but is not limited to, the following protocols:

- Theta burst (iTBS)
- Accelerated
- Deep
- Surface or superficial
- Unilateral

Definitions

Acute initial treatment- most studies support that an initial trial of rTMS should last at least three to four weeks before determining whether it is beneficial. However, the number of treatments necessary is not standardized, and the definition of clinical benefit has varied across studies.

The evidence for the efficacy of initial therapy with TMS consists of three randomized, sham controlled trials, with an aggregate 703 patients. Two were industry sponsored (Neurostar, Brainsway) and one was sponsored by the National Institute of Mental Health (NIMH). Only the NIMH-sponsored trial focused on a clinical endpoint of remission, based on the Hamilton Rating Scale for Depression (HAM-D24). In the Neuronetics and Brainsway trials, a meaningful improvement was defined by comparing the baseline to endpoint change on the Montgomery-Asberg Depression Rating Scale (MADRS) and/or HAM scales between the active and sham groups.

Extension of initial treatment for additional weeks is reasonable in patients who only achieve a partial response to initial therapy that has not plateaued and for those who have a prior history of late response to antidepressant medications (late response is defined as after 10 weeks).

The evidence for extension of initial treatment is from two observational studies (Avery et al 2008, McDonald et al 2011). In the McDonald study, patients who were part of a randomized, sham controlled trial of TMS were allowed to receive extension therapy if after three weeks of initial therapy they did not achieve remission (remission defined as a HAM-D score of ≤ 3 or two consecutive weekly HAM-D scores less than 10), but who improved (improvement defined as having at least a 30% reduction in HAM-D score from baseline but without an absolute score meeting remission criteria). This group continued extended treatment for up to 3 additional weeks if they continued to show improvement (defined as at least a two point drop in the HAM-D per week). Remission occurred in 31 percent. In the Avery study, 73 patients who did not respond to daily rTMS for four to six weeks were treated for six more weeks, and response (defined as improvement from baseline on a depression rating scale $\geq 50\%$) occurred in 26 percent.

Transcranial Magnetic Stimulation as a Treatment of Depression and Other Psychiatric/Neurologic Disorders

Maintenance therapy: Maintenance TMS may be defined as a course that begins after the end of the initial therapy and is intended to prevent recurrence of a new episode. However, the frequency of sessions for maintenance TMS has not been standardized, and evidence is not available to support one specific maintenance schedule over another. Nor is it known for how long a patient should receive maintenance.

There is a lack of randomized trials to assess and define maintenance TMS therapy. The best evidence for maintenance consists of one small prospective trial (Philip, 2015) which enrolled 67 patients with treatment resistant depression who underwent 6 weeks of acute TMS and were randomized after 6 weeks to scheduled maintenance or observation if they responded to the six weeks of acute therapy. Forty-nine patients who met the response criteria (acute phase endpoint HAM-D17 total score <15 and more than 25% improvement in total score HAM-D17 compared to baseline) were randomized to a single TMS session scheduled monthly as maintenance or observation only, with 12 months follow-up. There was a longer time to relapse in the group that received the scheduled maintenance, but it did not reach statistical significance.

The remaining published evidence for maintenance therapy consists of small observational studies.

Depression Assessment Instruments

The Hamilton Rating Scale for Depression (HAM-D24) is a 17-item rating scale to determine the severity level of depression in a patient before, during, and after treatment. The total score ranges from 0 to 52, with the score corresponding to the following classifications:

- 0-7: No depression (normal)
- 8-16: Mild depression
- 17-23: Moderate depression
- ≥24: Severe depression

Montgomery-Asberg Depression Rating Scale (MADRS), an adaptation of the HAM-D, is a 10-item scale with each item rated on a 0-6-point scale. MADRS investigates the presence of affective, somatic, cognitive and behavioral symptoms of depression. MADRS has a greater sensitivity to change in symptoms over time. The total score classifies the patient's level of severity of depression as normal or absent 0-6; mild 7-19; moderate 20-34; severe 35-60.

Patient Health Questionnaire 9 (PHQ-9) relies on patient self-report. There are nine questions based on the nine DSM-IV depression diagnostic criteria, and scores are based on frequency of symptoms. The possible range is 0-27, with minimal depression 0-4, mild 5-9, moderate 10-14, moderately severe 15-19, and severe 20-27.

Evidence summaries

For individuals who have TRD who receive rTMS, the evidence includes a large number of sham-controlled randomized trials and meta-analyses of these trials. Relevant outcomes are symptoms, functional outcomes, and quality of life. The meta-analyses found a clinical benefit associated with rTMS for TRD with improved response rates and rates of remission compared with sham. The most recent meta-analyses have concluded that the effect of rTMS, on average depression scores, is smaller than the effect of ECT on TRD and that the mean improvement in depression scores with rTMS did not reach the minimal clinically important difference; however, clinically meaningful improvements were noted in a subgroup of studies using higher frequency pulses. One potential area of benefit for rTMS is in accelerating or enhancing the response to antidepressant medications, and there is some evidence that rTMS, when given in conjunction with the initiation of pharmacologic therapy, improves the response rate compared with pharmacologic therapy alone. The effect of rTMS appears to be less robust when it is given in combination with a stable dose of antidepressant medication.

Transcranial Magnetic Stimulation as a Treatment of Depression and Other Psychiatric/Neurologic Disorders

Meta-analyses have also found that the efficacy of rTMS decreases with longer follow-up, though some studies have reported persistent response up to 6 months in some patients. There is limited evidence to compare the effects of these treatments on cognition, although the adverse events of rTMS appear to be minimal. While the most recent meta-analyses have reported that the effect of rTMS is smaller than the effect of ECT on TRD, because rTMS does not require general anesthesia or induce seizures, some individuals may decline ECT so the balance of incremental benefits and harms associated with rTMS may be a reasonable compared with ECT. Based on the short-term benefit observed in RCTs and the lack of alternative treatments, aside from ECT in patients with TRD, rTMS may be considered a treatment option in patients with TRD who meet specific criteria. The evidence for theta burst stimulation includes a large randomized trial showing noninferiority with another method of rTMS; no significant differences were noted in the number of adverse events. The evidence is sufficient to determine that the technology results in a meaningful improvement in the net health outcome.

For individuals who have psychiatric or neurologic disorders other than depression (eg, amyotrophic lateral sclerosis, chronic pain, epilepsy, fibromyalgia, migraine headache, obsessive-compulsive disorder, panic disorder, Parkinson disease, posttraumatic stress disorder, schizophrenia, stroke, substance abuse and craving) who receive rTMS, the evidence includes numerous small RCTs and meta-analyses of these randomized trials. Relevant outcomes are symptoms, functional outcomes, and quality of life. The trials included in the meta-analyses are typically small and of low methodologic quality. In addition, stimulation parameters have not been established, and trial results are heterogeneous. There are no large, high-quality trials for any of these conditions demonstrating efficacy or the durability of any treatment effects. The evidence is insufficient to determine the effects of the technology on health outcomes.

The American Psychiatric Association (2018) published consensus recommendations on repetitive transcranial magnetic stimulation (rTMS) for the treatment of depression. The guidelines state, "Multiple randomized controlled trials and published literature have supported the safety and efficacy of rTMS antidepressant therapy." The recommendations include information on the following variables: clinical environment, operator requirements, documentation, coils, cortical targets, coil positioning methods, determination of motor threshold, number of treatment sessions for a acute treatment, and allowable psychotropic medications during TMS treatment. The recommendations do not address use of one type of rTMS over another.

The Clinical TMS Society (2017) published consensus recommendations on TMS therapy for major depressive disorder. These recommendations concluded that clinical benefit was found within 6 weeks of daily treatment sessions. Continued treatment beyond 6 weeks in specific circumstances was also noted to have a clinical benefit.

The National Institute for Health and Care Excellence (2015) provided revised guidance, stating that evidence on the short-term efficacy of rTMS for depression is adequate, although the clinical response is variable and some patients may not benefit.

Billing/Coding/Physician Documentation Information

This policy may apply to the following codes. Inclusion of a code in this section does not guarantee that it will be reimbursed. For further information on reimbursement guidelines, please see Administrative Policies on the Blue Cross Blue Shield of North Carolina web site at www.bcbsnc.com. They are listed in the Category Search on the Medical Policy search page.

Applicable service codes: 90867, 90868, 90869

Transcranial Magnetic Stimulation as a Treatment of Depression and Other Psychiatric/Neurologic Disorders

BCBSNC may request medical records for determination of medical necessity. When medical records are requested, letters of support and/or explanation are often useful, but are not sufficient documentation unless all specific information needed to make a medical necessity determination is included.

Scientific Background and Reference Sources

Blue Cross and Blue Shield Association Technology Evaluation Center (TEC). Transcranial magnetic stimulation for depression. TEC Assessments. 2009; Volume 24: Tab 5.

Blue Cross and Blue Shield Association Technology Evaluation Center (TEC). Transcranial magnetic stimulation for depression. TEC Assessments. 2011; Volume 26: Tab 3.

Blue Cross and Blue Shield Association Technology Evaluation Center (TEC). Transcranial magnetic stimulation for depression. TEC Assessments. 2013; Volume 28: Tab 9.

BCBSA Medical Policy Reference Manual [Electronic Version]. 2.01.50, 10/10/2018

National Institute for Health and Care Excellence (NICE). Repetitive transcranial magnetic stimulation for depression [IPG542]. 2015; <https://www.nice.org.uk/guidance/ipg542>. Accessed February 17, 2020.

American Psychiatric Association. Practice guideline for the treatment of patients with obsessive-compulsive disorder. 2007; http://psychiatryonline.org/pb/assets/raw/sitewide/practice_guidelines/guidelines/ocd.pdf. Accessed February 17 2020.

Blumberger DM, Vila-Rodriguez F, Thorpe KE, et al. Effectiveness of theta burst versus high-frequency repetitive transcranial magnetic stimulation in patients with depression (THREE-D): a randomised non-inferiority trial. *Lancet*. Apr 28 2018;391(10131):1683-1692. PMID 29726344

Brunoni AR, Chaimani A, Moffa AH, et al. Repetitive Transcranial Magnetic Stimulation for the Acute Treatment of Major Depressive Episodes: A Systematic Review With Network Meta-analysis. *JAMA Psychiatry*. 2017;74(2):143–152. doi:10.1001/jamapsychiatry.2016.3644

Perera T, George MS, Grammer G, Janicak PG, Pascual-Leone A, Wirecki TS. The Clinical TMS Society Consensus Review and Treatment Recommendations for TMS Therapy for Major Depressive Disorder. *Brain Stimul*. 2016;9(3):336–346. doi:10.1016/j.brs.2016.03.010

Senova S, Cotovio G, Pascual-Leone A, Oliveira-Maia AJ. Durability of antidepressant response to repetitive transcranial magnetic stimulation: Systematic review and meta-analysis. *Brain Stimul*. 2019;12(1):119-128. doi:10.1016/j.brs.2018.10.001

Avery DH, Isenberg KE, Sampson SM, et al. Transcranial magnetic stimulation in the acute treatment of major depressive disorder: clinical response in an open-label extension trial. *J Clin Psychiatry*. 2008;69(3):441-451.

George MS, Lisanby SH, Avery D, et al. Daily left prefrontal transcranial magnetic stimulation therapy for major depressive disorder: a sham-controlled randomized trial. *Arch Gen Psychiatry*. 2010;67(5):507-516.

Levkovitz Y, Isserles M, Padberg F, et al. Efficacy and safety of deep transcranial magnetic stimulation for major depression: a prospective multicenter randomized controlled trial. *World Psychiatry*. 2015;14(1):64-73.

Transcranial Magnetic Stimulation as a Treatment of Depression and Other Psychiatric/Neurologic Disorders

McDonald WM, Durkalski V, Ball ER, et al. Improving the antidepressant efficacy of transcranial magnetic stimulation: maximizing the number of stimulations and treatment location in treatment-resistant depression. *Depress Anxiety*. 2011;28(11):973-980.

O'Reardon JP, Solvason HB, Janicak PG, et al. Efficacy and safety of transcranial magnetic stimulation in the acute treatment of major depression: a multisite randomized controlled trial. *Biol Psychiatry*. 2007;62(11):1208-1216.

Philip NS, Dunner DL, Dowd SM, et al. Can Medication Free, Treatment-Resistant, Depressed Patients Who Initially Respond to TMS Be Maintained Off Medications? A Prospective, 12-Month Multisite Randomized Pilot Study. *Brain Stimul*. 2016;9(2):251-257.

BCBSA Medical Policy Reference Manual [Electronic Version]. 2.01.50, 10/17/2019

Specialty Matched Consultant Advisory Panel – 6/2020

Carmi, L., Tendler, A., Bystritsky, A., Hollander, E., Blumberger, D. M., Daskalakis, J., Ward, H., Lapidus, K., Goodman, W., Casuto, L., Feifel, D., Barnea-Ygael, N., Roth, Y., Zangen, A., & Zohar, J. (2019). Efficacy and Safety of Deep Transcranial Magnetic Stimulation for Obsessive-Compulsive Disorder: A Prospective Multicenter Randomized Double-Blind Placebo-Controlled Trial. *The American journal of psychiatry*, 176(11), 931–938. <https://doi.org/10.1176/appi.ajp.2019.18101180>

Centers for Medicare and Medicaid Services (CMS). Local Coverage Determination (LCD): Repetitive Transcranial Magnetic Stimulation (rTMS) in Adults with Treatment Resistant Major Depressive Disorder (L34998). Accessed December 20, 2020. <https://www.cms.gov/medicare-coverage-database/details/lcd-details.aspx?LCDId=34998&ver=28&NCDId=230&ncdver=2&SearchType=Advanced&CoverageSelection=Both&NCSelection=NCD&PolicyType=Final&s=All&Keyword=vagus&KeywordLookUp=Doc&KeywordSearchType=Exact&kq=true&bc=IAAAACgAAAA&>

Dunner, D. L., Aaronson, S. T., Sackeim, H. A., Janicak, P. G., Carpenter, L. L., Boyadjis, T., Brock, D. G., Bonneh-Barkay, D., Cook, I. A., Lanocha, K., Solvason, H. B., & Demitrack, M. A. (2014). A multisite, naturalistic, observational study of transcranial magnetic stimulation for patients with pharmacoresistant major depressive disorder: durability of benefit over a 1-year follow-up period. *The Journal of clinical psychiatry*, 75(12), 1394–1401. <https://doi.org/10.4088/JCP.13m08977>

Fukuda, A. M., Tirrell, E., Gobin, A. P., & Carpenter, L. L. (2019). Repetitive Transcranial Magnetic Stimulation for depression relapse or recurrence: Naturalistic retreatment series outcomes. *Brain stimulation*, 12(5), 1328–1329. doi.org/10.1016/j.brs.2019.06.016

Trevizol AP, Shiozawa P, Cook IA, Sato IA, Kaku CB, Guimarães FB, Sachdev P, Sarkhel S, Cordeiro Q. Transcranial Magnetic Stimulation for Obsessive-Compulsive Disorder: An Updated Systematic Review and Meta-analysis. *J ECT*. 2016 Dec;32(4):262-266. doi: 10.1097/YCT.0000000000000335. PMID: 27327557.

BCBSA Medical Policy Reference Manual [Electronic Version]. 3.01.03, 10/15/2020

Specialty Matched Consultant Advisory Panel – 6/2021

Policy Implementation/Update Information

Transcranial Magnetic Stimulation as a Treatment of Depression and Other Psychiatric/Neurologic Disorders

- 10/15/19 New policy developed. TMS considered medically necessary as a treatment of major depressive disorder when criteria are met. Medical Director review 10/2019. (hb)
- 3/10/20 Description section updated. Clarified When covered statement to include all types of repetitive TMS. When covered section #4 added to require protocol and device be FDA approved. Note listing forms of rTMS added to Policy Guidelines for clarity. References added. (eel)
- 3/24/20 Regulatory section updated with FDA unit limits. Policy Guidelines updated with support of unit limits. Note added limiting units in When covered section. References added. **Policy noticed 3/24/20 for effective date 5/26/20.** (eel)
- 9/08/20 Clarification added to define the terms extension of initial therapy and maintenance to Policy Guidelines section. References added. Specialty Matched Consultant Advisory Panel review 6/17/2020. Medical Director review. No change to policy statement. (bb)
- 1/26/21 When covered section updated to clarify a age requirement as "18 years or older" based on current FDA approval and criteria added for repeat rTMS. Medical Director review. References added. (bb)
- 7/13/21 References added. Specialty Matched Consultant Advisory Panel review 6/2021. Medical Director review. No change to policy statement. (bb)

Medical policy is not an authorization, certification, explanation of benefits or a contract. Benefits and eligibility are determined before medical guidelines and payment guidelines are applied. Benefits are determined by the group contract and subscriber certificate that is in effect at the time services are rendered. This document is solely provided for informational purposes only and is based on research of current medical literature and review of common medical practices in the treatment and diagnosis of disease. Medical practices and knowledge are constantly changing and BCBSNC reserves the right to review and revise its medical policies periodically.