

Corporate Medical Policy

Venous and Arterial Thrombosis Risk Testing AHS – M2041

File Name: venous_and_arterial_thrombosis_risk_testing
Origination: 01/2019
Last Review: 04/2022

Description of Procedure or Service

Definitions

A thrombosis, also known as a blood clot, occurs within blood vessels in the body. The two main types of thrombosis include venous thrombosis, which is when a vein is blocked due to a blood clot, and arterial thrombosis, which is when an artery is blocked due to a blood clot. Thrombophilias refer to hereditary and/or acquired abnormalities of hemostasis that predispose patients to thrombosis (Stevens et al., 2016). The most common presentations of venous thromboembolism (VTE) are deep vein thrombosis (DVT) and pulmonary embolism (PE) (Bartholomew, 2017).

Related Policies

Cardiovascular Disease Risk Assessment AHS –G2050

Genetic Testing for Lipoprotein A Variant(s) as a Decision Aid for Aspirin Treatment and/or CVD Risk Assessment AHS- M2082

******Note: This Medical Policy is complex and technical. For questions concerning the technical language and/or specific clinical indications for its use, please consult your physician.***

Policy

BCBSNC will provide coverage for venous and arterial thrombosis risk testing when it is determined the medical criteria or reimbursement guidelines below are met.

Benefits Application

This medical policy relates only to the services or supplies described herein. Please refer to the Member's Benefit Booklet for availability of benefits. Member's benefits may vary according to benefit design; therefore, member benefit language should be reviewed before applying the terms of this medical policy.

When Venous and Arterial Thrombosis Risk Testing is covered

1. Genetic testing for Factor V Leiden and Prothrombin gene G20210A mutations is considered medically necessary in patients without recurrent VTE risk factors (for example, surgery, prolonged immobilization, collagen vascular disease, malignancy, certain hematologic disorders) in any of the following situations:
 - a. Age <50, any deep venous thrombosis
 - b. Deep venous thrombosis in unusual sites (such as hepatic, mesenteric, and cerebral veins)
 - c. Deep venous thrombosis and a strong family history of thrombotic disease
 - d. Deep venous thrombosis in pregnant women or in women taking oral contraceptives

Venous and Arterial Thrombosis Risk Testing AHS – M2041

- e. First- and second-degree relatives of individuals with deep venous thrombosis under age 50
 - f. Myocardial infarction in female smokers under age 50
 - g. Before administration of oral contraceptives, targeted testing of women with a personal or family history of deep venous thrombosis
 - h. Pediatric arterial ischemic stroke
2. Reimbursement for plasma testing for protein C deficiency, protein S deficiency and antithrombin III deficiency is allowed in patients without recurrent VTE risk factors (for example, surgery, prolonged immobilization, collagen vascular disease, malignancy, certain hematologic disorders) in any of the following situations. Testing should be performed at least six weeks after acute thrombotic event and while the patient is not taking anticoagulants. Assays for clotting inhibitors amount and function should be performed prior to any molecular testing.
- a. Age <50, any deep venous thrombosis
 - b. Deep venous thrombosis in unusual sites (such as hepatic, mesenteric, and cerebral veins)
 - c. Deep venous thrombosis and a strong family history of thrombotic disease
 - d. Deep venous thrombosis in pregnant women or in women taking oral contraceptives
 - e. Relatives of individuals with deep venous thrombosis under age 50
 - f. Myocardial infarction in female smokers under age 50
 - g. Before administration of oral contraceptives, targeted testing of women with a personal or family history of deep venous thrombosis
 - h. Individuals with warfarin induced skin necrosis
 - i. Infants who develop Neonatal Purpura Fulminans
 - j. Pediatric arterial ischemic stroke

When Venous and Arterial Thrombosis Risk Testing is not covered

1. Reimbursement is not allowed for MTHFR genetic testing for hypercoagulable evaluation or for “at risk” family members.
2. Reimbursement is not allowed for genetic testing for Factor V Leiden and Prothrombin gene G20210A mutations in patients with recurrent thrombotic events who are receiving a lifelong anticoagulation regimen.
2. Reimbursement is not allowed for venous thrombosis risk testing for superficial venous thrombosis (including superficial thrombophlebitis and varicosities).
3. Genetic Testing for inherited thrombophilia is considered not medically necessary for the following situations:
 - a. Evaluation of recurrent fetal loss, placental abruption, preeclampsia, or fetal growth restriction
 - b. Evaluation of arterial thrombosis not attributable to paradoxical emboli
 - c. Routine screening in the general population
 - d. Routine screening of asymptomatic women considering oral contraceptive use or hormone replacement therapy
 - e. Routine screening of asymptomatic pregnant women
 - f. Prenatal or preimplantation testing
 - g. Routine newborn screening
 - h. Testing more than once per lifetime

Venous and Arterial Thrombosis Risk Testing AHS – M2041

3. Testing for other factors, including the factor V HR2 variant, or prothrombin G1199A variant, or factor VII R353Q variant, or factor 13B V34L variant or PAI-1, as well as multi-gene panel testing is considered not medically necessary.
4. Deep venous thrombosis risk testing as part of a pre-transplant evaluation test is considered not medically necessary.
5. Genetic testing for Factor V Leiden mutation and Prothrombin gene G20210A mutation to determine arterial thrombosis risk is considered not medically necessary.

Policy Guidelines

Table of Terminology

Term	Definition
ACC	American College of Cardiology
ACMG	American College of Medical Genetics and Genomics
ACOG	American College of Obstetricians and Gynecologists
AHA/ASA	American Heart Association/American Stroke Association
APC	Activated protein c
APS	Antiphospholipid syndrome
ASCP	American Society for Clinical Pathology
ASH	American Society of Hematology
CTPA	Computed tomography pulmonary angiography
DVT	Deep vein thrombosis
EGAPP	Evaluation of genomic applications in practice and prevention
ESC	European Society of Cardiology
FVL	Factor V Leiden
HRT	Hormone replacement therapy
NAFT	The North American Thrombosis Forum
OC	Oral contraceptives
PE	Pulmonary embolism
SIGN	Scottish Intercollegiate Guidelines Network
SPESI	Simplified pulmonary embolism severity index
STEMI	Subsequent ST elevation myocardial infarction
SVM	Society for Vascular Medicine
VQ	Ventilation perfusion
VTE	Venous thromboembolism
VUS	Variant of unknown significance

Background

A thrombus is “an aggregate of coagulated blood within the vascular system or heart which contains platelets, fibrin, leukocytes, and red blood cells in varying amounts” (Herrmann, 2018). This aggregate of blood can be problematic as it may obstruct normal blood circulation throughout the

Venous and Arterial Thrombosis Risk Testing AHS – M2041

body and even travel to peripheral areas. The primary manifestations of venous thromboembolisms (VTE) are deep vein thrombosis and pulmonary embolism. These conditions affect an estimated one million individuals in the United States annually (Bartholomew, 2017).

Thrombosis is widely theorized to develop due to Virchow's Triad, which consists of abnormalities in blood flow, a vascular endothelial injury, and alterations in the blood constituents. Changes in any of these characteristics may cause the clot to form (Bauer & Lip, 2018, 2020). For example, sickle red blood cells may cause increased clumping or decreased adhesion to the vessel walls (Byrnes & Wolberg, 2017). There are two main types of thrombosis: venous thrombosis (when a vein is blocked due to a blood clot) and arterial thrombosis (when an artery is blocked due to a blood clot).

A deep vein thrombosis (DVT) refers to a thrombus in a "deep" vein whereas a pulmonary embolism (PE) refers to an obstruction of the pulmonary artery (or one of its branches) by foreign material (Bauer, 2020; Thompson, 2018, 2020). DVT of the lower extremities may cause symptoms, such as swelling or edema in the lower extremities, pain, and warmth in the affected area (Bauer, 2020). This thrombus may travel to the lungs (becoming an embolus) and cause a PE. A PE has similar symptoms to DVT but may include pulmonary issues, such as shortness of breath. The risk factors for VTE, PE, and DVT are similar (Thompson, 2020). The two primary categories of risk factors for VTE are hereditary and acquired, and the genetic tendency toward VTE is referred to as inherited thrombophilia. Hereditary risk factors include genetic mutations such as Factor V Leiden (FVL) mutations. The five most common genetic risk factors for VTE are FVL mutations, prothrombin mutations, protein S defect, protein C defect, and antithrombin defect (Bauer & Lip, 2020). Approximately 50–60% of the variance in VTE incidence are attributed to genetic effects (Crous-Bou, Harrington, & Kabrhel, 2016).

FVL mutations cause coagulation factor V to be unresponsive to activated protein C. A single nucleotide change (G1691A) results in a point mutation of glutamine to arginine at position 506. Approximately 99% of carriers of this mutation are heterozygous, and only 5% of these heterozygotes will experience a VTE in their lifetime. These mutations are often suspected in patients experiencing a VTE at a young age (under 50), a VTE in unusual areas such as a portal vein, or recurrent VTEs (Bauer, 2019a). Protein C may also be genetically deficient, but this mutation is only seen in 2-5% of individuals with a VTE (Bauer, 2020). Protein S, a cofactor for the activated protein C control mechanism, and deficiencies in this protein may also confer additional risk for VTE (Bauer, 2021).

The second most common inherited thrombophilia is the G20210A mutation of prothrombin. This mutation is a gain of function mutation where clotting activity is increased by creating more thrombin and fibrin. The overall prevalence of this mutation is about 2% (Bauer, 2019c). Genetic defects of antithrombin (an inhibitor of thrombin) may also occur, but the estimated prevalence of antithrombin defects is only a maximum of 0.2% (Bauer, 2019b).

Acquired risk factors or predisposing conditions for thrombosis include a prior thrombotic event, recent major surgery, presence of a central venous catheter, trauma, immobilization, malignancy, pregnancy, the use of oral contraceptives or heparin, myeloproliferative disorders, antiphospholipid syndrome (APS), and a number of other major medical illnesses (Bauer & Lip, 2020). Patients with acquired hypercoagulability have an increased risk of venous thrombosis, arterial thrombosis, or both; however, there is a low risk of recurrence, regardless of thrombophilia status (Connors, 2017).

Risk factors for arterial thrombosis are lesser known. The relationship between FVL and arterial thrombosis is controversial with studies reporting varying results; overall, FVL is not currently considered a major risk factor for arterial thrombosis (Carroll & Piazza, 2018; Kujovich, 2011). Kujovich (2018) states that FVL testing should not be performed on persons with any type of arterial thrombosis including myocardial infarction and stroke in children or adults. It has also been reported that while inherited antithrombin, protein S and protein C deficiencies are important risk factors for venous thrombosis, "they have little or no effect on arterial thrombosis" (Previtali, Bucciarelli, Passamonti, & Martinelli, 2011). Further, prothrombin gene mutation is not

Venous and Arterial Thrombosis Risk Testing AHS – M2041

consistently shown to increase the risk of an arterial thromboembolism, and “There is no association of antithrombin deficiency with arterial thrombosis” (Carroll & Piazza, 2018). It has been proposed that venous thrombosis risk testing may be beneficial as a pre-transplant evaluation test. However, no studies have been identified suggesting this. The North American Thrombosis Forum (NAFT) states that even though certain genetic conditions predispose a small proportion of the population to the development of blood clots, “few people with thrombophilias develop symptoms”; further, there is no cost-effective, safe or long-term method to prevent a blood clot from forming even if a genetic predisposition is identified (NATF, 2019).

Thrombotic events such as thrombophilia and stroke have become increasingly documented in hospitalized pediatric patients with underlying medical conditions such as prematurity, cancer, and congenital heart disease, but they are rarely identified in healthy children. Furthermore, in most cases of pediatric venous thromboembolism, there exist other underlying risk factors such as indwelling central venous catheter and inherited thrombophilia that are worthy of further investigation. The incidence of venous thromboembolisms is highest in neonates and infants, but there is a second peak recorded in adolescence, coinciding with the use of oral contraceptives. However, as in the case with adults, little to no evidence suggests that the use of venous thrombosis risk testing in children will affect the acute management of venous thromboembolisms. In a study including a total of 271 children with VTE, it was found that the relative frequencies of individually inherited thrombophilias were low—for example, the highest recorded frequency of IT disorders was of Factor V Leiden, occurring in only 5 to 10 percent of the samples. Moreover, a study of 52 children with thromboembolic events during the acute phase did not urge any changes to acute management, regardless of the result of the test (Raffini, Mahoney, & Armsby, 2019).

Venous thrombosis risk testing has also been entertained as a manner of combatting pediatric stroke, which can be characterized in a variety of ways, such as by age and by presentation. Arterial ischemic infarctions are the most common, comprising approximately 80% of all perinatal strokes, and this form of stroke can occur in up to 1 in 3500 of newborns. However, though it would seem reasonable for venous thrombosis risk testing to be employed here, recent prospective case-control studies suggest that routine thrombophilia testing is not warranted. The study showed that conditions associated with thrombophilia rarely coincided with arterial ischemic strokes, and these conditions included, but were not limited to, decreased levels of protein C, protein S, or prothrombin, and genotyping of factor V Leiden (FVL) and factor II (FII, prothrombin) G20210A. Of the 14 parameters examined, 12 showed no difference, including all common thrombophilias examined, with specific mention that FVL and FII were comparable to population norms (Curtis et al., 2017; Ferriero et al., 2019). Subsequent evaluation deemed thrombophilia evaluation in neonates as having limited clinical utility because “levels of protein C, protein S, antithrombin, and factor XI are normally decreased to 30% of adults levels, and these levels only approach adult levels at various time points during childhood”. Therefore, the use of thrombophilia testing for these proteins may be misleading in the neonatal period, and MRIs instead should be used to diagnose the thrombosis (Ferriero et al., 2019). Moreover, studies focusing on the roles of thrombophilia, arteriopathy, and cardiac abnormalities in perinatal ischemic stroke find that these risk factors were at best unclear, weakening what predictive power they were believed to contain for even recurrent events after perinatal stroke and leading researchers to conclude that thrombophilia evaluation should rarely be considered in cases of perinatal stroke (Lehman et al., 2017).

Clinical Utility and Validity

A D-dimer assay is a blood test that is used in clinical practice to assist in identifying if a patient has a DVT or PE; this test may also help patients experiencing unprovoked VTE to determine if anticoagulation treatment should continue or halt after initial treatment is complete (Linkins & Takach Lapner, 2017). A D-dimer assay may vary greatly based on the type of antibody used, the method of capture, calibration, and instrumentation. Currently, 30 different assays are available which use 20 different monoclonal antibodies; various studies have reported a broad sensitivity and specificity range for D-dimer assays from 69-97% and 43-99% respectively (Linkins & Takach

Venous and Arterial Thrombosis Risk Testing AHS – M2041

Lapner, 2017). Hence, all D-dimer assays differ and need to be validated within the population of interest. Because of this, comparing study results is challenging.

Factor VIII is a blood clotting protein encoded by the F8 gene. A case report by Algahtani and Stuckey (2019) suggests that high factor VIII levels may also assist in risk factor determination for thrombosis or ischemic heart disease. “We conclude that high factor VIII levels are a risk factor for thrombosis, with a greater impact on venous than on arterial thrombosis. However, due to a lack of international consensus on methods for the laboratory testing of factor VIII levels in plasma, we would not currently recommend the measurement of factor VIII levels as part of routine thrombophilia screening (Algahtani & Stuckey, 2019).” This relationship has been shown previously as elevated levels of coagulation factor VIII:C were identified in a retrospective study of 584 first-degree relatives of 177 patients with high coagulation factor VIII:C levels; the researchers found that 40% of first degree relatives also had high VIII:C levels and were at an increased risk for VTE and arterial thrombosis when compared to other first-degree relatives with normal VIII:C levels (Bank et al., 2005).

Lee et al. (2017) performed whole exome sequencing on 64 patients with VTE to assess the types of mutations of inherited thrombophilias. Of these 64 patients, 39 of them were found to have a pathogenic variant or variant of unknown significance (VUS). Further, eight were found to have a Factor V mutation (6 with FVL and 2 with less common mutations), two were found to have a prothrombin G20210A mutation, six were found to have a protein S mutation, two were found to have a protein C mutation, and three were found to have an antithrombin mutation (Lee et al., 2017).

Segal et al. (2009) reviewed the utility of FVL and prothrombin G20210A testing. The authors reviewed 124 articles and concluded that although genetic testing for these two risk factors is very accurate (valid), the clinical utility is lacking due to lack of evidence demonstrating improvement in clinical outcomes (Segal et al., 2009).

Onda studied the clinical utility of a new diagnostic algorithm based on serum D-dimer levels for VTE after hepatectomy. 742 patients who underwent hepatectomy were enrolled in the study and measured for serum D-dimer levels post-op. CT scan was performed for patients who had a D-dimer level of greater than 20 µg/mL. Based on D-dimer and CT scan, VTE was diagnosed in 26 patients and pulmonary embolism (PE) was diagnosed in 18 patients. Multivariate analysis also showed that a resected liver weight of more than 120 grams is a significant predictor of VTE. Overall, “patients who undergo hepatectomy are at high risk for VTE, especially when the resected liver weight is high. The proposed diagnostic algorithm based on serum D-dimer levels for VTE after hepatectomy can be useful for early diagnosis” (Onda et al., 2021).

Analytical Validity

Murphy and Sabath (2019) have compared the accuracy and reliability of two tests: a genotypic assay which identifies FVL mutations, and a phenotypic activated protein C resistance assay. Data from 1596 patients was analyzed; each patient had received both types of testing. The authors state that the phenotypic testing exhibited both high sensitivity and specificity compared to genotypic testing. “Phenotypic assays had close to total concordance with genotypic assays over 16 years of testing. Changing ordering practices could result in up to an 80% reduction in testing costs (Murphy & Sabath, 2019).”

A systematic review and meta-analysis by Chiasakul et al. (2019) researched the relationship between inherited thrombophilia and the risk of arterial ischemic stroke in adults. Inherited thrombophilias included FVL, protein C and S deficiency, antithrombin deficiency and prothrombin G20210A mutation. For this study, 11,916 stroke patients and 96,057 controls were identified. The authors concluded that “Compared with controls, patients with arterial ischemic stroke were significantly more likely to have the following inherited thrombophilias: factor V Leiden (OR, 1.25; 95% CI, 1.08-1.44; I²=0%), prothrombin G20210A mutation (OR, 1.48; 95% CI, 1.22-1.80;

Venous and Arterial Thrombosis Risk Testing AHS – M2041

I2=0%), protein C deficiency (OR, 2.13; 95% CI, 1.16-3.90; I2=0%), and protein S deficiency (OR, 2.26; 95% CI, 1.34-3.80; I2=8.8%) (Chiasakul et al., 2019).” Antithrombin deficiency did not reach statistical significance in this study. Hence, in this review, inherited thrombophilias were found to be associated with an increased risk of arterial ischemic stroke in adults.

In a systematic review, Ortega studied the predictive value of D-dimer testing for venous thrombosis diagnosis in unusual locations. 3378 patients from 23 articles with thrombosis in unusual sites, such as upper extremity deep vein thrombosis (DVT), cerebral vein thrombosis (CVT) and splanchnic vein thrombosis (SVT), were studied. 12 articles on CVT concluded that timing of D-dimer testing is important and patients with short duration of symptoms displayed higher D-dimer levels. Sensitivity and specificity in these patients ranged from 58% to 97% and from 77% to 97.5%, respectively. The authors conclude that "D-dimer testing should not be currently recommended for the diagnosis of thrombosis in unusual sites as a first line diagnostic tool. The development of algorithms combining biomarkers such as D-dimer and clinical decision tools could improve the diagnosis" (Ordieres-Ortega et al., 2020).

Guidelines and Recommendations

American Heart Association/American Stroke Association

The AHA/ASA has issued a scientific statement for the management of stroke in neonates and children, wherein testing for thrombophilic abnormalities are discussed. The AHA/ASA admits that due to the lack of “an adequately powered study to detect the impact of genetic thrombophilia on recurrence risk in pediatric AIS [arterial ischemic stroke], definite recommendations about evaluation remain challenging”, but acknowledges that “laboratory testing outside of clinical studies may provide guidance for long-term management of the patient”. For cases of thrombophilia the AHA/ASA provides an algorithm for the “Targeted Evaluation of a Child With AIS for Rare Causes or Causes Requiring Additional Evaluation” that includes the examination of Factor VIII level Lipoprotein(a), *MTHFR* mutation, and homocysteine levels, and it is suggested that “non-DNA testing may need to be repeated when the child is older to ensure that adult levels of proteins have been attained” and “measurement of proteins or homocysteine levels in the acute phase of stroke may not be accurate and should be repeated after the acute event”. Finally, for the evaluation of a child with AIS, it is believed that “A thrombophilia evaluation is helpful in every case of childhood stroke, especially if there is no identifiable cause, medical history of thrombosis, or a first-degree relative with thrombosis history” (Ferriero et al., 2019).

In 2021, the AHA released guidelines on stroke prevention. The AHA brushes on testing for hematologic traits in the context of secondary stroke prevention. "If in certain clinical scenarios (eg, paradoxical emboli caused by venous thrombosis or recurrent venous thrombosis) testing for thrombophilic states is considered, testing for protein C, protein S, or anti thrombin levels should be deferred or repeated at least 4 to 6 weeks (or up to 6 months for factor VIII609) after the acute stroke given that these protein levels may be altered during the acute stroke phase” (Kleindorfer et al., 2021).

American College of Medical Genetics and Genomics (ACMG)

ACMG has released guidelines for laboratory testing of venous thromboembolism (VTE). This 2018 edition superseded the 2005 edition. The guidelines are as follows:

Testing for factor V Leiden and factor II c.*97G>A (this mutation is also known as G20210A) is recommended in the following circumstances:

- A first unprovoked VTE, especially <50 years old
- VTE at unusual sites (such as hepatic portal, mesenteric, and cerebral veins)

Venous and Arterial Thrombosis Risk Testing AHS – M2041

- Recurrent VTE
- Personal history of VTE with (a) two or more family members with a history of VTE or (b) one first-degree relative with VTE at a young age
- Patients with low activated protein C (APC) resistance activity

Testing may be considered in following circumstances:

- Females under the age of 50 who smoke tobacco and have a history of acute myocardial infarction
- Siblings of individuals known to be homozygous for factor V Leiden or factor II c.*97G>A, because they have a 1 in 4 chance of being a homozygote
- Asymptomatic pregnant female or female contemplating pregnancy, with a first-degree relative with unprovoked VTE or VTE provoked by pregnancy or contraceptive use
- Pregnant female or female contemplating pregnancy or estrogen use who has a first-degree relative with a history of VTE and is a known carrier for factor V Leiden and/or factor II c.97*G>A variant
- Pregnant female or female contemplating pregnancy with a previous non-estrogen-related VTE or VTE provoked by a minor risk factor, because knowledge of the factor V Leiden or factor II c.*97G>A status may alter pregnancy related thrombophylaxis (Zhang et al., 2018).

The ACMG found several clinical scenarios requiring special considerations worth mentioning, involving different populations. One involved the testing of asymptomatic versus symptomatic individuals, in which they assert that “It is generally not recommended to test asymptomatic minors as VTE rarely occurs before young adulthood even in the homozygous state.” For prenatal testing and population screening, the ACMG suggests that “prenatal testing and population screening are not indicated due to the low penetrance of these variants, later age of onset, and lack of genotype-directed prophylaxis”. Lastly, in women considering taking estrogen-containing oral contraceptives (OC) or hormone replacement therapy (HRT), the ACMG indicates that “A family and personal history of thrombosis should be carefully evaluated for all women before initiating HRT and a positive history may warrant thrombophilia screening” (Zhang et al., 2018). ACMG does not support testing for *MTHFR* variants in thrombophilia assessment due to the lack of correlation with negative pregnancy outcomes (Hickey, Curry, & Toriello, 2013). This statement was reaffirmed in 2020 (Bashford et al., 2020).

American Society of Hematology (ASH)

The 2013 ASH recommends against testing “for thrombophilia in adult patients with venous thromboembolism (VTE) occurring in the setting of major transient risk factors (surgery, trauma or prolonged immobility) (ASH, 2013).”

In 2018, ASH released their guidelines for management of venous thromboembolism, which included the following recommendations (Lim et al., 2018):

- “*Recommends* using a strategy starting with D-dimer for excluding PE in a population with low prevalence/PTP ($\leq 5\%$), followed by ventilation-perfusion (VQ) scan or computed tomography pulmonary angiography (CTPA) for patients requiring additional testing.
- *Recommends against* using a positive D-dimer alone to diagnose PE, and against additional testing following negative CTPA or normal VQ scan in a population with low prevalence/PTP ($\leq 5\%$).
- *Suggests* using a strategy starting with D-dimer for excluding PE in a population with intermediate prevalence/PTP ($\sim 20\%$), followed by VQ scan or CTPA for patients requiring additional testing.

Venous and Arterial Thrombosis Risk Testing AHS – M2041

- *Recommends against* using a positive D-dimer alone to diagnose PE, and against additional testing following negative CTPA or normal VQ scan in a population with intermediate prevalence/PTP (~20%).
- *Recommends against* using a positive D-dimer alone to diagnose PE, and against using D-dimer as a subsequent test following a negative CT scan in a population with high prevalence/PTP (≥50%).
- *Suggests* using a strategy starting with D-dimer for excluding recurrent PE in a population with unlikely PTP.
- *Recommends* using a strategy starting with D-dimer for excluding DVT in a population with low prevalence/PTP (≤10%), followed by proximal lower extremity ultrasound or whole-leg ultrasound for patients requiring additional testing.
- *Recommends against* using a positive D-dimer alone to diagnose DVT, and against additional testing following negative proximal or whole-leg ultrasound in a population with low prevalence/PTP (≤10%).
- *Recommends against* using a positive D-dimer alone to diagnose DVT in a population with intermediate prevalence/PTP (~25%).
- *Recommends against* using a positive D-dimer alone to diagnose DVT in a population with high prevalence/PTP (≥50%).
- *Suggests* using a strategy starting with D-dimer for excluding recurrent DVT in a population with unlikely PTP.
- *Suggests* a strategy starting with D-dimer for excluding upper extremity DVT in a population with low prevalence/unlikely PTP (10%), followed by duplex ultrasound if D-dimer is positive.
- *Recommends against* using a positive D-dimer alone to diagnose upper extremity DVT in a population with low prevalence/unlikely PTP (10%).
- *Suggests* a strategy of either D-dimer followed by duplex ultrasound/serial duplex ultrasound, or duplex ultrasound/serial duplex ultrasound alone for assessing patients suspected of having upper extremity DVT in a population with high prevalence/likely PTP (40%).
- *Recommends against* using a positive D-dimer alone to diagnose upper extremity DVT in a population with high prevalence/likely PTP (40%) (Lim et al., 2018).”

In 2020, The ASH guidelines released guidelines on management of venous thromboembolism. ASH suggest “against the routine use of prognostic scores, D-dimer testing, or venous ultrasound to guide the duration of anticoagulation” (Ortel et al., 2020).

Society for Vascular Medicine

This society recommends against workup for clotting disorders for patients with DVT as treatment will not change based on any abnormalities (SVM, 2013).

American College of Obstetricians and Gynecologists

The 2013 ACOG clinical management guidelines recommend that screening for inherited thrombophilia “may be considered in the following clinical settings:

1. “A personal history of venous thromboembolism that was associated with a non-recurrent risk factor
2. A first degree relative (parent or sibling) with a history of high-risk thrombophilia (ACOG, 2013).”

The 2018 ACOG Practice Bulletin Summary Number 197 supersedes the above 2013 guidelines (Practice Bulletin Number 138). In this update, the ACOG makes the following recommendations regarding screening based on “limited or inconsistent scientific evidence”:

Venous and Arterial Thrombosis Risk Testing AHS – M2041

“Screening for inherited thrombophilias is not recommended for women with a history of fetal loss or adverse pregnancy outcomes including abruption, preeclampsia, or fetal growth restriction because there is insufficient clinical evidence that antepartum prophylaxis with unfractionated heparin or low molecular-weight heparin prevents recurrence in these patients.”

“Because of the lack of association between either heterozygosity or homozygosity for the *MTHFR* C677T polymorphism and any negative pregnancy outcomes, including any increased risk of VTE, screening with either *MTHFR* mutation analyses or fasting homocysteine levels is not recommended.”

The 2018 ACOG recommends the following screening guideline based on “consensus and expert opinion”:

“Among women with personal histories of VTE, recommended screening tests for inherited thrombophilias should include factor V Leiden mutation; prothrombin G20210A mutation; and antithrombin, protein S, and protein C deficiencies.” (ACOG, 2018)

Evaluation of Genomic Applications in Practice and Prevention

“The Evaluation of Genomic Applications in Practice and Prevention (EGAPP) Working Group found adequate evidence to recommend against *routine* testing for Factor V Leiden (FVL) and/or prothrombin 20210G>A (PT) in the following circumstances: (1) adults with idiopathic venous thromboembolism (VTE). In such cases, longer term secondary prophylaxis to avoid recurrence offers similar benefits to patients with and without one or more of these mutations. (2) Asymptomatic adult family members of patients with VTE and an FVL or PT mutation, for the purpose of considering primary prophylactic anticoagulation. Potential benefits are unlikely to exceed potential harms. The evidence was insufficient to determine whether FVL/PT testing might have clinical utility in some circumstances, such as for identifying FVL homozygosity among asymptomatic family members of adults with idiopathic VTE or counseling patients about the risks and benefits of antithrombotic therapy. The recommendations do not extend to patients with other risk factors for thrombosis, such as contraceptive use, as the evidence review that serves as the basis for the recommendations focused primarily on idiopathic VTE (EGAPP, 2011).”

The Anticoagulation Forum

The Anticoagulation Forum published guidance in the Journal of Thrombosis and Thrombolysis on (Stevens et al., 2016):

- “Do not perform thrombophilia testing following an episode of provoked VTE. A positive thrombophilia evaluation is not a sufficient basis to offer extended anticoagulation following an episode of provoked VTE.
- Do not perform thrombophilia testing in patients following an episode of unprovoked VTE. If a patient with unprovoked VTE and low bleeding risk is planning to stop anticoagulation, test for thrombophilia if test results would change this decision. A negative thrombophilia evaluation is not a sufficient basis to stop anticoagulants following an episode of unprovoked VTE in a patient with low bleeding risk and willingness to continue therapy.

Heterozygosity for FVL or PGM does not increase the predicted risk of recurrence after unprovoked VTE to a clinically significant degree.

- Do not test for thrombophilia in asymptomatic family members of patients with VTE or hereditary thrombophilia. As a family history of VTE confers an excess risk of

Venous and Arterial Thrombosis Risk Testing AHS – M2041

thrombosis, relatives should be counseled regarding use of prophylaxis in high risk situations.

- Do not test for thrombophilia in asymptomatic family members of patients with VTE or hereditary thrombophilia who are contemplating use of estrogen. If a woman contemplating estrogen use has a first-degree relative with VTE and a known hereditary thrombophilia, test for that thrombophilia if the result would change the decision to use estrogen.
- Do not perform thrombophilia testing at the time of VTE diagnosis or during the initial 3-month course of anticoagulant therapy. When testing for thrombophilias following VTE, use either a 2-stage testing approach or perform testing after a minimum of 3 months of anticoagulant therapy has been completed, and anticoagulants have been held.
- Do not test for thrombophilia in asymptomatic family members of patients with VTE or hereditary thrombophilia who are contemplating pregnancy. If a woman contemplating pregnancy has a first-degree relative with VTE and a known hereditary thrombophilia... test for that thrombophilia if the result would change VTE prophylaxis decisions (Stevens et al., 2016)."

American College of Cardiology (ACC)

In 2017 guidance published in the New England Journal of medicine by Gupta was summarized by Barnes for the American College of Cardiology:

1. "Venous thromboembolism (VTE) affects an estimated 300,000-600,000 patients annually in the United States.
2. The risk of VTE recurrence is best predicted by whether the initial VTE episode was provoked or unprovoked, not the results of inherited thrombophilia testing.
3. Most patients with a provoked VTE have recently undergone surgery, immobility, trauma, or have a concurrent cancer diagnosis. Concurrent use of hormones (e.g., estrogen-containing contraceptive pills) is also frequently considered a provoking factor for VTE development.
4. For patients with a first provoked VTE event, guidelines recommend anticoagulation for only 3 months (not longer). Prolonged anticoagulation is associated with an increased risk of bleeding that outweighs the risk of VTE recurrence for these patients.
5. Patients with an unprovoked VTE (none of the provoking risk factors listed above) require longer anticoagulation due to a higher risk of recurrence that outweighs the risk of bleeding associated with long-term anticoagulation therapy.
6. Thrombophilia testing performed in the setting of an acute clot or ongoing anticoagulation therapy will often result in spurious results (usually false positive). For example, natural anticoagulants (e.g., protein C and S, antithrombin) are consumed during an acute thrombotic event and the levels can be reduced by ongoing anticoagulant therapy.
7. A recent study identified that up to 55% of Medicare patients with provoked VTE had undergone inappropriate thrombophilia testing, associated with significant cost to the healthcare system.
8. While thrombophilia testing rarely impacts management decisions about anticoagulation therapy, it may be beneficial for genetic testing purposes in patients presenting with a first unprovoked VTE at a young age (e.g., <45 years) or at an unusual site.
9. For patients with unprovoked VTE at a young age, VTE at an unusual site, arterial thrombosis, or pregnancy morbidity, testing for antiphospholipid antibodies, JAK2 mutation, and paroxysmal nocturnal hemoglobinuria may be beneficial.

Venous and Arterial Thrombosis Risk Testing AHS – M2041

10. There is no role for extensive cancer screening (e.g., computed tomography scanning) in patients with VTE. Only routine, age-appropriate cancer screening is recommended (G. Barnes, 2017; Gupta et al., 2017).”

Again in 2017, key points—inclusive of guiding points—published in the New England Journal of Medicine by Connors were captured by Barnes for the American College of Cardiology:

1. “The majority of patients with venous thromboembolism (VTE) should not be tested for thrombophilia. Data supporting clinical usefulness and benefits are limited or nonexistent.
2. Most patients with inherited thrombophilia can be identified by coagulation experts based on the patient’s personal and family history of VTE. Thrombophilia testing is usually not required.
3. Factors associated with an inherited thrombophilia include VTE at a young age (<40-50 years), a strong family history of VTE, VTE in conjunction with weak provoking factors at a young age, recurrent VTE, and VTE in an unusual site (e.g., cerebral or splanchnic veins).
4. Do not perform thrombophilia testing at the time of a VTE event, as it can be inaccurate (often false positive). Perform testing (when indicated) after completion of initial therapy and if it might change management strategies.
5. Do not perform thrombophilia testing while a patient is receiving anticoagulation. Instead, wait until 2 weeks after discontinuing warfarin, or 2 days for direct oral anticoagulants and heparin.
6. The goal of thrombophilia testing should be to aid decision making regarding future VTE prophylaxis, to guide testing of family members, and to determine the cause in severe or fatal VTE. Test results alone should not be used to decide on the duration of anticoagulation therapy.
7. Most VTE recurrence risk tools do not incorporate thrombophilia test results into their risk stratification schemes.
8. For patients with provoked VTE, even if they have homozygous factor V Leiden, prothrombin gene mutations, or deficiencies of protein S, C, or antithrombin, they do not require lifelong anticoagulation.
9. Currently available thrombophilia tests are insufficient to identify inherited risks of VTE. Therefore, a negative test should not be interpreted as a patient being free of thrombophilia.
10. Testing for the antiphospholipid antibody syndrome may be useful in patients with unprovoked VTE if there is clinical equipoise about extended anticoagulation courses. It can also be useful to determine warfarin versus direct oral anticoagulant therapy (G. D. Barnes, 2017; Connors, 2017).”

American Society for Clinical Pathology

ASCP has published guidelines with Choosing Wisely which state: “Do not test for Protein C, Protein S, or Antithrombin (ATIII) levels during an active clotting event to diagnose a hereditary deficiency because these tests are not analytically accurate during an active clotting event (ASCP, 2017).”

In 2019, an additional guideline was put forth by the ASCP on Choosing Wisely: “Do not perform a hypercoagulable workup in patients taking direct factor Xa or direct thrombin inhibitors.” The

Venous and Arterial Thrombosis Risk Testing AHS – M2041

guideline explained that the use of certain direct oral anticoagulants may render the results of such workups uninterpretable and inaccurate (ASCP, 2019).

European Society of Cardiology (ESC)

The ESC has published guidelines for the diagnosis and management of acute PE. These guidelines state:

- “D-dimer measurement and clinical prediction rules should be considered to rule out PE during pregnancy or the post-partum period
- Plasma D-dimer measurement, preferably using a highly sensitive assay, is recommended in outpatients/emergency department patients with low or intermediate clinical probability, or those that are PE-unlikely, to reduce the need for unnecessary imaging and irradiation
- A D-dimer test, using an age-adjusted cut-off or adapted to clinical probability, should be considered as an alternative to the fixed cut-off level
- D-dimer measurement is not recommended in patients with high clinical probability, as a normal result does not safely exclude PE, even when using a highly sensitive assay
- Assessment of the RV [right ventricle] by imaging methods or laboratory biomarkers should be considered, even in the presence of a low PESI [Pulmonary Embolism Severity Index] or a negative sPESI [simplified Pulmonary Embolism Severity Index] (Konstantinides et al., 2019).”

In 2021, the ESC Working Group released guidelines on diagnosis and management of acute deep vein thrombosis. These guidelines suggest that “ELISA D-dimer or highly sensitive immunoturbidimetric tests should be measured in ‘unlikely’ clinical probability patients to exclude DVT diagnosis” (Mazzolai et al., 2021).

Scottish Intercollegiate Guidelines Network

Regarding laboratory tests in the assessment of thrombosis risk, SIGN has stated that “Routine laboratory screening for heritable thrombophilias is not recommended (SIGN, 2014).”

American Society for Clinical Laboratory Science (ASCLS)

In 2021, ASCLS published guidelines on Choosing Wisely to suggest against ordering a homocysteine assay as part of the thrombophilia work up. “An elevated homocysteine level is not a clotting disorder and should not be included in thrombophilia testing panels” (ASCLS, 2021).

Applicable Federal Regulations

Many labs have developed specific tests that they must validate and perform in house. These laboratory-developed tests (LDTs) are regulated by the Centers for Medicare and Medicaid (CMS) as high-complexity tests under the Clinical Laboratory Improvement Amendments of 1988 (CLIA '88). LDTs are not approved or cleared by the U. S. Food and Drug Administration; however, FDA clearance or approval is not currently required for clinical use.

Billing/Coding/Physician Documentation Information

Venous and Arterial Thrombosis Risk Testing AHS – M2041

This policy may apply to the following codes. Inclusion of a code in this section does not guarantee that it will be reimbursed. For further information on reimbursement guidelines, please see Administrative Policies on the Blue Cross Blue Shield of North Carolina web site at www.bcbsnc.com. They are listed in the Category Search on the Medical Policy search page.

Applicable service codes: 81240, 81241, 81291, 81400, 81479, 85300, 85301, 85302, 85303, 85305, 85306, 85307, 0278U

BCBSNC may request medical records for determination of medical necessity. When medical records are requested, letters of support and/or explanation are often useful but are not sufficient documentation unless all specific information needed to make a medical necessity determination is included.

Scientific Background and Reference Sources

For policy titled: Venous Thrombosis Risk Testing AHS – M2041

ACOG. (2013). ACOG Practice Bulletin No. 138: Inherited thrombophilias in pregnancy. *Obstet Gynecol*, 122(3), 706-717. doi:10.1097/01.AOG.0000433981.36184.4e

ASH. (2013). ASH - Testing for thromboembolism | Choosing Wisely. <http://www.choosingwisely.org/clinician-lists/american-society-hematology-testing-for-thrombophilia-in-adults/>

Barnes, G. (2017, 06/05/2017). Thrombophilia Testing for Provoked VTE. Retrieved from <https://www.acc.org/latest-in-cardiology/ten-points-to-remember/2017/06/05/12/46/thrombophilia-testing-in-provoked-venous-thromboembolism>

Bartholomew, J. R. (2017). Update on the management of venous thromboembolism. *Cleve Clin J Med*, 84(12 Suppl 3), 39-46. doi:10.3949/ccjm.84.s3.04

Bauer, K. (2017). Protein S deficiency. Retrieved from https://www.uptodate.com/contents/protein-s-deficiency?topicRef=1361&source=see_link

Bauer, K. (2018a). Factor V Leiden and activated protein C resistance. Retrieved from https://www.uptodate.com/contents/factor-v-leiden-and-activated-protein-c-resistance?topicRef=1361&source=see_link

Bauer, K. (2018b). Protein C deficiency. Retrieved from https://www.uptodate.com/contents/protein-c-deficiency?topicRef=1361&source=see_link

Bauer, K. (2018c). Prothrombin G20210A mutation. Retrieved from https://www.uptodate.com/contents/prothrombin-g20210a-mutation?topicRef=1361&source=see_link

Bauer, K., & Lip, G. (2018). Overview of the causes of venous thrombosis - UpToDate. In G. Finlay (Ed.), UpToDate. Retrieved from https://www.uptodate.com/contents/overview-of-the-causes-of-venous-thrombosis?search=thrombophilia&usage_type=default&source=search_result&selectedTitle=1~150&display_rank=1.

Byrnes, J. R., & Wolberg, A. S. (2017). Red blood cells in thrombosis. *Blood*, 130(16), 1795-1799. doi:10.1182/blood-2017-03-745349

Connors, J. M. (2017). Thrombophilia Testing and Venous Thrombosis. In *N Engl J Med* (Vol. 377, pp. 2298). United States.

Venous and Arterial Thrombosis Risk Testing AHS – M2041

Crous-Bou, M., Harrington, L. B., & Kabrhel, C. (2016). Environmental and genetic risk factors associated with venous thromboembolism. *Semin Thromb Hemost*, 42(8), 808-820. doi:10.1055/s-0036-1592333

EGAPP. (2011). Recommendations from the EGAPP Working Group: routine testing for Factor V Leiden (R506Q) and prothrombin (20210G>A) mutations in adults with a history of idiopathic venous thromboembolism and their adult family members. *Genet Med*, 13(1), 67-76. doi:10.1097/GIM.0b013e3181f8e46f

Gupta, A., Sarode, R., & Nagalla, S. (2017). Thrombophilia Testing in Provoked Venous Thromboembolism: A Teachable Moment. *JAMA Intern Med*, 177(8), 1195-1196. doi:10.1001/jamainternmed.2017.1815

Herrmann, J. (2018). *Clinical Cardio-Oncology*: Elsevier.

Hickey, S. E., Curry, C. J., & Toriello, H. V. (2013). ACMG Practice Guideline: lack of evidence for MTHFR polymorphism testing. *Genet Med*, 15(2), 153-156. doi:10.1038/gim.2012.165

Kearon, C., Bauer, Kenneth. (2018). Clinical presentation and diagnosis of the nonpregnant adult with suspected deep vein thrombosis of the lower extremity. Retrieved from https://www.uptodate.com/contents/clinical-presentation-and-diagnosis-of-the-nonpregnant-adult-with-suspected-deep-vein-thrombosis-of-the-lower-extremity?search=deep%20vein%20thrombosis&source=search_result&selectedTitle=2~150&usage_type=default&display_rank=2

Lee, E. J., Dykas, D. J., Leavitt, A. D., Camire, R. M., Ebberink, E., García de Frutos, P., . . . Lee, A. I. (2017). Whole-exome sequencing in evaluation of patients with venous thromboembolism. *Blood Adv*, 1(16), 1224-1237. doi:10.1182/bloodadvances.2017005249

Lim, W., Le Gal, G., Bates, S. M., Righini, M., Haramati, L. B., Lang, E., . . . Mustafa, R. A. (2018). American Society of Hematology 2018 guidelines for management of venous thromboembolism: diagnosis of venous thromboembolism. *Blood Adv*, 2(22), 3226. doi:10.1182/bloodadvances.2018024828

Segal, J. B., Brotman, D. J., Emadi, A., Necochea, A. J., Samal, L., Wilson, L. M., . . . Bass, E. B. (2009). Outcomes of genetic testing in adults with a history of venous thromboembolism. *Evid Rep Technol Assess (Full Rep)*(180), 1-162.

Stevens, S. M., Woller, S. C., Bauer, K. A., Kasthuri, R., Cushman, M., Streiff, M., . . . Douketis, J. D. (2016). Guidance for the evaluation and treatment of hereditary and acquired thrombophilia. *J Thromb Thrombolysis*, 41, 154-164. doi:10.1007/s11239-015-1316-1

SVM. (2013, 02/21/2013). Don't do work up for clotting disorder (order hypercoagulable testing) for patients who develop first episode of deep vein thrombosis (DVT) in the setting of a known cause. Retrieved from <http://www.choosingwisely.org/clinician-lists/society-vascular-medicine-clotting-disorder-workup-after-first-episode-of-deep-vein-thrombosis/>

Thompson, B. T., Kabrhel, Christopher. (2018). Overview of acute pulmonary embolism in adults. Retrieved from https://www.uptodate.com/contents/overview-of-acute-pulmonary-embolism-in-adults?search=pulmonary%20embolism&source=search_result&selectedTitle=1~150&usage_type=default&display_rank=1

Zhang, S., Taylor, A. K., Huang, X., Luo, B., Spector, E. B., Fang, P., & Richards, C. S. (2018). Venous thromboembolism laboratory testing (factor V Leiden and factor II c.*97G>A), 2018 update: a technical standard of the American College of Medical Genetics and Genomics (ACMG). *Genet Med*, 20(12), 1489-1498. doi:10.1038/s41436-018-0322-

Venous and Arterial Thrombosis Risk Testing AHS – M2041

For policy titled: Venous and Arterial Thrombosis Risk Testing AHS – M2041

- ACOG. (2013). ACOG Practice Bulletin No. 138: Inherited thrombophilias in pregnancy. *Obstet Gynecol*, 122(3), 706-717. doi:10.1097/01.AOG.0000433981.36184.4e
- ACOG. (2018). ACOG Practice Bulletin No. 197 Summary: Inherited Thrombophilias in Pregnancy. *Obstet Gynecol*, 132(1), 249-251. doi:10.1097/aog.0000000000002705
- Algahtani, F. H., & Stuckey, R. (2019). High factor VIII levels and arterial thrombosis: illustrative case and literature review. *Ther Adv Hematol*, 10, 2040620719886685. doi:10.1177/2040620719886685
- ASCLS. (2021). American Society for Clinical Laboratory Science. Retrieved from <https://www.choosingwisely.org/clinician-lists/ascls7-do-not-order-a-homocysteine-assay-as-part-of-the-thrombophilia-work-up/>
- ASCP. (2017). American Society for Clinical Pathology. Retrieved from <http://www.choosingwisely.org/clinician-lists/ascp-testing-for-protein-c-protein-s-or-antithrombin-during-active-clotting-event/>
- ASCP. (2019). American Society of Clinical Pathology. Retrieved from <https://www.choosingwisely.org/clinician-lists/ascp-hypercoagulable-workup/>
- ASH. (2013). ASH - Testing for thromboembolism | Choosing Wisely. Retrieved from <http://www.choosingwisely.org/clinician-lists/american-society-hematology-testing-for-thrombophilia-in-adults/>. <http://www.choosingwisely.org/clinician-lists/american-society-hematology-testing-for-thrombophilia-in-adults/>
- Bank, I., Libourel, E. J., Middeldorp, S., Hamulyak, K., van Pampus, E. C., Koopman, M. M., . . . Buller, H. R. (2005). Elevated levels of FVIII:C within families are associated with an increased risk for venous and arterial thrombosis. *J Thromb Haemost*, 3(1), 79-84. doi:10.1111/j.1538-7836.2004.01033.x
- Barnes, G. (2017, 06/05/2017). Thrombophilia Testing for Provoked VTE. Retrieved from <https://www.acc.org/latest-in-cardiology/ten-points-to-remember/2017/06/05/12/46/thrombophilia-testing-in-provoked-venous-thromboembolism>
- Barnes, G. D. (2017). Thrombophilia Testing and Venous Thrombosis. Retrieved from <https://www.acc.org/latest-in-cardiology/ten-points-to-remember/2017/10/20/11/18/thrombophilia-testing-and-venous-thrombosis>.
- Bartholomew, J. R. (2017). Update on the management of venous thromboembolism. *Cleve Clin J Med*, 84(12 Suppl 3), 39-46. doi:10.3949/ccjm.84.s3.04
- Bashford, M. T., Hickey, S. E., Curry, C. J., Toriello, H. V., American College of Medical, G., Genomics Professional, P., & Guidelines, C. (2020). Addendum: ACMG Practice Guideline: lack of evidence for MTHFR polymorphism testing. *Genetics in Medicine*, 22(12), 2125-2125. doi:10.1038/s41436-020-0843-0
- Bauer, K. (2019a). Factor V Leiden and activated protein C resistance. Retrieved from https://www.uptodate.com/contents/factor-v-leiden-and-activated-protein-c-resistance?topicRef=1361&source=see_link
- Bauer, K. (2019b). Protein C deficiency. Retrieved from https://www.uptodate.com/contents/protein-c-deficiency?topicRef=1361&source=see_link
- Bauer, K. (2019c). Prothrombin G20210A mutation. Retrieved from https://www.uptodate.com/contents/prothrombin-g20210a-mutation?topicRef=1361&source=see_link
- Bauer, K. (2020). Clinical presentation and diagnosis of the nonpregnant adult with suspected deep vein thrombosis of the lower extremity. Retrieved from https://www.uptodate.com/contents/clinical-presentation-and-diagnosis-of-the-nonpregnant-adult-with-suspected-deep-vein-thrombosis-of-the-lower-extremity?search=deep%20vein%20thrombosis&source=search_result&selectedTitle=2~150&usage_type=default&display_rank=2
- Bauer, K. (2021). Protein S deficiency. Retrieved from https://www.uptodate.com/contents/protein-s-deficiency?topicRef=1361&source=see_link
- Bauer, K., & Lip, G. (2018). Overview of the causes of venous thrombosis - UpToDate. In G. Finlay (Ed.), *UpToDate*. Retrieved from <https://www.uptodate.com/contents/overview-of-the-causes-of-venous->

Venous and Arterial Thrombosis Risk Testing AHS – M2041

[thrombosis?search=thrombophilia&usage_type=default&source=search_result&selectedTitle=1~150&display_rank=1](https://www.uptodate.com/contents/overview-of-the-causes-of-venous-thrombosis?search=thrombophilia&usage_type=default&source=search_result&selectedTitle=1~150&display_rank=1)

Bauer, K., & Lip, G. (2020). Overview of the causes of venous thrombosis - UpToDate. In G. Finlay (Ed.), *UpToDate*. Retrieved from [https://www.uptodate.com/contents/overview-of-the-causes-of-venous-](https://www.uptodate.com/contents/overview-of-the-causes-of-venous-thrombosis?search=thrombophilia&usage_type=default&source=search_result&selectedTitle=1~150&display_rank=1)

[thrombosis?search=thrombophilia&usage_type=default&source=search_result&selectedTitle=1~150&display_rank=1](https://www.uptodate.com/contents/overview-of-the-causes-of-venous-thrombosis?search=thrombophilia&usage_type=default&source=search_result&selectedTitle=1~150&display_rank=1)

Byrnes, J. R., & Wolberg, A. S. (2017). Red blood cells in thrombosis. *Blood*, *130*(16), 1795-1799. doi:10.1182/blood-2017-03-745349

Carroll, B. J., & Piazza, G. (2018). Hypercoagulable states in arterial and venous thrombosis: When, how, and who to test? *Vasc Med*, *23*(4), 388-399. doi:10.1177/1358863x18755927

Chiasakul, T., De Jesus, E., Tong, J., Chen, Y., Crowther, M., Garcia, D., . . . Cuker, A. (2019). Inherited Thrombophilia and the Risk of Arterial Ischemic Stroke: A Systematic Review and Meta-Analysis. *J Am Heart Assoc*, *8*(19), e012877. doi:10.1161/jaha.119.012877

Connors, J. M. (2017). Thrombophilia Testing and Venous Thrombosis. In *N Engl J Med* (Vol. 377, pp. 2298). United States.

Crous-Bou, M., Harrington, L. B., & Kabrhel, C. (2016). Environmental and genetic risk factors associated with venous thromboembolism. *Semin Thromb Hemost*, *42*(8), 808-820. doi:10.1055/s-0036-1592333

Curtis, C., Mineyko, A., Massicotte, P., Leaker, M., Jiang, X. Y., Floer, A., & Kirton, A. (2017). Thrombophilia risk is not increased in children after perinatal stroke. *Blood*, *129*(20), 2793-2800. doi:10.1182/blood-2016-11-750893

EGAPP. (2011). Recommendations from the EGAPP Working Group: routine testing for Factor V Leiden (R506Q) and prothrombin (20210G>A) mutations in adults with a history of idiopathic venous thromboembolism and their adult family members. *Genet Med*, *13*(1), 67-76.

doi:10.1097/GIM.0b013e3181f8e46f

Ferriero, D. M., Fullerton, H. J., Bernard, T. J., Billingham, L., . . . Nursing, A. H. A. S. C. a. C. o. C. a. S. (2019). Management of Stroke in Neonates and Children: A Scientific Statement From the American Heart Association/American Stroke Association. *Stroke*, *50*(3). doi:

<http://dx.doi.org/10.1161/str.000000000000183>

Gupta, A., Sarode, R., & Nagalla, S. (2017). Thrombophilia Testing in Provoked Venous Thromboembolism: A Teachable Moment. *JAMA Internal Medicine*, *177*(8), 1195-1196. doi:10.1001/jamainternmed.2017.1815

Herrmann, J. (2018). *Clinical Cardio-Oncology*: Elsevier.

Hickey, S. E., Curry, C. J., & Toriello, H. V. (2013). ACMG Practice Guideline: lack of evidence for MTHFR polymorphism testing. *Genet Med*, *15*(2), 153-156. doi:10.1038/gim.2012.165

Kleindorfer, D. O., Towfighi, A., Chaturvedi, S., Cockcroft, K. M., Gutierrez, J., Lombardi-Hill, D., . . . Williams, L. S. (2021). 2021 Guideline for the Prevention of Stroke in Patients With Stroke and Transient Ischemic Attack: A Guideline From the American Heart Association/American Stroke Association. *Stroke*, *52*(7), e364-e467. doi:10.1161/str.0000000000000375

Konstantinides, S. V., Meyer, G., Becattini, C., Bueno, H., Geersing, G. J., Harjola, V. P., . . . Zamorano, J. L. (2019). 2019 ESC Guidelines for the diagnosis and management of acute pulmonary embolism developed in collaboration with the European Respiratory Society (ERS): The Task Force for the diagnosis and management of acute pulmonary embolism of the European Society of Cardiology (ESC). *Eur Respir J*, *54*(3). doi:10.1183/13993003.01647-2019

Kujovich, J. L. (2011). Factor V Leiden thrombophilia. *Genet Med*, *13*(1), 1-16. doi:10.1097/GIM.0b013e3181faa0f2

Kujovich, J. L. (2018). Factor V Leiden Thrombophilia. In M. P. Adam, H. H. Ardinger, R. A. Pagon, S. E. Wallace, L. J. H. Bean, K. Stephens, & A. Amemiya (Eds.), *GeneReviews((R))*. Seattle (WA): University of Washington, Seattle. GeneReviews is a registered trademark of the University of Washington, Seattle. All rights reserved.

Lee, E. J., Dykas, D. J., Leavitt, A. D., Camire, R. M., Ebberink, E., García de Frutos, P., . . . Lee, A. I. (2017). Whole-exome sequencing in evaluation of patients with venous thromboembolism. *Blood Adv*, *1*(16), 1224-1237. doi:10.1182/bloodadvances.2017005249

Venous and Arterial Thrombosis Risk Testing AHS – M2041

- Lehman, L. L., Beaute, J., Kapur, K., Danehy, A. R., Bernson-Leung, M. E., Malkin, H., . . . Trenor, C. C. (2017). Workup for Perinatal Stroke Does Not Predict Recurrence. *Stroke*, *48*(8), 2078-2083. doi:10.1161/STROKEAHA.117.017356
- Lim, W., Le Gal, G., Bates, S. M., Righini, M., Haramati, L. B., Lang, E., . . . Mustafa, R. A. (2018). American Society of Hematology 2018 guidelines for management of venous thromboembolism: diagnosis of venous thromboembolism. *Blood Adv*, *2*(22), 3226. doi:10.1182/bloodadvances.2018024828
- Linkins, L. A., & Takach Lapner, S. (2017). Review of D-dimer testing: Good, Bad, and Ugly. *Int J Lab Hematol*, *39 Suppl 1*, 98-103. doi:10.1111/ijlh.12665
- Mazzolai, L., Ageno, W., Alatri, A., Bauersachs, R., Becattini, C., Brodmann, M., . . . Aboyans, V. (2021). Second consensus document on diagnosis and management of acute deep vein thrombosis: updated document elaborated by the ESC Working Group on aorta and peripheral vascular diseases and the ESC Working Group on pulmonary circulation and right ventricular function. *European Journal of Preventive Cardiology*. doi:10.1093/eurjpc/zwab088
- Murphy, C. H., & Sabath, D. E. (2019). Comparison of Phenotypic Activated Protein C Resistance Testing With a Genetic Assay for Factor V Leiden. *Am J Clin Pathol*, *151*(3), 302-305. doi:10.1093/ajcp/aqy142
- NATF. (2019). Genetic Risk Factors for Blood Clots and the Role of Genetic Testing. Retrieved from <https://natfonline.org/2019/01/genetic-risk-factors-blood-clots-role-genetic-testing/>
- Onda, S., Furukawa, K., Haruki, K., Hamura, R., Shirai, Y., Yasuda, J., . . . Ikegami, T. (2021). d-dimer-based screening for early diagnosis of venous thromboembolism after hepatectomy. *Langenbeck's Archives of Surgery*, *406*(3), 883-892. doi:10.1007/s00423-020-02058-9
- Ordieres-Ortega, L., Demelo-Rodríguez, P., Galeano-Valle, F., Kremers, B. M. M., ten Cate-Hoek, A. J., & ten Cate, H. (2020). Predictive value of D-dimer testing for the diagnosis of venous thrombosis in unusual locations: A systematic review. *Thrombosis Research*, *189*, 5-12. doi:<https://doi.org/10.1016/j.thromres.2020.02.009>
- Ortel, T. L., Neumann, I., Ageno, W., Beyth, R., Clark, N. P., Cuker, A., . . . Zhang, Y. (2020). American Society of Hematology 2020 guidelines for management of venous thromboembolism: treatment of deep vein thrombosis and pulmonary embolism. *Blood Adv*, *4*(19), 4693-4738. doi:10.1182/bloodadvances.2020001830
- Previtali, E., Bucciarelli, P., Passamonti, S. M., & Martinelli, I. (2011). Risk factors for venous and arterial thrombosis. *Blood Transfus*, *9*(2), 120-138. doi:10.2450/2010.0066-10
- Raffini, L., Mahoney, D. H., & Armsby, C. (2019). Thrombophilia testing in children and adolescents. *UpToDate*. Retrieved from https://www.uptodate.com/contents/thrombophilia-testing-in-children-and-adolescents?topicRef=1354&source=see_link#H1628041305
- Segal, J. B., Brotman, D. J., Emadi, A., Necochea, A. J., Samal, L., Wilson, L. M., . . . Bass, E. B. (2009). Outcomes of genetic testing in adults with a history of venous thromboembolism. *Evid Rep Technol Assess (Full Rep)*(180), 1-162.
- SIGN. (2014). Prevention and management of venous thromboembolism Retrieved from <https://www.sign.ac.uk/assets/sign122.pdf>
- Stevens, S. M., Woller, S. C., Bauer, K. A., Kasthuri, R., Cushman, M., Streiff, M., . . . Douketis, J. D. (2016). Guidance for the evaluation and treatment of hereditary and acquired thrombophilia. *J Thromb Thrombolysis*, *41*, 154-164. doi:10.1007/s11239-015-1316-1
- SVM. (2013, 02/21/2013). Don't do work up for clotting disorder (order hypercoagulable testing) for patients who develop first episode of deep vein thrombosis (DVT) in the setting of a known cause. Retrieved from <http://www.choosingwisely.org/clinician-lists/society-vascular-medicine-clotting-disorder-workup-after-first-episode-of-deep-vein-thrombosis/>
- Thompson, B. T., Kabrhel, Christopher. (2018). Overview of acute pulmonary embolism in adults. Retrieved from https://www.uptodate.com/contents/overview-of-acute-pulmonary-embolism-in-adults?search=pulmonary%20embolism&source=search_result&selectedTitle=1~150&usage_type=default&display_rank=1
- Thompson, B. T., Kabrhel, Christopher. (2020). Overview of acute pulmonary embolism in adults. Retrieved from https://www.uptodate.com/contents/overview-of-acute-pulmonary-embolism-in-adults?search=pulmonary%20embolism&source=search_result&selectedTitle=1~150&usage_type=default&display_rank=1

Venous and Arterial Thrombosis Risk Testing AHS – M2041

Zhang, S., Taylor, A. K., Huang, X., Luo, B., Spector, E. B., Fang, P., & Richards, C. S. (2018). Venous thromboembolism laboratory testing (factor V Leiden and factor II c.*97G>A), 2018 update: a technical standard of the American College of Medical Genetics and Genomics (ACMG). *Genet Med*, 20(12), 1489-1498. doi:10.1038/s41436-018-0322-z

Specialty Matched Consultant Advisory Panel review 10/2020

Medical Director review 10/2020

Specialty Matched Consultant Advisory Panel review 10/2021

Medical Director review 10/2021

Policy Implementation/Update Information

For policy titled: Venous Thrombosis Risk Testing AHS – M2041

- 1/1/2019 New policy developed. BCBSNC will provide coverage for venous thrombosis risk testing when it is determined to be medically necessary because the criteria and guidelines have been met. Medical Director review 1/1/2019. (jd)
- 5/14/19 Reviewed by Avalon 1st Quarter 2019 CAB. Minor revision to Description section and “Related Policies” section added. When Not Covered section reformatted, no change to policy intent. Policy guidelines and references updated. Medical Director review 5/2019. (jd)
- 10/29/19 Wording in the Policy, When Covered, and/or Not Covered section(s) changed from Medical Necessity to Reimbursement language, where needed. (gm)
- 3/31/20 Specialty Matched Consultant Advisory Panel review 3/2020. Medical Director review 3/2020. (jd)

For policy titled: “Venous and Arterial Thrombosis Risk Testing AHS – M2041

- 5/12/20 Title changed from Venous Thrombosis Risk Testing to Venous and Arterial Thrombosis Risk Testing. Under When Not Covered section, changed items #2 and #3 from “investigational” to “not medically necessary; added items #4 and #5. Policy guidelines and references updated. Medical Director review 4/2020. (jd)
- 11/10/20 Specialty Matched Consultant Advisory Panel review 10/2020. Medical Director review 10/2020. (jd)
- 7/13/21 Reviewed by Avalon 1st Quarter 2021 CAB. Specified “deep” venous thrombosis throughout policy for clarity. The following changes made to the When Covered section: Item 1 – added item i.- “Pediatric arterial ischemic stroke”; item 2 – added statement “Assays for clotting inhibitors amount and function should be performed prior to any molecular testing.” And item k. “Pediatric arterial ischemic stroke”. The following changes were made to the When Not Covered section: Item 2 – added statement “Reimbursement is not allowed for venous thrombosis risk testing for superficial venous thrombosis (including superficial thrombophlebitis and varicosities).”; to item 3 – added h. “Testing more than once per lifetime”. Policy guidelines and references updated. Policy noticed 5/18/21; effective 7/13/21. Medical Director review 4/2021. (jd)
- 9/21/21 Title correction as follows: added the #1 to read M2041. (jd)
- 11/2/21 Specialty Matched Consultant Advisory Panel review 10/2021. Medical Director review 10/2021. (jd)

Venous and Arterial Thrombosis Risk Testing AHS – M2041

5/17/22 Reviewed by Avalon 1st Quarter 2022 CAB. Under the When Covered section: reworded item 1, adding “mutations” and removed criteria “Recurrent deep vein thrombosis” from both item 1 and 2. Added item 2 to When Not Covered section as follows: “Reimbursement is not allowed for genetic testing for Factor V Leiden and Prothrombin gene G20210A mutations in patients with recurrent thrombotic events who are receiving a lifelong anticoagulation regimen”. Policy guidelines and references updated. Added code 0278U to the Billing/Coding section. Medical Director review 4/2022. (jd)

Medical policy is not an authorization, certification, explanation of benefits or a contract. Benefits and eligibility are determined before medical guidelines and payment guidelines are applied. Benefits are determined by the group contract and subscriber certificate that is in effect at the time services are rendered. This document is solely provided for informational purposes only and is based on research of current medical literature and review of common medical practices in the treatment and diagnosis of disease. Medical practices and knowledge are constantly changing and BCBSNC reserves the right to review and revise its medical policies periodically.